


CONVENTION ON INTERNATIONAL TRADE IN ENDANGERED SPECIES
OF WILD FAUNA AND FLORA



Twenty-seventh meeting of the Animals Committee
Veracruz (Mexico), 28 April – 3 May 2014

Interpretation and implementation of the Convention

Species trade and conservation

Standard nomenclature [Resolution Conf. 12.11 (Rev. CoP16)]

REPORT OF THE SPECIALIST ON ZOOLOGICAL NOMENCLATURE

1. This document has been prepared by the specialist on zoological nomenclature of the Animals Committee¹.

Nomenclatural tasks referred to the Animals Committee by CoP16

2. *Hippocampus* taxonomy

At CoP16 Australia had asked for the recognition of a number of *Hippocampus* species. As this request had been made after Annex 6 (Rev.1) of CoP16 Doc 43.1 (Rev.1) had been adopted already it was decided to refer the discussion of this issue to the next Animal Committee meeting. The specialist on zoological nomenclature has contacted Australia to clarify the issue. Australia requests the following species to be recognized as valid species under CITES based on Kuitert, R.H. (2001): Revision of the Australian seahorses of the genus *Hippocampus* (Syngnathiformes: Syngnathidae) with a description of nine new species - Records of the Australian Museum, 53: 293-340.

Hippocampus bleekeri FOWLER, 1907 - split from *Hippocampus abdominalis* LESSON, 1827
Hippocampus dahli OGILBY, 1908 - split from *Hippocampus trimaculatus* LEACH, 1814
Hippocampus elongatus CASTELNAU, 1873 (to be reinstated for *H. subelongatus* CASTELNAU, 1873)
Hippocampus kampylotrachelos BLEEKER, 1854 - split from *Hippocampus trimaculatus* LEACH, 1814
Hippocampus planifrons PETERS, 1877 - split from *Hippocampus kuda* BLEEKER, 1852
Hippocampus taeniopterus BLEEKER, 1852 - split from *Hippocampus kuda* BLEEKER, 1852
Hippocampus tristis CASTELNAU, 1872 - split from *Hippocampus kuda* BLEEKER, 1852
Hippocampus tuberculatus CASTELNAU, 1875 - split from *Hippocampus breviceps* PETERS, 1869

H. bleekeri, *H. dahli*, *H. planifrons*, *H. tristis* and *H. tuberculatus* are endemic to Australian waters; *H. taeniopterus* is a western pacific species and *H. kampylotrachelos* occurs in South Indonesia (according to ESCHMEYER & FRICKE 2014 (Catalog of Fishes, see below).

Except for the reinstating of *H. elongatus* for *H. subelongatus*, all taxa are also currently accepted as valid taxa in the "Catalog of Fishes" by ESCHMEYER, W.N. & FRICKE, R. (eds.): Catalog of Fishes, an online reference (<http://research.calacademy.org/redirect?url=http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>), version downloaded 2014-02-17).

¹ The geographical designations employed in this document do not imply the expression of any opinion whatsoever on the part of the CITES Secretariat or the United Nations Environment Programme concerning the legal status of any country, territory, or area, or concerning the delimitation of its frontiers or boundaries. The responsibility for the contents of the document rests exclusively with its author.

If recommended for adoption the list of nomenclatural standard references for the genus *Hippocampus* would have to be changed as outlined in Annex 1 to this document. Extracts from the Catalog of Fishes for these species have been included for background information in Annex 1 as well.

3. Coral Nomenclature reference

Decision 15.64 (a)) requires the Animals Committee to "identify existing coral reference materials that could be adopted as standard nomenclatural references for CITES-listed corals". As it had not been possible to identify references serving this purpose in the term between CoP 15 and CoP 16 a coral species list provided by WCMC has been adopted at CoP 16 as an interim solution leaving the task still open.

Other Nomenclatural issues

4. Layout of nomenclatural references in Res. Conf. 12.11(rev. CP16)

During the last ten years the number of the various publications adopted to define the nomenclature used for different animal groups has increased tremendously. It has become quite difficult to identify which is the reference literature for separate group such as Cetacea, Primates, Iguanidae, Boidae, Elapidae or Trionychidae. Adopting new overall checklists for higher taxa such as birds or Testudines only solve this problem for short interim time sections. Annex 2 presents a different layout for the fauna references in the Annex of Res. Conf. 12.11 (Rev. CoP16) which is recommended by the nomenclature specialist in order to facilitate the use of this source of information in the future.

5. Nomenclature references for species suggested for inclusion in the CITES Appendices not covered by the nomenclature references listed in Res. Conf. 12.11

With new species suggested for inclusion in the CITES Appendices it happens often that the respective taxa are not covered by the references listed in Res. Conf. 12.11 (see new shark species and the genus *Hoplostodactylus* listed at CoP16 in Appendix II).

This problem will increase in the future as more and more taxonomic checklists covering higher taxa such as the classes Mammalia, Aves, Reptilia, Amphibia or the various fish classes (cover all species of the world and not only the CITES ones) will no longer be published as single publications but turned into online databases that are constantly updated. However, CITES requires as much stability as possible with regard to the nomenclature used under this convention. As a result extracts from these online databases downloaded at a given date such as already done for amphibian, fish and spider species will also increase to serve as taxonomic checklists for the nomenclature used under CITES. These naturally will only cover the CITES species listed so far.

Section C.1.4. of Annex 6 of Resolution 9.24 includes a recommendation with regard to the taxonomy and nomenclature of a taxon suggested for inclusion in the Appendices, however, this usually does not result in a formal request of the proponent Party to adopt a certain nomenclature standard reference in cases where the new taxon is not covered by already adopted standard references. It is therefore recommended to reword the existing text:

" If the species concerned is included in one of the standard lists of names or taxonomic references adopted by the Conference of the Parties, the name provided by that reference should be entered here. If the species concerned is not included in one of the adopted standard references, the proponent should provide references as to the source of the name used"

in a way that the proponent is definitely requested to formally suggest a nomenclature standard reference to be included in the Annex 6 of Resolution 12.11 for the proposed taxon in case it is missing so far. An option to be considered by the AC may be: "... the proponent is requested to provide taxonomic information as to the source of the name used and suggest a new nomenclature standard reference to be included in the Annex of Resolution 12.11 if necessary".

6. *Bradypus pygmaeus*

In the end of 2013 it had been noticed that animals known under the name of *Bradypus pygmaeus* were previously included in Appendix II by the Conference of the Parties under the name of *Bradypus variegatus* and that the omission of the species from the Appendices had been an oversight. Consequently the Secre-

ariat corrected CITES Appendices on the CITES website to include *Bradypus pygmaeus*. For details see CITES Notification 2013/052.

7. Taxonomic changes in non-passerines bird species between the current basic nomenclature reference for birds and its latest published edition

In 2013 a new edition of "The Howard and Moore Complete Checklist of the Birds of the World" has been published in a first volume covering the Non-Passerines. The EU Commission commissioned WCMC to identify the changes that would result with regard to the nomenclature of the bird species covered by CITES and the EU Regulation implementing CITES within the member states of the European Union. The report is presented in Annex 3.

8. *Nilssonia leithii* (Gray, 1872)

At CoP16 a publication by PRASCHAG & al. (2007)² has been adopted as new nomenclature standard reference for the species *Nilssonia gangeticus*, *N. hurum* and *N. nigricans*, formerly placed into the genus *Aspideretes*. This publication also places the former species *Aspideretes leithii* into the genus *Nilssonia* which has been overlooked when changing the nomenclature standard reference for the *Aspideretes* taxa at CoP16.

9. Taxonomic Checklist for all Chamaeleonidae species and the genus *Phelsuma*

As the nomenclature standard references for chameleon species as well as the species of the genus *Phelsuma* have increased significantly (with regard to chameleons there are currently altogether 32 single nomenclature standard references) and it has become difficult to keep track of them. Therefore the German Scientific Authority commissioned Frank GLAW to compile a taxonomic checklist for all species of the family Chamaeleonidae as well as the genus *Phelsuma*. It will include all species or taxonomic changes within these taxa published until July 2014. It is planned to publish the checklist in the first volume of Vertebrate Zoology in 2015. Until today 11 chameleon genera with together 198 species and 52 *Phelsuma* species have been identified to be covered by the checklist (see Annex 4). Like the main nomenclature reference for chelonians of the world the checklist will include all species and subspecies, list the most important synonyms and the distribution of the species concerned.

10. Other identified nomenclature changes in reptile species

The EU Commission commissioned WCMC as well to identify recent changes that would result with regard to the nomenclature of the reptile species (other than chameleons and *Phelsuma* species) covered by CITES and the EU Regulation implementing CITES within the member states of the European Union. The report is presented in Annex 5.

11. Amphibian species adopted by the parties but missing from current Taxonomic Checklist for Amphibians which has been adopted as nomenclature standard reference for amphibian species

At CoP16 quite a number of changes in the amphibian taxonomy and resulting nomenclature have been adopted. Unfortunately two species (*Altiphrynoides malcolmi* and *Hyloxalus azureiventris*) have been missed by error to be added to the extract from the online-database "Amphibian Species of the World" now used as Taxonomic Checklist for CITES listed Amphibian species (see Res.Conf.12.11). The missing information for these two species has been compiled in Annex 6 based on the current online version of the mentioned database and completed by adding as well the respective information for *Hynobius amjiensis* listed on Appendix III by China on June 12 2013.

12. *Sphyrna gilberti*, *Scleropages inscriptus* and necessary other additions to the Taxonomic Checklist of all CITES listed Shark and Fish species

Sphyrna gilberti QUATTRO, DRIGGERS III, GRADY, ULRICH & ROBERTS, 2013³, has been described 2013 as new species of the genus *Sphyrna*. The designation of this species is based on 54 specimens collected in the coastal waters of South Carolina, USA. Morphologically, *S. gilberti* is separable from *S. lewini* (GRIFFITH &

² PRASCHAG, P., HUNSDÖRFER, A.K., REZA, A.H.M.A. & FRITZ, U. (2007): Genetic evidence for wild-living *Aspideretes nigricans* and a molecular phylogeny of South Asian softshell turtles (Reptilia: Trionychidae: Aspideretes, Nilssonia). - *Zoologica Scripta*, **36**:301-310.

³ QUATTRO, J.M., DRIGGERS III, W.B., GRADY, J.M., ULRICH, G.F. & ROBERTS, M.A. (2013): *Sphyrna gilberti*, sp. nov., a new hammerhead shark (Carcharhiniformes, Sphyrnidae) from the western Atlantic Ocean. -- *Zootaxa*, **3702**(2): 159-178.

SMITH, 1834) only in the number of precaudal vertebrae. Due to the rarity of specimens and the highly migratory behaviour of most sphyrynids, the range of *S. gilberti* is still unknown.

Taking into account the fact that the distribution area of the few specimens caught of *S. gilberti* lies within the distribution area of *S. lewini*, that there is a resemblance between *S. lewini* and *S. gilberti* which only differ by internal morphological features which implies that many specimens identified so far as *S. lewini* may in fact belong to *S. gilberti*, the nomenclature specialist of the AC considers *S. gilberti* to be covered by the current listing of *S. lewini* in Appendix II of CITES.

In 2012 the CITES Secretariat informed the Parties by Notification 2012/43 of the publication of a new *Scleropages* species, *Scleropages inscriptus* ROBERTS, 2012, and its status with regard to the CITES Appendices. Taking into account the information provided in the original listing proposal for *Scleropages formosus* and the information provided in the publication describing the new species *Scleropages inscriptus*⁴ it seems obvious that *Scleropages inscriptus* has to be regarded as covered by the current listing of *Scleropages formosus* in Appendix I.

ROBERTS (2012) indicates its close relationship to *S. formosus* as well as the possibility to easily distinguish between these two taxa by different color patterns with *S. inscriptus* having unique maze-like markings : "It differs from the two Australian species of the subgenus *Scleropages* in meristic and morphometric characters, while at the same time agreeing closely in these characters with its Southeast Asian congener *S. formosus* of the subgenus *Delsmania*" (page 115) and "Thus the striking difference in color patterns in *S. inscriptus* strongly indicates that it is not the same species as *S. formosus*" (page 117). Therefore it is recommended to accept *S. inscriptus* as a valid species being part of the former listing of *S. formosus* and it is consequently recommended to add the species to Appendix I at the occasion of the next CoP.

At CoP 16 a number of shark species have been included in Appendix II of CITES. These are naturally missing in the checklist for fishes (CoP16 Doc 43.1 Annex 2) adopted as major nomenclature standard reference for fish species. In addition, *Neoceratodus forsteri* as well as the genus *Laterimeria* have also been missing from the checklist so far.

Annex 7 comprises the missing taxonomic information extracted from the online database "Catalogue of Fish Species" by ESCHMEYER & FRICKE for all the mentioned fish species.

Harmonization of nomenclature with other biodiversity-related multilateral environmental agreements

This section has been prepared by the CITES Secretariat.

13. Resolution Conf. 12.11 (Rev. CoP16) on Standard nomenclature acknowledges the desirability of harmonizing, to the extent possible, the species nomenclature used by the biodiversity-related multilateral environmental agreements and directs the Secretariat, in close cooperation with the nomenclature specialists of the Animals and Plants Committees to promote such harmonization. This objective was supported by the Chairs of the Scientific Advisory Bodies of Biodiversity-related Conventions (CSAB) at its 2nd meeting (Bonn, May 2008).
14. The principal species-based multilateral environmental agreements is the Convention on the Conservation of Migratory Species of Wild Animals (CMS). In recent years, through a process of mutual adjustment CITES and CMS have achieved almost complete harmonization of the nomenclature used for the Class Mammalia.
15. As reported at the 26th meeting of the Animals Committee in document AC26 Doc. 20 the 10th meeting of the Conference of the Parties to CMS (Bergen, November 2011) requested the Chair of the CMS Scientific Council to liaise with CSAB and others with the aim of evaluating the possible adoption by CMS Parties of a single nomenclature and taxonomy for birds for adoption at CMS CoP11 in late 2014.
16. In fulfilment of this instruction CMS convened an *ad hoc* meeting on Harmonization of bird taxonomy held in Formia (Italy) on 8 October 2013. The Secretariat participated at this meeting. At the time of writing (February 2014) the final report of the meeting is not available, but it is expected to be so before the present meeting.

⁴ ROBERTS, T.R. (2012): *Scleropages inscriptus*, a new fish species from the Tanathayi or Tenasserim River basin, Malay Peninsula of Myanmar (Osteoglossidae: Osteoglossiformes). -- *aqua, International Journal of Ichthyology*, vol. 18 (2): 113-118.

17. At the meeting, the tendency amongst CMS and its daughter Agreements that were present was that they had an interest in following the nomenclature used in the IUCN Red List of Threatened Species because many existing CMS instruments had direct links to it. With respect to birds, the IUCN Red List of Threatened Species uses *The BirdLife checklist of the birds of the world, with conservation status and taxonomic sources* which is maintained by a BirdLife Taxonomic Working Group.
18. The current version of *The BirdLife checklist of the birds of the world, with conservation status and taxonomic sources* is Version 6 of November 2013 (http://www.birdlife.org/datazone/userfiles/file/Species/Taxonomy/BirdLife_Checklist_Version_6.zip)
19. BirdLife International intends to publish its checklist, with a volume on non-passerines due in August 2014 and a volume on passerines to be published in 2016. (<http://www.lynxeds.com/product/hbw-and-birdlife-international-illustrated-checklist-birds-world>). BirdLife International have kindly compared the current CITES Appendices with the present draft of the list of species which will be found in the non-passerine checklist to be published later in 2014. The differences can be seen in Annex 8 of the present document.

Recommendations for the work of the Nomenclature Working Group at AC 26

20. It is suggested
 - to develop recommendations on all nomenclatural changes identified in this paper under points 2, 7-8, 10-12
 - to develop recommendations with regard to the suggestions of the nomenclature specialist under points 4 and 5
 - to develop a recommendation on how to proceed with the request of the Parties outlined in Decision 15.64(a) under point 3
 - to take account of the developments reported in points 13-19 and respond as necessary.

1. List of standard reference for the nomenclature of *Hippocampus* species in case the request of Australia will be recommended

LOURIE, S. A., VINCENT, A. C. J. & HALL, H. J. (1999): *Seahorses. An identification guide to the world's species and their conservation*. Project Seahorse (ISBN 0 9534693 0 1) (Second edition available on CD-ROM). [for *Hippocampus* with the exception of the species mentioned below]

FOSTER, R. & GOMON, M. F. (2010): **A new seahorse (Teleostei: Syngnathidae: *Hippocampus*) from south-western Australia. – Zootaxa, 2613: 61-68. [for *Hippocampus paradoxus*]**

GOMON, M. F. & KUITER, R. H. (2009): Two new pygmy seahorses (Teleostei: Syngnathidae: *Hippocampus*) from the Indo-West Pacific. – *Aqua, Int. J. of Ichthyology*, 15(1): 37-44. [for *Hippocampus debelius*, *Hippocampus waleanus*]

HORNE, M. L. (2001): [A new seahorse species \(Syngnathidae: *Hippocampus*\) from the Great Barrier Reef](#) – *Records of the Australian Museum*, 53: 243-246. [for *Hippocampus queenslandicus*]

KUITER, R. H. (2001): [Revision of the Australian seahorses of the genus *Hippocampus* \(Syngnathiformes: Syngnathidae\) with a description of nine new species](#) – *Records of the Australian Museum*, 53: 293-340. [for *Hippocampus alatus*, *H. biocellatus*, *H. bleekeri*, *H. dahli*, *H. elongatus*⁵, *H. grandiceps*, *H. hendriki*, *H. jugumus*, *H. kamylotrachelos*, *H. montebelloensis*, *H. multispinus*, *H. planifrons*, *H. procerus*, *H. semispinosus*, *H. taeniopterus*, *H. tristis*, *H. tuberculatus*]

KUITER, R. H. (2003): [A new pygmy seahorse \(Pisces: Syngnathidae: *Hippocampus*\) from Lord Howe Island](#) – *Records of the Australian Museum*, 55: 113-116. [for *Hippocampus colemani*]

LOURIE, S. A. & KUITER, R. H. (2008): Three new pygmy seahorse species from Indonesia (Teleostei: Syngnathidae: *Hippocampus*). – *Zootaxa*, 1963: 54-68. [for *Hippocampus pontohi*, *Hippocampus satomiae*, *Hippocampus severnsi*]

LOURIE, S. A. & RANDALL, J. E. (2003): A new pygmy seahorse, *Hippocampus denise* (Teleostei: Syngnathidae), from the Indo-Pacific – *Zoological Studies*, 42: 284-291. [for *Hippocampus denise*]

PIACENTINO, G. L. M. AND LUZZATTO, D. C. (2004): *Hippocampus patagonicus* sp. nov., new seahorse from Argentina (Pisces, Syngnathiformes). – *Revista del Museo Argentino de Ciencias Naturales*, 6(2): 339-349. [for *Hippocampus patagonicus*]

RANDALL, J. & LOURIE, S. A. (2009): *Hippocampus tyro*, a new seahorse (Gasterosteiformes: Syngnathidae) from the Seychelles. – *Smithiana Bulletin*, 10: 19-21. [for *Hippocampus tyro*]

⁵ Reinstating *Hippocampus elongatus* for *Hippocampus subelongatus*.

2. Extract for the Australian species from

"Catalogue of Fishes "

by ESCHMEYER, W.N. & FRICKE, R. (eds.): **Catalog of Fishes, an online reference**

(<http://research.calacademy.org/redirect?url=http://researcharchive.calacademy.org/research/Ichthyology/catalog/fishcatmain.asp>), version downloaded 2014-02-17).

Hippocampus bleekeri FOWLER, 1907

agnesae, *Hippocampus* Fowler [H. W.] 1907:429, Fig. 5 [Proceedings of the Academy of Natural Sciences of Philadelphia v. 59; ref. [1377](#)] Victoria, Australia. Holotype: ANSP 33123. Paratypes: ANSP 33162 (1, in jar with holotype). Type catalog: Böhlke 1984:158 [ref. [13621](#)]. •Synonym of *Hippocampus abdominalis* Lesson 1827 -- (Paxton et al. 1989:421 [ref. [12442](#)], Gomon et al. 1994:448 [ref. [22532](#)], Lourie et al. 1999:69 [ref. [23993](#)]). •Synonym of *Hippocampus bleekeri* Fowler 1907 -- (Kuitert 2001:323 [ref. [25900](#)], Paxton et al. 2006:826 [ref. [29073](#)], Kuitert 2009:54 [ref. [30404](#)]). **Current status:** Synonym of *Hippocampus bleekeri* Fowler 1907. Syngnathidae: Hippocampinae. Habitat: marine.

bleekeri, *Hippocampus* Fowler [H. W.] 1907:426, Fig. 4 [Proceedings of the Academy of Natural Sciences of Philadelphia v. 59; ref. [1377](#)] Victoria, Australia. Holotype: ANSP 33122 (in 2 pieces). Paratypes: ANSP 33184-87 (6, 1 with head missing, in jar with holotype). Type catalog: Böhlke 1984:159 [ref. [13621](#)]. •Synonym of *Hippocampus abdominalis* Lesson 1827 -- (Paxton et al. 1989:421 [ref. [12442](#)], Gomon et al. 1994:448 [ref. [22532](#)], Lourie et al. 1999:69 [ref. [23993](#)]). •Valid as *Hippocampus bleekeri* Fowler 1907 -- (Kuitert 2001:323 [ref. [25900](#)], Paxton et al. 2006:826 [ref. [29073](#)] dated 1908, Kuitert 2008:454 [ref. [30642](#)], Kuitert 2009:54 [ref. [30404](#)]). **Current status:** Valid as *Hippocampus bleekeri* Fowler 1907. Syngnathidae: Hippocampinae. Distribution: Southeastern Australia: Victoria, Tasmania and South Australia. Habitat: marine.

graciliformis, *Hippocampus* McCulloch [A. R.] 1911:29, Pl. 6 (fig. 2) [Biological Results Endeavour [v. 1] (pt 1); ref. [2936](#)] Near Bass Strait, Australia. Holotype (unique): AMS E.429. •Synonym of *Hippocampus abdominalis* Lesson 1827 -- (Paxton et al. 1989:421 [ref. [12442](#)], Gomon et al. 1994:448 [ref. [22532](#)], Lourie et al. 1999:69 [ref. [23993](#)]). •Synonym of *Hippocampus bleekeri* Fowler 1907 -- (Kuitert 2001:323 [ref. [25900](#)], Paxton et al. 2006:826 [ref. [29073](#)], Kuitert 2009:18 [ref. [30404](#)]). **Current status:** Synonym of *Hippocampus bleekeri* Fowler 1907. Syngnathidae: Hippocampinae. Habitat: marine.

Hippocampus dahl OGILBY, 1908

dahl, *Hippocampus* Ogilby [J. D.] 1908:17 [Annals of the Queensland Museum No. 9 (pt 1); ref. [3285](#)] Moreton Bay, Noosa, southern Queensland, Australia. Holotype: QM I.788. Paratypes: QM (missing). •Synonym of *Hippocampus planifrons* Peters 1877. •Synonym of *Hippocampus trimaculatus* Leach 1814 -- (Lourie et al. 1999:125 [ref. [23993](#)]). •Valid as *Hippocampus dahl* Ogilby 1908 -- (Kuitert 2001:308 [ref. [25900](#)], Horne 2001:245 [ref. [25696](#)], Paxton et al. 2006:826 [ref. [29073](#)], Kuitert 2009:124 [ref. [30404](#)], Larson et al. 2013:77 [ref. [32988](#)]). **Current status:** Valid as *Hippocampus dahl* Ogilby 1908. Syngnathidae: Hippocampinae. Distribution: Northern and eastern Australia. Habitat: marine.

lenis, *Hippocampus* De Vis [C. W.] 1908 Not available, name only. From Kuitert 2001:308, 309 [ref. [25900](#)]; unpublished museum name, based on specimen label of QM I.788, which later became the holotype of

Hippocampus dahli Ogilby 1908. •In the synonymy of *Hippocampus dahli* Ogilby 1908 -- (Kuitert 2009:18 [ref. 30404]). **Current status:** Synonym of *Hippocampus dahli* Ogilby 1908. Syngnathidae: Hippocampinae.

***Hippocampus subelongatus* CASTELNAU, 1873**

elongatus, Hippocampus Castelnau [F. L.] 1873:144 [Proceedings of the Zoological and Acclimatisation Society of Victoria, Melbourne v. 2; ref. 758] Fremantle, Western Australia. Holotype (unique): MNHN A-4536. Kuitert's 2001:330 [ref. 25900] reasons for reversing the selection of *subelongatus* over *elongatus* by Lourie et al. 1999 is not supported by the Code. See Kuitert 2001:330 [ref. 25900] for discussion of type. •Synonym of *Hippocampus angustus* Günther 1870 -- (Paxton et al. 1989:421 [ref. 12442]). •Synonym of *Hippocampus subelongatus* Castelnau 1873 -- (Lourie et al. 1999:123 [ref. 23993], Paxton et al. 2006:828 [ref. 29073], Kuitert 2009:94 [ref. 30404]). •Wrongly valid as *Hippocampus elongatus* Castelnau 1873 -- (Kuitert 2001:330 [ref. 25900]). **Current status:** Synonym of *Hippocampus subelongatus* Castelnau 1873. Syngnathidae: Hippocampinae. Habitat: marine.

subelongatus, Hippocampus Castelnau [F. L.] 1873:145 [Proceedings of the Zoological and Acclimatisation Society of Victoria, Melbourne v. 2; ref. 758] Fremantle, Western Australia. Holotype (unique): MNHN A-4536. Type catalog: Bertin & Estève 1950:53 [ref. 19576]. None of the reasons cited by Kuitert 2001:330 [ref. 25900] to change the subsequent selection of Lourie et al. 1999, selecting *subelongatus* over *elongatus* are valid; Kuitert's reasons are not supported by the Code -- there is no such thing as page preference for example, or because *subelongatus* is a derivative of *elongatus*. Type follows discussion in Kuitert 2001:330 [ref. 25900]. •Valid as *Hippocampus subelongatus* Castelnau 1873 -- (Lourie et al. 1999:123 [ref. 23993], Paxton et al. 2006:828 [ref. 29073], Kuitert 2009:94 [ref. 30404]). •Wrongly as a synonym of *Hippocampus elongatus* Castelnau 1873 -- (Kuitert 2001:330 [ref. 25900]). **Current status:** Valid as *Hippocampus subelongatus* Castelnau 1873. Syngnathidae: Hippocampinae. Distribution: Australia: Western Australia. Habitat: marine.

***Hippocampus kampylotrachelos* BLEEKER, 1854**

kampylotrachelos, Hippocampus Bleeker [P.] 1854:107 [Natuurkundig Tijdschrift voor Nederlandsch Indië v. 7; ref. 16939] Priaman, Sumatra, Indonesia. Holotype (unique): RMNH 7257. •Synonym of *Hippocampus trimaculatus* Leach 1814 -- (Lourie et al. 1999:125 [ref. 23993]). •Valid as *Hippocampus kampylotrachelos* Bleeker 1854 -- (Kuitert 2001:307 [ref. 25900], Allen & Adrim 2003:28 [ref. 26830], Paxton et al. 2006:827 [ref. 29073], Kuitert 2009:122 [ref. 30404]). **Current status:** Valid as *Hippocampus kampylotrachelos* Bleeker 1854. Syngnathidae: Hippocampinae. Distribution: Southern Indonesia. Habitat: marine.

***Hippocampus planifrons* PETERS, 1877**

planifrons, Hippocampus Peters [W. (C. H.)] 1877:851 [Monatsberichte der Königlich Preuss[ischen] Akademie der Wissenschaften zu Berlin 1876; ref. [3454](#)] "Naturalists Channel", northwestern Australia, depth 10 fathoms. Holotype (unique): ZMB 9387. •Questionably a synonym of *Hippocampus trimaculatus* Leach 1814 -- (Lourie et al. 1999:171 [ref. [23993](#)]). •Valid as *Hippocampus planifrons* Peters 1877 -- (Paxton et al. 1989:422 [ref. [12442](#)], Larson & Williams 1997:352 [ref. [23967](#)], Francis 1993:159 [ref. [25479](#)], Paulus 1999:2269 [ref. [24794](#)], Johnson 1999:726 [ref. [25471](#)], Kuitert 2001:310 [ref. [25900](#)], Hutchins 2001:27 [ref. [25847](#)], Paxton et al. 2006:827 [ref. [29073](#)], Kuitert 2009:126 [ref. [30404](#)]). **Current status:** Valid as *Hippocampus planifrons* Peters 1877. Syngnathidae: Hippocampinae. Distribution: Western Australia, Australia: Shark Bay to Exmouth. Habitat: marine.

***Hippocampus taeniopterus* BLEEKER, 1852**

melanospilos, Hippocampus Bleeker [P.] 1854:505 [Natuurkundig Tijdschrift voor Nederlandsch Indië v. 6; ref. [344](#)] Ambon Island, Molucca Islands, Indonesia. Holotype (unique): RMNH 5165 (1 of 4). Bleeker specimens: BMNH 1867.11.28.362 (1), RMNH 5165 (3 of 4). •Synonym of *Hippocampus kuda* Bleeker 1852 -- (Lourie et al. 1999:109 [ref. [23993](#)]). •Synonym of *Hippocampus taeniopterus* Bleeker 1852 -- (Kuitert 2001:314 [ref. [25900](#)], Kuitert 2009:112 [ref. [30404](#)]). **Current status:** Synonym of *Hippocampus taeniopterus* Bleeker 1852. Syngnathidae: Hippocampinae. Habitat: brackish, marine.

novaehebudorum, Hippocampus Fowler [H. W.] 1944:162, Fig. 12 [Proceedings of the Academy of Natural Sciences of Philadelphia v. 96; ref. [1451](#)] Vanuatu. Holotype (unique): ANSP 71352. Type catalog: Böhlke 1984:159 [ref. [13621](#)]. Originally as *novae-hebudorum*. •Synonym of *Hippocampus kuda* Bleeker 1852 -- (Lourie et al. 1999:109 [ref. [23993](#)]). •Synonym of *Hippocampus taeniopterus* Bleeker 1852 -- (Kuitert 2009:19 [ref. [30404](#)]). **Current status:** Synonym of *Hippocampus taeniopterus* Bleeker 1852. Syngnathidae: Hippocampinae. Habitat: brackish, marine.

taeniopterus, Hippocampus Bleeker [P.] 1852:306 [Natuurkundig Tijdschrift voor Nederlandsch Indië v. 3; ref. [16825](#)] Ambon Island, Molucca Islands, Indonesia. Syntypes: (3) whereabouts unknown. •Synonym of *Hippocampus kuda* Bleeker 1852 -- (Lourie et al. 1999:109 [ref. [23993](#)]). •Valid as *Hippocampus taeniopterus* Bleeker 1852 -- (Horne 2001:245 [ref. [25696](#)], Kuitert 2001:314 [ref. [25900](#)], Allen & Adrim 2003:28 [ref. [26830](#)], Fricke 2004:42 [ref. [28269](#)], Paxton et al. 2006:828 [ref. [29073](#)], Kuitert 2009:112 [ref. [30404](#)], Fricke et al. 2011:376 [ref. [31242](#)], Allen & Erdmann 2012:206 [ref. [31980](#)], Larson et al. 2013:77 [ref. [32988](#)]). **Current status:** Valid as *Hippocampus taeniopterus* Bleeker 1852. Syngnathidae: Hippocampinae. Distribution: Western Pacific: Indonesia, Papua New Guinea and to Fiji, south to Australia and New Caledonia. Habitat: brackish, marine.

***Hippocampus tristis* CASTELNAU, 1872**

tristis, ***Hippocampus*** Castelnau [F. L.] 1872:197 [Proceedings of the Zoological and Acclimatisation Society of Victoria, Melbourne v. 1; ref. 757] Melbourne market, Victoria, Australia. Syntypes: MNHN A-4537 and 4538 (2). Type catalog: Bertin & Estève 1950:53 [ref. 19576]. •Probably a synonym of *Hippocampus whitei* Bleeker 1855 -- (Paxton et al. 1989:42 [ref. 12442], Gomon et al. 1994:449 [ref. 22532]). •Synonym of *Hippocampus kuda* Bleeker 1852 -- (Lourie et al. 1999:109 [ref. 23993]). •Valid as *Hippocampus tristis* Castelnau 1872 -- (Kuitert 2001:316 [ref. 25900], Paxton et al. 2006:829 [ref. 29073], Kuitert 2009:107 [ref. 30404]). **Current status:** Valid as *Hippocampus tristis* Castelnau 1872. Syngnathidae: Hippocampinae. Distribution: South Australia. Habitat: marine.

***Hippocampus tuberculatus* CASTELNAU, 1875**

tuberculatus*, *Hippocampus Castelnau [F. L.] 1875:48 [Researches on the fishes of Australia No. 2; ref. 768] Swan River, Western Australia. Holotype (unique): MNHN A-4539. •Synonym of *Hippocampus breviceps* Peters 1869 -- (Paxton et al. 1989:421 [ref. 12442], Gomon et al. 1994:448 [ref. 22532], Lourie et al. 1999:81 [ref. 23993]). •Valid as *Hippocampus tuberculatus* Castelnau 1875 -- (Hutchins 2001:27 [ref. 25847], Kuitert 2001:313 [ref. 25900], Paxton et al. 2006:829 [ref. 29073], Kuitert 2009:51 [ref. 30404]). **Current status:** Valid as *Hippocampus tuberculatus* Castelnau 1875. Syngnathidae: Hippocampinae. Distribution: Western Australia, Australia: Perth region north to Onslow. Habitat: brackish, marine.

List of standard references adopted by the Conference of the Parties

FAUNA

MAMMALIA

WILSON, D. E. & REEDER, D. M. (ed.) (2005): Mammal Species of the World. A Taxonomic and Geographic Reference. Third edition, Vol. 1-2, xxxv + 2142 pp. Baltimore (John Hopkins University Press). [for all mammals – with the exception of the recognition of the following names for wild forms of species (in preference to names for domestic forms): *Bos gaurus*, *Bos mutus*, *Bubalus arnee*, *Equus africanus*, *Equus przewalskii*, *Ovis orientalis ophion*; and with the exception of the taxa mentioned below]

WILSON, D. E. & REEDER, D. M. (1993): Mammal Species of the World: a Taxonomic and Geographic Reference. Second edition. xviii + 1207 pp., Washington (Smithsonian Institution Press). [for *Loxodonta africana*, *Puma concolor*, *Lama guanicoe* and *Ovis vignei*]

Cetaceae (in addition to the main reference noted under Mammalia above)

BEASLY, I., ROBERTSON, K. M. & ARNOLD, P. W. (2005): Description of a new dolphin, the Australian Snubfin Dolphin, *Orcaella heinsohni* sp. n. (Cetacea, Delphinidae). – Marine Mammal Science, **21**(3): 365-400. [for *Orcaella heinsohni*]

CABALLERO, S., TRUJILLO, F., VIANNA, J. A., BARRIOS-GARRIDO, H., MONTIEL, M. G., BELTRÁN-PEDREROS, S., MARMONTEL, M., SANTOS, M. C., ROSSI-SANTOS, M. R. & BAKER, C. S. (2007). Taxonomic status of the genus *Sotalia*: species level ranking for "tucuxi" (*Sotalia fluviatilis*) and "costero" (*Sotalia guianensis*) dolphins. - Marine Mammal Science, **23**: 358-386 [for *Sotalia fluviatilis* and *Sotalia guianensis*]

RICE, D. W., (1998): Marine Mammals of the World: Systematics and Distribution - Society of Marine Mammalogy Special Publication Number 4, The Society for Marine Mammalogy, Lawrence, Kansas [for *Physeter macrocephalus* and *Platanista gangetica*]

WADA, S., OISHI, M. & YAMADA, T. K. (2003): A newly discovered species of living baleen whales. – Nature, **426**: 278-281. [for *Balaenoptera omura*]

Primates (in addition to the main reference noted under Mammalia above)

BOUBLI, J. P., DA SILVA, M. N. F., AMADO, M. V., HRBEK, T., PONTUAL, F. B. & FARIAS, I. P. (2008): A taxonomic reassessment of *Cacajao melanocephalus* Humboldt (1811), with the description of two new species. – International Journal of Primatology, **29**: 723-741. [for *Cacajao ayresi*, *C. hosomi*]

BRANDON-JONES, D., EUDEY, A. A., GEISSMANN, T., GROVES, C. P., MELNICK, D. J., MORALES J. C., SHEKELLE, M. & STEWARD, C.-B. (2004): Asian primate classification. - International Journal of Primatology, **25**: 97-163. [for *Trachypithecus villosus*] S.

DAVENPORT, T. R. B., STANLEY, W. T., SARGIS, E. J., DE LUCA, D. W., MPUNGA, N. E., MACHAGA, S. J. & OLSON, L. E. (2006): A new genus of African monkey, *Rungwecebus*: Morphology, ecology, and molecular phylogenetics. – Science, **312**: 1378-1381. [for *Rungwecebus kipunjii*]

DEFLER, T. R. & BUENO, M. L. (2007): *Aotus* diversity and the species problem. – Primate Conservation, **22**: 55-70. [for *Aotus jorgehernandezii*]

DEFLER, T. R., BUENO, M. L. & GARCÍA, J. (2010): *Callicebus caquetensis*: a new and Critically Endangered titi monkey from southern Caquetá, Colombia. – Primate Conservation, **25**: 1-9. [for *Callicebus caquetensis*]

FERRARI, S. F., SENA, L., SCHNEIDER, M. P. C. & JÚNIOR, J. S. S. (2010): Rondon's Marmoset, *Mico rondoni* sp. n., from southwestern Brazilian Amazonia. – International Journal of Primatology, **31**: 693-714. [for *Mico rondoni*]

GEISMANN, T., LWIN, N., AUNG, S. S., AUNG, T. N., AUNG, Z. M., HLA, T. H., GRINDLEY, M. & MOMBERG, F. (2011): A new species of snub-nosed monkey, genus *Rhinopithecus* Milne-Edwards, 1872 (Primates, Colobinae), from Northern Kachin State, Northeastern Myanmar. – *Amer. J. Primatology*, **73**: 96-107. [for *Rhinopithecus strykeri*]

MERKER, S. & GROVES, C.P. (2006): *Tarsius lariang*: A new primate species from Western Central Sulawesi. – *International Journal of Primatology*, **27**(2): 465-485. [for *Tarsius lariang*]

OLIVEIRA, M. M. DE & LANGGUTH, A. (2006): Rediscovery of Marcgrave's Capuchin Monkey and designation of a neotype for *Simia flava* Schreber, 1774 (Primates, Cebidae). – *Boletim do Museu Nacional do Rio de Janeiro, N.S., Zoologia*, **523**: 1-16. [for *Cebus flavius*]

SHEKELLE, M., GROVES, C., MERKER, S. & SUPRIATNA, J. (2010): *Tarsius tumpara*: A new tarsier species from Siau Island, North Sulawesi. – *Primate Conservation*, **23**: 55-64. [for *Tarsius tumpara*]

SINHA, A., DATTA, A., MADHUSUDAN, M. D. & MISHRA, C. (2005): *Macaca munzala*: A new species from western Arunachal Pradesh, northeastern India. – *International Journal of Primatology*, **26**(4): 977-989: doi:10.1007/s10764-005-5333-3. [for *Macaca munzala*]

VAN NGOC THINH, MOOTNICK, A. R., VU NGOC THANH, NADLER, T. & ROOS, C. (2010): A new species of crested gibbon from the central Annamite mountain range. - *Vietnamese Journal of Primatology*, **4**: 1-12. [for *Nomascus annamensis*]

WALLACE, R. B., GÓMEZ, H., FELTON, A. & FELTON, A. (2006): On a new species of titi monkey, genus *Callicebus* Thomas (Primates, Pitheciidae), from western Bolivia with preliminary notes on distribution and abundance. – *Primate Conservation*, **20**: 29-39. [for *Callicebus aureipalatii*]

AVES

MORONY, J. J., BOCK, W. J. & FARRAND, J., Jr. (1975): [Reference List of the Birds of the World](#). American Museum of Natural History. 207 pp. [for order- and family-level names for birds]

DICKINSON, E.C. (ed.) (2003): *The Howard and Moore Complete Checklist of the Birds of the World. Revised and enlarged 3rd Edition. 1039 pp. London (Christopher Helm). [together with DICKINSON 2005 for all bird species – with the exception of the taxa mentioned below]*

DICKINSON, E.C. (2005): [Corrigenda 4 \(02.06.2005\) to Howard & Moore Edition 3 \(2003\)](#). http://www.naturalis.nl/sites/naturalis.en/contents/i000764/corrigenda%204_final.pdf (available on the CITES website) [together with DICKINSON 2003 for all bird species – except for the taxa mentioned below]

Apodiformes (in addition to the main reference noted under Aves above)

CORTÉS-DIAGO, A., ORTEGA, L. A., MAZARIEGOS-HURTADO, L. & WELLER, A.-A. (2007): A new species of *Eriocnemis* (Trochilidae) from southwest Colombia. – *Ornitologia Neotropical*, **18**: 161-170. [for *Eriocnemis isabellae*]

PACHECO, J. F. & WHITNEY, B. M. (2006): Mandatory changes to the scientific names of three Neotropical birds. - *Bull. Brit. Orn. Club*, **126**: 242-244. [for *Chlorostilbon lucidus*]

PIACENTINI, V. Q., ALEIXO, A. & SILVEIRA, L. F. (2009): Hybrid, subspecies or species? The validity and taxonomic status of *Phaethornis longuemareus aethopyga* Zimmer, 1950 (Trochilidae). - *Auk*, **126**: 604-612. [for *Phaethornis aethopyga*]

Falconiformes (in addition to the main reference noted under Aves above)

PARRY, S. J., CLARK, W. S. & PRAKASH, V. (2002) On the taxonomic status of the Indian Spotted Eagle *Aquila hastata*. – *Ibis*, **144**: 665-675. [for *Aquila hastata*]

PORTER, R. F. & KIRWAN, G. M. (2010): Studies of Socotran birds VI. The taxonomic status of the Socotra Buzzard. – *Bulletin of the British Ornithologists' Club*, **130** (2): 116–131. [for *Buteo socotraensis*]

WHITTAKER, A. (2002): A new species of forest-falcon (Falconidae: *Micrastur*) from southeastern Amazonia and the Atlantic rainforests of Brazil. – *Wilson Bulletin*, **114**: 421-445. [for *Micrastur mintoni*]

Passeriformes (in addition to the main reference noted under Aves above)

COLLAR, N. J. (2006): A partial revision of the Asian babblers (Timaliidae). – Forktail, **22**: 85-112. [for *Garrulax taewanus*]

Psittaciformes (in addition to the main reference noted under Aves above)

ARNDT, T. (2008): Anmerkungen zu einigen *Pyrrhura*-Formen mit der Beschreibung einer neuen Art und zweier neuer Unterarten. – Papageien, **8**: 278-286. [for *Pyrrhura parvifrons*]

COLLAR, N. J. (1997) Family Psittacidae (Parrots). In DEL HOYO, J., ELLIOT, A. AND SARGATAL, J. (eds.), Handbook of the Birds of the World, **4** (Sandgrouse to Cuckoos): 280-477. Barcelona (Lynx Edicions). [for *Psittacula intermedia* and *Trichoglossus haematodus*]

GABAN-LIMA, R., RAPOSO, M. A. & HOFLING, E. (2002): Description of a new species of *Pionopsitta* (Aves: Psittacidae) endemic to Brazil. – Auk, **119**: 815-819. [for *Pionopsitta aurantiocephala*]

NEMESIO, A. & RASMUSSEN, C. (2009): The rediscovery of Buffon's "Guarouba" or "Perriche jaune": two senior synonyms of *Aratinga pinto* SILVEIRA, LIMA & HÖFLING, 2005 (Aves: Psittaciformes). – Zootaxa, **2013**: 1-16. [for *Aratinga maculata*]

OLMOS, F., SILVA, W. A. G. & ALBANO, C. (2005): Grey-breasted Conure *Pyrrhura griseipectus*, an overlooked endangered species. – Cotinga, **24**: 77-83. [for *Pyrrhura griseipectus*]

PACHECO, J. F. & WHITNEY, B. M. (2006): Mandatory changes to the scientific names of three Neotropical birds. – Bull. Brit. Orn. Club, **126**: 242-244. [for *Forpus modestus*]

ROSELAAR, C. S. & MICHELS, J. P. (2004): Nomenclatural chaos untangled, resulting in the naming of the formally undescribed *Cacatua* species from the Tanimbar Islands, Indonesia (Psittaciformes: Cacatuidae). – Zoologische Verhandlungen, **350**: 183-196. [for *Cacatua goffiniana*]

Strigiformes (in addition to the main reference noted under Aves above)

DA SILVA, J. M. C., COELHO, G. & GONZAGA, P. (2002): Discovered on the brink of extinction: a new species of pygmy owl (Strigidae: Glaucidium) from Atlantic forest of northeastern Brazil. – Ararajuba, **10**(2): 123-130. [for *Glaucidium mooreorum*]

INDRAWAN, M. & SOMADIKARTA, S. (2004): A new hawk-owl from the Togian Islands, Gulf of Tomini, central Sulawesi, Indonesia. – Bulletin of the British Ornithologists' Club, **124**: 160-171. [for *Ninox burhani*]

WARAKAGODA, D. H. & RASMUSSEN, P. C. (2004): A new species of scops-owl from Sri Lanka. – Bulletin of the British Ornithologists' Club, **124**(2): 85-105. [for *Otus thilohoffmanni*]

REPTILIA

Crocodylia and Rhynchocephalia

WERMUTH, H. & MERTENS, R. (1996) (reprint): *Schildkröte, Krokodile, Brückenechsen*. xvii + 506 pp. Jena (Gustav Fischer Verlag). [for Crocodylia and Rhynchocephalia except for the taxon listed below]

TUCKER, A. D. (2010): The correct name to be applied to the Australian freshwater crocodile, *Crocodylus johnstoni* [Krefft, 1873]. – Australian Zoologist, **35**(2): 432-434. [for *Crocodylus johnstoni*]

Sauria

POUGH, F. H., ANDREWS, R. M., CADLE, J. E., CRUMP, M. L., SAVITZKY, A. H. & WELLS, K. D. (1998): Herpetology. Upper Saddle River/New Jersey (Prentice Hall). [for delimitation of families within the Sauria]

Agamidae

WILMS, T. M., BÖHME, W., WAGNER, P., LUTZMANN, N. & SCHMITZ, A. (2009): On the phylogeny and taxonomy of the genus *Uromastix* Merrem, 1820 (Reptilia: Squamata: Agamidae: Uromastycinae) – resurrection of the genus *Saara* Gray, 1845. – *Bonner zool. Beiträge*, **56**(1-2): 55-99. [for *Uromastix*, *Saara*]

Chamaeleonidae

KLAVER, C. J. J. & BÖHME, W. (1997): Chamaeleonidae – Das Tierreich, **112**, 85 pp. [for *Bradypodion*, *Brookesia*, *Calumma*, *Chamaeleo* and *Furcifer* – except for the taxa mentioned below, and except for the recognition of *Calumma andringitraense*, *C. guillaumeti*, *C. hilleniusi* and *C. marojezense* as valid species]

ANDREONE, F., MATTIOLI, F., JESU, R. & RANDRIANIRINA, J. E. (2001): [Two new chameleons of the genus *Calumma* from north-east Madagascar, with observations on hemipenial morphology in the *Calumma furcifer* group \(Reptilia, Squamata, Chamaeleonidae\)](#) – *Herpetological Journal*, **11**: 53-68. [for *Calumma vatosoa* and *Calumma vencesi*]

BAREJ, M. F., INEICH, I., GVOŽDÍK, V., LHERMITTE-VALLARINO, N., GONWOU, N. L., LEBRETON, M., BOTT, U. & SCHMITZ, A. (2010): Insights into chameleons of the genus *Trioceros* (Squamata: Chamaeleonidae) in Cameroon, with the resurrection of *Chamaeleo serratus* Mertens, 1922. – *Bonn zool. Bull.*, **57**(2): 211-229. [for *Trioceros perretti*, *Trioceros serratus*]

BÖHME, W. (1997): [Eine neue Chamäleon-Art aus der *Calumma gastrotaenia* – Verwandtschaft Ost-Madagaskars](#) – *Herpetofauna* (WeinStadt), **19** (107): 5-10. [for *Calumma glawi*]

BRANCH, W. R. & TOLLEY, K. A. (2010): A new species of chameleon (Sauria: Chamaeleonidae: *Nadzikambia*) from Mount Mabu, central Mozambique. – *Afr. J. Herpetology*, **59**(2): 157-172. doi: 10.1080/21564574.2010.516275. [for *Nadzikambia baylissi*]

BRANCH, W. R., TOLLEY, K. A. & TILBURY, C. R. (2006): A new Dwarf Chameleon (Sauria: *Bradypodion* Fitzinger, 1843) from the Cape Fold Mountains, South Africa. – *African Journal Herpetology*, **55**(2): 123-141. [for *Bradypodion atromontanum*]

GEHRING, P.-S., PABIJAN, M., RATSOAVINA, F. M., KÖHLER, J., VENCES, M. & GLAW, F. (2010): A Tarzan yell for conservation: a new chameleon, *Calumma tarzan* sp. n., proposed as a flagship species for the creation of new nature reserves in Madagascar. – *Salamandra*, **46**(3): 167-179. [for *Calumma tarzan*]

GEHRING, P.-S., RATSOAVINA, F. M., VENCES, M. & GLAW, F. (2011): *Calumma vohibola*, a new chameleon species (Squamata: Chamaeleonidae) from the littoral forests of eastern Madagascar. – *Afr. J. Herpetology*, **60**(2): 130-154. [for *Calumma vohibola*]

GLAW, F., KÖHLER, J. & VENCES, M. (2009b): A distinctive new species of chameleon of the genus *Furcifer* (Squamata: Chamaeleonidae) from the Montagne d'Ambre rainforest of northern Madagascar. – *Zootaxa*, **2269**: 32-42. [for *Furcifer timoni*]

GLAW, F. & M. VENCES (2007): A field guide to the amphibians and reptiles of Madagascar, third edition. Vences & Glaw Verlag, 496 pp. [for *Brookesia ramanantsoai*, *Calumma ambrense*]

GLAW, F., VENCES, M., ZIEGLER, T., BÖHME, W. & KÖHLER, J. 1999. Specific distinctiveness and biogeography of the dwarf chameleons *Brookesia minima*, *B. peyrierasi* and *B. tuberculata* (Reptilia: Chamaeleonidae): evidence from hemipenial and external morphology. – *J. Zool. Lond.*, **247**: 225-238. [for *Brookesia peyrierasi*, *B. tuberculata*]

JESU, R., MATTIOLI, F. & SCHIMENTI, G. (1999): [On the discovery of a new large chameleon inhabiting the limestone outcrops of western Madagascar: *Furcifer nicosiai* sp. nov. \(Reptilia, Chamaeleonidae\)](#) – *Doriana*, **7**(311): 1-14. [for *Furcifer nicosiai*]

KRAUSE, P. & BÖHME, W. (2010): A new chameleon of the *Trioceros bitaeniatus* complex from Mt. Hanang, Tanzania, East Africa (Squamata, Chamaeleonidae). – *Bonn Zoological Bulletin*, **57**: 19-29. [for *Trioceros hanangensis*]

LUTZMANN, N. & LUTZMANN, H. (2004): Das grammatikalische Geschlecht der Gattung *Calumma* (Chamaeleonidae) und die nötigen Anpassungen einiger Art- und Unterartbezeichnungen. – *Reptilia* (Münster), **9**(4): 4-5 (Addendum in issue 5: 13). [for *Calumma cucullatum*, *Calumma nasutum*]

MARIAUX, J., LUTZMANN, N. & STIPALA, J. (2008): The two-horned chameleons of East Africa. – *Zoological Journal Linnean Society*, **152**: 367-391. [for *Kinyongia vosseleri*, *Kinyongia boehmei*]

MENEGON, M., TOLLEY, K. A., JONES, T., ROVERO, F., MARSHALL, A. R. & TILBURY, C. R. (2009): A new species of chameleon (Sauria: Chamaeleonidae: *Kinyongia*) from the Magombera forest and the Udzungwa Mountains National Park, Tanzania. – *African Journal of Herpetology*, **58**(2): 59-70. [for *Kinyongia magomberae*]

- NECAS, P. (2009): Ein neues Chamäleon der Gattung *Kinyongia* Tilbury, Tolley & Branch 2006 aus den Poroto-Bergen, Süd-Tansania (Reptilia: Sauria: Chamaeleonidae). – *Sauria*, **31**(2): 41-48. [for *Kinyongia vanheygeni*]
- NECAS, P., MODRY, D. & SLAPETA, J. R. (2003): *Chamaeleo (Trioceros) narraioca* n. sp. (Reptilia Chamaeleonidae), a new chamaeleon species from a relict montane forest of Mount Kulal, northern Kenya. – *Tropical Zool.*, **16**:1-12. [for *Chamaeleo narraioca*]
- NECAS, P., MODRY, D. & SLAPETA, J. R. (2005): *Chamaeleo (Trioceros) ntunte* n. sp. a new chamaeleon species from Mt. Nyiru, northern Kenya (Squamata: Sauria: Chamaeleonidae). – *Herpetozoa*, **18**(3/4): 125-132. [for *Chamaeleo ntunte*]
- NECAS, P., SINDACO, R., KOŘENÝ, L., KOPEČNÁ, J., MALONZA, P. K. & MODRY, D. (2009): *Kinyongia asheorum* sp. n., a new montane chameleon from the Nyiro Range, northern Kenya (Squamata: Chamaeleonidae). – *Zootaxa*, **2028**: 41-50. [for *Kinyongia asheorum*]
- RAW, L. & BROTHERS, D. J. (2008): Redescription of the South African dwarf chameleon, *Bradypodion nemorale* Raw 1978 (Sauria: Chamaeleonidae), and description of two new species. – *ZooNova*, **1**(1): 1-7. [for *Bradypodion caeruleogula*, *Bradypodion nkandlae*]
- RAXWORTHY, C.J. & NUSSBAUM, R.A. (2006): Six new species of Occipital-Lobed *Calumma* Chameleons (Squamata: Chamaeleonidae) from Montane Regions of Madagascar, with a New Description and Revision of *Calumma brevicorne*. – *Copeia*, **4**: 711-734. [for *Calumma amber*, *Calumma brevicorne*, *Calumma crypticum*, *Calumma hafahafa*, *Calumma jeji*, *Calumma peltierorum*, *Calumma tsycorne*]
- STIPALA, J., LUTZMANN, N., MALONZA, P.K., BORGHESIO, L., WILKINSON, P., GODLEY, B. & EVANS, M.R. (2011): A new species of chameleon (Sauria: Chamaeleonidae) from the highlands of northwest Kenya. – *Zootaxa*, **3002**: 1-16. [for *Trioceros nyirif*]
- TILBURY, C. (1998): [Two new chameleons \(Sauria: Chamaeleonidae\) from isolated Afromontane forests in Sudan and Ethiopia](#) – *Bonner Zoologische Beiträge*, **47**: 293-299. [for *Chamaeleo balebicornutus* and *Chamaeleo conirostratus*]
- TILBURY, C. R. & TOLLEY, K. A. (2009a): A new species of dwarf chameleon (Sauria; Chamaeleonidae, *Bradypodion* Fitzinger) from KwaZulu Natal South Africa with notes on recent climatic shifts and their influence on speciation in the genus. – *Zootaxa*, **2226**: 43-57. [for *Bradypodion ngomeense*, *B. nkandlae*]
- TILBURY, C. R. & TOLLEY, K. A. (2009b): A re-appraisal of the systematics of the African genus *Chamaeleo* (Reptilia: Chamaeleonidae). – *Zootaxa*, **2079**: 57-68. [for *Trioceros*]
- TILBURY, C. R., TOLLEY, K. A. & BRANCH, R. B. (2007): Corrections to species names recently placed in *Kinyongia* and *Nadzikambia* (Reptilia: Chamaeleonidae). – *Zootaxa*, **1426**: 68. [for correct spelling of *Kinyongia uluguruensis*, *Nadzikambia mlanjensis*]
- TILBURY, C. R., TOLLEY, K. A. & BRANCH, W. R. (2006): A review of the systematics of the genus *Bradypodion* (Sauria: Chamaeleonidae), with the description of two new genera. – *Zootaxa*, **1363**: 23-38. [for *Kinyongia adolfifriderici*, *Kinyongia carpenteri*, *Kinyongia excubitor*, *Kinyongia fischeri*, *Kinyongia matschiei*, *Kinyongia multituberculata*, *Kinyongia oxyrhina*, *Kinyongia tavetana*, *Kinyongia tenuis*, *Kinyongia ulugurensis*, *Kinyongia uthmoelleri*, *Kinyongia xenorhina*, *Nadzikambia mlanjense*]
- TOLLEY, K. A., TILBURY, C. R., BRANCH, W. R. & MATHEE, C. A. (2004): Phylogenetics of the southern African dwarf chameleons, *Bradypodion* (Squamata: Chamaeleonidae). – *Molecular Phylogen. Evol.*, **30**: 351-365. [for *Bradypodion caffrum*, *Bradypodion damaranum*, *Bradypodion gutturale*, *Bradypodion occidentale*, *Bradypodion taeniobronchum*, *Bradypodion transvaalense*, *Bradypodion ventrale*]
- TOWNSEND, T. M., TOLLEY, K. A., GLAW, F., BÖHME, W. & VENCES, M. (2010): Eastward from Africa: paleocurrent-mediated chameleon dispersal to the Seychelles Islands. – *Biol. Lett.*, published online 8 September 2010, doi: 10.1098/rsbl.2010.0701 [for *Archaius tigris*]
- ULLENBRUCH, K., KRAUSE, P. & BÖHME, W. (2007): A new species of the *Chamaeleo dilepis* group (Sauria Chamaeleonidae) from West Africa. – *Tropical Zool.*, **20**: 1-17. [for *Chamaeleo necas*]
- WALBRÖL, U. & WALBRÖL, H. D. (2004): Bemerkungen zur Nomenklatur der Gattung *Calumma* (Gray, 1865) (Reptilia: Squamata: Chamaeleonidae). – *Sauria*, **26** (3): 41-44. [for *Calumma andringitraense*, *Calumma marojezense*, *Calumma tsaratanaense*]

Cordylidae

BROADLEY, D. G. (2006): [CITES Standard reference for the species of Cordylus \(Cordylidae, Reptilia\)](#) prepared at the request of the CITES Nomenclature Committee [for *Cordylus*]

Gekkonidae

Nactus serpensinsula

KLUGE, A.G. (1983): Cladistic relationships among gekkonid lizards. – *Copeia*, **1983**(no. 2): 465-475. [for *Nactus serpensinsula*]

Phelsuma spp.

HALLMANN, G., KRÜGER, J. & TRAUTMANN, G. (2008). Faszinierende Taggeckos. Die Gattung *Phelsuma*. 2. überarbeitete und erweiterte Auflage, 253 pp., Münster (Natur und Tier – Verlag). ISBN 978-3-86659-059-5. [for *Phelsuma* spp., however, with the retention of *Phelsuma ocellata* and except for the taxa mentioned below]

BERGHOF, H.-P. & TRAUTMANN, G. (2009): Eine neue Art der Gattung *Phelsuma* Gray, 1825 (Sauria: Gekkonidae) von der Ostküste Madagaskars. – *Sauria*, **31**(1): 5-14. [for *Phelsuma hoeschi*]

CROTTINI, A., GEHRING, P.-S., GLAW, F., HARRIS, D.J., LIMA, A. & VENCES, M. (2011): Deciphering the cryptic species diversity of dull-coloured day geckos *Phelsuma* (Squamata: Gekkonidae) from Madagascar, with description of a new species. – *Zootaxa*, **2982**: 40-48. [for *Phelsuma gouldi*]

GLAW, F., GEHRING, P.-S., KÖHLER, J., FRANZEN, M. & VENCES, M. (2010): A new dwarf species of day gecko, genus *Phelsuma*, from the Ankarana pinnacle karst in northern Madagascar. – *Salamandra*, **46**: 83-92. [for *Phelsuma roesleri*]

GLAW, F., KÖHLER, J. & VENCES, M. (2009a): A new species of cryptically coloured day gecko (*Phelsuma*) from the Tsingy de Bemaraha National Park in western Madagascar. – *Zootaxa*, **2195**: 61-68. [for *Phelsuma boraï*]

ROCHA, S., RÖSLER, H., GEHRING, P.-S., GLAW, F., POSADA, D., HARRIS, D. J. & VENCES, M. (2010): Phylogenetic systematics of day geckos, genus *Phelsuma*, based on molecular and morphological data (Squamata: Gekkonidae). – *Zootaxa*, **2429**: 1-28. [for *Phelsuma dorsovittata*, *P. parva*]

Uroplatus spp.

RAXWORTHY, C.J. (2003): Introduction to the reptiles. – In: Goodman, S.M. & Bernstead, J.P. (eds.), *The natural history of Madagascar*, : 934-949. Chicago. [for *Uroplatus* spp. except for the taxa mentioned below]

BÖHLE, A. & SCHÖNECKER, P. (2003): Eine neue Art der Gattung *Uroplatus* Duméril, 1805 aus Ost-Madagaskar (Reptilia: Squamata: Gekkonidae). – *Salamandra*, **39**(3/4): 129-138. [for *Uroplatus pietschmanni*]

GLAW, F., KOSUCH, J., HENKEL, W. F., SOUND, P. AND BÖHME, W. (2006): Genetic and morphological variation of the leaf-tailed gecko *Uroplatus fimbriatus* from Madagascar, with description of a new giant species. – *Salamandra*, **42**: 129-144. [for *Uroplatus giganteus*]

RATSOAVINA, F.M., LOUIS JR., E.E., CROTTINI, A., RANDRIANIAINA, R.-D., GLAW, F. & VENCES, M. (2011): A new leaf tailed gecko species from northern Madagascar with a preliminary assessment of molecular and morphological variability in the *Uroplatus ebenau* group. – *Zootaxa*, **3022**: 39-57. [for *Uroplatus finiavana*]

RAXWORTHY, C.J., PEARSON, R.G., ZIMKUS, B.M., REDDY, S., DEO, A.J., NUSSBAUM, R.A. & INGRAM, C.M. (2008): Continental speciation in the tropics: contrasting biogeographic patterns of divergence in the *Uroplatus* leaf-tailed gecko radiation of Madagascar. – *Journal of Zoology*, **275**: 423–440. [for *Uroplatus sameiti*]

Iguanidae

HOLLINGSWORTH, B. D. (2004): The Evolution of Iguanas: An Overview of Relationships and a Checklist of Species. pp. 19-44. In: Alberts, A. C., Carter, R. L., Hayes, W. K. & Martins, E. P. (Eds), *Iguanas: Biology and Conservation*. Berkeley (University of California Press). [for Iguanidae except for taxa mentioned below]

BURTON, F. J. (2004): [Revision to Species *Cyclura nubila lewisi*, the Grand Cayman Blue Iguana](#) – *Caribbean Journal of Science*, **40**(2): 198-203. [for *Cyclura lewisi*]

GENTILE, G. & SNELL, H. (2009): *Conolophus marthae* sp. nov. (Squamata, Iguanidae), a new species of land iguana from the Galápagos archipelago. – *Zootaxa*, **2201**: 1-10. [for *Conolophus marthae*]

KEOGH, J. S., EDWARDS, D. L., FISHER, R. N. & HARLOW, P. S. (2008): Molecular and morphological analysis of the critically endangered Fijian iguanas reveals cryptic diversity and a complex biogeographic history. – Phil. Trans. R. Soc. B, **363**(1508): 3413-3426. [for *Brachylophus bulabula*]

MONTANUCCI, R.R. (2004): Geographic variation in *Phrynosoma coronatum* (Lacertilia, Phrynosomatidae): further evidence for a peninsular archipelago. – Herpetologica, **60**: 117. [for *Phrynosoma blainvillii*, *Phrynosoma cerroense*, *Phrynosoma wigginsi*]

Teiidae

AVILA PIRES, T. C. S. (1995): Lizards of Brazilian Amazonia (Reptilia: Squamata) – Zoologische Verhandelingen, **299**: 706 pp. [for *Tupinambis* except for the taxa mentioned below]

CEI, J. M. (1993): Reptiles del noroeste, nordeste y este de la Argentina – herpetofauna de las selvas subtropicales, Puna y Pampa – Monografía XIV, Museo Regionale di Scienze Naturali. [for *Tupinambis* except for the taxa mentioned below]

COLLI, G. R., PÉRES, A. K. & DA CUNHA, H. J. (1998): A new species of *Tupinambis* (Squamata: Teiidae) from central Brazil, with an analysis of morphological and genetic variation in the genus – Herpetologica, **54**: 477-492. [for *Tupinambis cerradensis*]

FITZGERALD, L. A., COOK, J. A. & LUZ AQUINO, A. (1999): Molecular Phylogenetics and Conservation of *Tupinambis* (Sauria: Teiidae). – Copeia, **4**: 894-905. [for *Tupinambis duseni*]

MANZANI, P. R. & ABE, A. S. (1997): A new species of *Tupinambis* Daudin, 1802 (Squamata, Teiidae) from central Brazil – Boletim do Museu Nacional Nov. Ser. Zool., **382**: 1-10. [for *Tupinambis quadrilineatus*]

MANZANI, P. R. & ABE, A. S. (2002): A new species of *Tupinambis* Daudin, 1803 from southeastern Brazil – Arquivos do Museu Nacional, Rio de Janeiro, **60**(4): 295-302. [for *Tupinambis palustris*]

MASSARY, J.-C. DE & HOOGMOED, M. (2001): [The valid name for *Crocodylurus lacertinus auctorum* \(nec Daudin, 1802\) \(Squamata: Teiidae\)](#) – Journal of Herpetology, **35**: 353-357. [for *Crocodylurus amazonicus*]

Varanidae

BÖHME, W. (2003): Checklist of the living monitor lizards of the world (family Varanidae) – Zoologische Verhandelingen. Leiden, **341**: 1-43. [for Varanidae]

KOCH, A., AULIYA, M. & ZIEGLER, T. (2010): Updated Checklist of the living monitor lizards of the world (Squamata: Varanidae). - Bonn zool. Bull., **57**(2): 127-136. [for Varanidae]

Serpentes

McDIARMID, R. W., CAMPBELL, J. A. & TOURÉ, T. A. (1999): Snake Species of the World. A Taxonomic and Geographic Reference. Volume 1, Washington, DC. (The Herpetologists' League). [for Loxocemidae, Pythonidae, Boidae, Bolyeriidae, Tropicophiidae and Viperidae – except for the retention of the genera *Acrantophis*, *Sanzinia*, *Calabaria* and *Lichanura*, the recognition of *Epicrates maurus* as valid species and except for the species mentioned below]

Boidae (in addition to the main reference noted under Serpentes above)

DIRKSEN, L. (2002): *Anakondas*. NTV Wissenschaft. [for *Eunectes beniensis*]

HENDERSON, R. W., PASSOS, P. & FEITOSA, D. (2009): Geographic variation in the Emerald Treeboa, *Corallus caninus* (Squamata: Boidae). – Copeia, **2009** (3): 572-582. [for *Corallus batesii*]

LANZA, B. & NISTRÌ, A. (2005): Somali Boidae (genus *Eryx* Daudin 1803) and Pythonidae (genus *Python* Daudin 1803) (Reptilia Serpentes). – Tropical Zoology, **18**(1): 67-136. [for *Eryx borrii*]

PASSOS, P. & FERNANDES, R. (2008): Revision of the *Epicrates cenchria* complex (Serpentes: Boidae). – Herpetol. Monographs, **22**: 1-30. [for *Epicrates crassus*, *E. assisi*, *E. alvarezii*]

SMITH, H. M., CHISZAR, D., TEPEDELEN, K. & VAN BREUKELLEN, F. (2001): A revision of the bevelnosed boas (*Candoia carinata* complex) (Reptilia: Serpentes). – Hamadryad, **26**(2): 283-315. [for *Candoia paulsoni*, *C. superciliosa*]

Elapidae (in addition to the main reference noted under Serpentes above)

SLOWINSKI, J. B. & WÜSTER, W. (2000.): [A new cobra \(Elapidae: *Naja*\) from Myanmar \(Burma\)](#) – *Herpetologica*, **56**: 257-270. [for *Naja mandalayensis*]

WÜSTER, W. (1996): Taxonomic change and toxinology: systematic revisions of the Asiatic cobras (*Naja naja* species complex) – *Toxicon*, **34**: 339-406. [for *Naja atra*, *Naja kaouthia*, *Naja oxiana*, *Naja philippinensis*, *Naja sagittifera*, *Naja samarensis*, *Naja siamensis*, *Naja sputatrix* and *Naja sumatrana*]

Pythonidae (in addition to the main reference noted under Serpentes above)

BROADLEY, D. G. (1999): The southern African python, *Python natalensis* A. Smith 1840, is a valid species. – *African Herp News*, **29**: 31-32. [for *Python natalensis*]

HARVEY, M. B., BARKER, D. B., AMMERMAN, L. K. & CHIPPINDALE, P. T. (2000): Systematics of pythons of the *Morelia amethystina* complex (Serpentes: Boidae) with the description of three new species – [Herpetological Monographs](#), **14**: 139-185. [for *Morelia clastolepis*, *Morelia nauta* and *Morelia tracyae*, and elevation to species level of *Morelia kinghorni*]

JACOBS, H. J., AULIYA, M. & BÖHME, W. (2009): Zur Taxonomie des Dunklen Tigerpythons, *Python molurus bivittatus* KUHL, 1820, speziell der Population von Sulawesi. – *Sauria*, **31**: 5-16. [for *Python bivittatus*]

KEOGH, J. S., BARKER, D. G. & SHINE, R. 2001. Heavily exploited but poorly known: systematics and biogeography of commercially harvested pythons (*Python curtus* group) in Southeast Asia – *Biological Journal of the Linnean Society*, **73**: 113-129. [for *Python breitensteini* and *Python brongersmai*]

SCHLEIP, W. D. (2008): Revision of the genus *Leiopython* Hubrecht 1879 (Serpentes: Pythonidae) with the redescription of taxa recently described by Hoser (2000) and the description of new species. – *Journal of Herpetology*, **42**(4): 645–667. [for *Leiopython bennetorum*, *L. biakensis*, *L. fredparkeri*, *L. huonensis*, *L. hoserae*]

ZUG, G.R., GROTTÉ, S. W. & JACOBS, J. F. (2011): Pythons in Burma: Short-tailed python (Reptilia: Squamata). – *Proc. biol. Soc. Washington*, **124**(2): 112-136. [for *Python kyaiktiyo*]

Tropidophiidae (in addition to the main reference noted under Serpentes above)

DOMÍNGUEZ, M., MORENO, L. V. & HEDGES, S. B. (2006): A new snake of the genus *Tropidophis* (Tropidophiidae) from the Guanahacabibes Peninsula of Western Cuba. – *Amphibia-Reptilia*, **27**(3): 427-432. [for *Tropidophis xanthogaster*]

HEDGES, B. S. & GARRIDO, O. (1999): [A new snake of the genus *Tropidophis* \(Tropidophiidae\) from central Cuba](#) – *Journal of Herpetology*, **33**: 436-441. [for *Tropidophis spiritus*]

HEDGES, B. S. & GARRIDO, O. (2002): [A new snake of the genus *Tropidophis* \(Tropidophiidae\) from Eastern Cuba](#) – *Journal of Herpetology*, **36**: 157-161. [for *Tropidophis hendersoni*]

HEDGES, B. S., ESTRADA, A. R. & DIAZ, L. M. (1999): [New snake \(*Tropidophis*\) from western Cuba](#) – *Copeia*, **1999**(2): 376-381. [for *Tropidophis celiae*]

HEDGES, B. S., GARRIDO, O. & DIAZ, L. M. (2001): [A new banded snake of the genus *Tropidophis* \(Tropidophiidae\) from north-central Cuba](#) – *Journal of Herpetology*, **35**: 615-617. [for *Tropidophis morenoi*]

Testudines

WERMUTH, H. & MERTENS, R. (1996) (reprint): Schildkröte, Krokodile, Brückenechsen. xvii + 506 pp. Jena (Gustav Fischer Verlag). [for Testudines order names]

FRITZ, U. & HAVAŠ, P. (2007): Checklist of Chelonians of the World. – *Vertebrate Zoology*, **57**(2): 149-368. Dresden. ISSN 1864-5755 [without its appendix; for Testudines for species and family names – with the exception of the retention of the following names *Mauremys iversoni*, *Mauremys pritchardi*, *Ocadia glyphistoma*, *Ocadia philippeni*, *Sacalia pseudocellata*, and except for the taxa mentioned below]

Testudinidae (in addition to the main reference noted under Testudines above)

BRANCH, W. R. (2007): A new species of tortoise of the genus *Homopus* (Chelonia: Testudinidae) from southern Namibia. – African Journal of Herpetology, **56**(1): 1-21. [for *Homopus solus*]

MURPHY, R. W., BERRY, K. H., EDWARDS, T., LEVITON, A. E., LATHROP, A. & RIEDLE, J. D. (2011): The dazed and confused identity of Agassiz's land tortoise, *Gopherus agassizii* (Testudines, Testudinidae) with the description of a new species, and its consequences for conservation. – Zookeys, **113**: 39-71. [for *Gopherus morafka*]

Emydidae (in addition to the main reference noted under Testudines above)

ENNEN, J. R., LOVICH, J. E., KREISER, B. R., SELMAN, W. & QUALLS, C. P. (2010): Genetic and morphological variation between populations of the Pascagoula Map Turtle (*Graptemys gibbonsi*) in the Pearl and Pascagoula Rivers with description of a new species. – Chelonian Conservation and Biology, **9**(1): 98-113. [for *Graptemys pearlensis*]

Geoemydidae (in addition to the main reference noted under Testudines above)

PRASCHAG, P., HUNSDÖRFER, A. K. & FRITZ, U. (2007): Phylogeny and taxonomy of endangered South and South-east Asian freshwater turtles elucidated by mtDNA sequence variation (Testudines: Geoemydidae: *Batagur*, *Callagur*, *Hardella*, *Kachuga*, *Pangshura*). – Zoologica Scripta, **36**: 429-442. [for *Batagur borneoensis*, *Batagur dhongoka*, *Batagur kachuga*, *Batagur trivittata*]

PRASCHAG, P., SOMMER, R. S., MCCARTHY, C., GEMEL, R. & FRITZ, U. (2008): Naming one of the world's rarest chelonians, the southern Batagur. – Zootaxa, **1758**: 61-68. [for *Batagur affinis*]

Trionychidae (in addition to the main reference noted under Testudines above)

PRASCHAG, P., HUNSDÖRFER, A.K., REZA, A.H.M.A. & FRITZ, U. (2007): Genetic evidence for wild-living *Aspideretes nigricans* and a molecular phylogeny of South Asian softshell turtles (Reptilia: Trionychidae: *Aspideretes*, *Nilssonina*). – Zoologica Scripta, **36**: 301-310. [for *Nilssonina gangeticus*, *N. hurum*, *N. nigricans*]

PRASCHAG, P., STUCKAS, H., PÄCKERT, M., MARAN, J. & FRITZ, U. (2011): Mitochondrial DNA sequences suggest a revised taxonomy of Asian flapshell turtles (*Lissemys* Smith, 1931) and the validity of previously unrecognized taxa (Testudines: Trionychidae). – Vertebrate Zoology, **61**(1): 147-160. [for *Lissemys ceylonensis*]

AMPHIBIA

Taxonomic Checklist of CITES-listed Amphibians, information extracted from FROST, D. R. (ed.) (2011), Amphibian Species of the World: a taxonomic and geographic reference, an online reference (<http://research.amnh.org/herpetology/amphibia/index.html>) Version 5.5 as of December 2011 in combination with BROWN, J. L., TWOMEY, E., AMÉZQUITA, A., BARBOSA DE SOUZA, M., CALDWELL, L. P., LÖTTERS, S., VON MAY, R., MELO-SAMPAIO, P. R., MEJÍA-VARGAS, D., PEREZ-PEÑA, P., PEPPER, M., POELMAN, E. H., SANCHEZ-RODRIGUEZ, M. & SUMMERS, K. (2011): A taxonomic revision of the Neotropical poison frog genus *Ranitomeya* (Amphibia: Dendrobatidae). – Zootaxa, **3083**: 1-120. [for all Amphibian species]

ELASMOBRANCHII, ACTINOPTERYGII and SARCOPTERYGII

Taxonomic Checklist of all CITES listed Fish species (Elasmobranchii and Actinopterygii, except the genus *Hippocampus*), information extracted from ESCHMEYER, W.N. & FRICKE, R. (eds.): Catalog of Fishes, an online reference (<http://research.calacademy.org/redirect?url=http://researcharchive.calacademy.org/research/Ichthyology/catalog/fishcatmain.asp>), version downloaded 30 November 2011. [for all fish species, except the genus *Hippocampus*]

Hippocampus spp.

HORNE, M. L. (2001): [A new seahorse species \(Syngnathidae: *Hippocampus*\) from the Great Barrier Reef](#) – Records of the Australian Museum, **53**: 243-246. [for *Hippocampus*]

KUITER, R. H. (2001): [Revision of the Australian seahorses of the genus *Hippocampus* \(Syngnathiformes: Syngnathidae\) with a description of nine new species](#) – Records of the Australian Museum, **53**: 293-340. [for *Hippocampus*]

KUITER, R. H. (2003): [A new pygmy seahorse \(Pisces: Syngnathidae: Hippocampus\) from Lord Howe Island](#) – Records of the Australian Museum, **55**: 113-116. [for *Hippocampus*]

LOURIE, S. A. & RANDALL, J. E. (2003): A new pygmy seahorse, *Hippocampus denise* (Teleostei: Syngnathidae), from the Indo-Pacific – Zoological Studies, **42**: 284-291. [for *Hippocampus*]

LOURIE, S. A., VINCENT, A. C. J. & HALL, H. J. (1999): Seahorses. An identification guide to the world's species and their conservation. Project Seahorse (ISBN 0 9534693 0 1) (Second edition available on CD-ROM). [for *Hippocampus*]

FOSTER, R. & GOMON, M. F. (2010): A new seahorse (Teleostei: Syngnathidae: Hippocampus) from south-western Australia. – Zootaxa, **2613**: 61-68. [for *Hippocampus paradoxus*]

GOMON, M. F. & KUITER, R. H. (2009): Two new pygmy seahorses (Teleostei: Syngnathidae: *Hippocampus*) from the Indo-West Pacific. – Aqua, Int. J. of Ichthyology, **15**(1): 37-44. [for *Hippocampus debelius*, *Hippocampus waleanus*]

LOURIE, S. A. & KUITER, R. H. (2008): Three new pygmy seahorse species from Indonesia (Teleostei: Syngnathidae: *Hippocampus*). – Zootaxa, **1963**: 54-68. [for *Hippocampus pontohi*, *Hippocampus satomiae*, *Hippocampus severnsi*]

RANDALL, J. & LOURIE, S. A. (2009): *Hippocampus tyro*, a new seahorse (Gasterosteiformes: Syngnathidae) from the Seychelles. – Smithiana Bulletin, **10**: 19-21. [for *Hippocampus tyro*]

PIACENTINO, G. L. M. AND LUZZATTO, D. C. (2004): *Hippocampus patagonicus* sp. nov., new seahorse from Argentina (Pisces, Syngnathiformes). – Revista del Museo Argentino de Ciencias Naturales, **6**(2): 339-349. [for *Hippocampus patagonicus*]

ARACHNIDA

Araneae

[Taxonomic Checklist of CITES listed Spider Species](#), information extracted from PLATNICK, N. (2006), The World Spider Catalog, an online reference, Version 6.5 as of 7 April 2006 [for Theraphosidae except for the taxon mentioned below]

RUDLOFF, J.-P. (2008): Eine neue *Brachypelma*-Art aus Mexiko (Araneae: Mygalomorphae: Theraphosidae: Theraphosinae). – Arthropoda, **16**(2): 26-30. [for *Brachypelma kahlenbergi*]

Scorpiones

LOURENÇO, W. R. & CLOUDSLEY-THOMPSON, J. C. (1996): [Recognition and distribution of the scorpions of the genus Pandinus Thorell. 1876 accorded protection by the Washington Convention](#) – Biogeographica, **72**(3): 133-143. [for scorpions of the genus *Pandinus*]

INSECTA

BARTOLOZZI, L. (2005): Description of two new stag beetle species from South Africa (Coleoptera: Lucanidae). – African Entomology, **13**(2): 347-352. [for *Colophon endroedyi*]

MATSUKA, H. (2001): Natural History of Birdwing Butterflies. 367 pp. Tokyo (Matsuka Shuppan). (ISBN 4-9900697-0-6). [for birdwing butterflies of the genera *Ornithoptera*, *Trogonoptera* and *Troides*]

HIRODINOIDEA

NESEMANN, H. & NEUBERT, E. (1999): Annelida: Clitellata: Branchiobdellida, Acanthobdellea, Hirudinea. – Süßwasserfauna von Mitteleuropa, vol. **6**/2, 178 pp., Berlin (Spektrum Akad. Verlag). ISBN 3-8274-0927-6. [for *Hirudo medicinalis* and *Hirudo verbana*]

ANTHOZOA and HYDROZOA

Taxonomic Checklist of all CITES listed Coral Species, based on information compiled by UNEP-WCMC 2012.

UNEP-WCMC **technical report**

Bird taxonomy

Comparison of the generic and species taxonomies in the 3rd and 4th editions of *The Howard & Moore complete checklist of the birds of the world*, relating to taxa listed in the EU Wildlife Trade Regulations (which includes all CITES listed species)



Comparison of the generic and species taxonomies in the 3rd and 4th editions of *The Howard & Moore complete checklist of the birds of the world*, relating to taxa listed in the EU Wildlife Trade Regulations (including CITES listed species)

Prepared for

The European Commission, Directorate General Environment, Directorate E - Global & Regional Challenges, LIFE ENV.E.2. – Global Sustainability, Trade & Multilateral Agreements, Brussels, Belgium

Prepared January 2014

Copyright

European Commission 2014

Citation

UNEP-WCMC. 2014. Bird taxonomy: Comparison of the generic and species taxonomies in the 3rd and 4th editions of *The Howard & Moore complete checklist of the birds of the world*, relating to taxa listed in the EU Wildlife Trade Regulations (including CITES listed species). UNEP-WCMC, Cambridge.



The UNEP World Conservation Monitoring Centre (UNEP-WCMC) is the specialist biodiversity assessment of the United Nations Environment Programme, the world's foremost intergovernmental environmental organization. The Centre has been in operation for over 30 years, combining scientific research with policy advice and the development of decision tools.

We are able to provide objective, scientifically rigorous products and services to help decision-makers recognize the value of biodiversity and apply this knowledge to all that they do. To do this, we collate and verify data on biodiversity and ecosystem services that we analyze and interpret in comprehensive assessments, making the results available in appropriate forms for national and international level decision-makers and businesses. To ensure that our work is both sustainable and equitable we seek to build the capacity of partners where needed, so that they can provide the same services at national and regional scales.

The contents of this report do not necessarily reflect the views or policies of UNEP, contributory organisations or editors. The designations employed and the presentations do not imply the expressions of any opinion whatsoever on the part of UNEP, the European Commission or contributory organisations, editors or publishers concerning the legal status of any country, territory, city area or its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of a commercial entity or product in this publication does not imply endorsement by UNEP.

UNEP World Conservation Monitoring Centre (UNEP-WCMC)

219 Huntingdon Road,
Cambridge CB3 0DL, UK
Tel: +44 1223 277314
www.unep-wcmc.org

UNEP promotes environmentally sound practices globally and in its own activities. Printing on paper from environmentally sustainable forests and recycled fibre is encouraged.

Introduction

This report provides an overview of taxonomic updates for bird species that the CITES Animals Committee may wish to consider in the context of updating the CITES Standard Reference.

The revised and enlarged 3rd edition of *The Howard and Moore Complete Checklist of the Birds of the World* (Dickinson, 2003)⁶ is the current CITES standard nomenclatural reference for most birds (Resolution Conf. 12.11 Rev. CoP16). This is also the standard reference determining the nomenclature of birds in the EU Wildlife Trade Regulations. In 2013, a 4th edition of this Checklist was published for non-passerines (Dickinson & Remsen, 2013)⁷.

The table below provides a comparison of the generic and species taxonomy adopted in Dickinson (2003) (and its 4th corrigenda) with that adopted in Dickinson & Remsen (2013), relating to taxa listed in CITES and in the EU Wildlife Trade Regulations.

⁶ Dickinson, E.C. (ed.) 2003. *The Howard and Moore Complete Checklist of the Birds of the World. Revised and enlarged 3rd Edition*. 1039 pp. London: Christopher Helm.

⁷ Dickinson, E.C. & Remsen Jr. J.V. (eds.) 2013. *The Howard and Moore Complete Checklist of the Birds of the World. 4th Edition. Volume one: Non-passerines*. 461 pp. Eastbourne: Aves Press.

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
ANSERIFORMES				
ANATIDAE				
<i>Anas chlorotis</i> G. R. Gray, 1845	I/A	< ⁸	<i>Anas aucklandica</i> (G. R. Gray, 1844)	Species lump
<i>Anas nesiotis</i> (J. H. Fleming, 1935)	I/A	<	<i>Anas aucklandica</i> (G. R. Gray, 1844)	Species lump
<i>Branta canadensis leucopareia</i> (von Brandt, 1836)	I/A	= ⁹	<i>Branta hutchinsii leucopareia</i> (von Brandt, 1836)	Subspecies transfer
APODIFORMES				
TROCHILIDAE				
<i>Aglaiocercus kingi</i> (Lesson, 1832)	II/B	> ¹⁰	<i>Aglaiocercus kingii</i> (Lesson, 1832)	Spelling correction
			<i>Aglaiocercus berlepschi</i> (E. Hartert, 1898)	Species split
<i>Amazilia alfaroana</i> Underwood, 1896	II/B	<	<i>Amazilia cyanifrons</i> (Bourcier, 1843)	Species lump
<i>Amazilia alticola</i> Gould, 1860	II/B	<	<i>Amazilia amazilia</i> (Lesson & Garnot, 1827)	Species lump
<i>Amazilia amabilis</i> (Gould, 1853)	II/B	>	<i>Amazilia amabilis</i> (Gould, 1853)	
			<i>Amazilia decora</i> (Salvin, 1891)	Species split
<i>Amazilia cupreicauda</i> Salvin & Godman, 1884	II/B	<	<i>Amazilia viridigaster</i> (Bourcier, 1843)	Species lump
<i>Amazilia rondoniae</i> Ruschi, 1982	II/B	<	<i>Amazilia versicolor</i> (Vieillot, 1818)	Species lump
<i>Anthracothorax prevostii iridescens</i> (Gould, 1861)	II/B	=	<i>Anthracothorax nigricollis iridescens</i> (Gould, 1861)	Subspecies transfer
<i>Anthracothorax recurvirostris</i> (Swainson, 1822)	II/B	=	<i>Avocettula recurvirostris</i> (Swainson, 1822)	Generic change
<i>Basilinna leucotis</i> (Vieillot, 1818)	II/B	=	<i>Hylocharis leucotis</i> (Vieillot, 1818)	Generic change
<i>Basilinna xantusii</i> (Lawrence, 1860)	II/B	=	<i>Hylocharis xantusii</i> (Lawrence, 1861)	Generic change and date correction
<i>Campylopterus curvipennis</i> (Deppe, 1830)	II/B	>	<i>Campylopterus curvipennis</i> (Deppe, 1830) (including ssp. <i>curvipennis</i> , <i>pampa</i> , <i>yucatanensis</i>)	
			<i>Campylopterus excellens</i> (Wetmore, 1941)	Species split
<i>Campylopterus cuvierii</i> (DeLattre & Bourcier, 1846)	II/B	=	<i>Phaeochroa cuvierii</i> (DeLattre & Bourcier, 1846)	Generic change
<i>Chaetocercus mulsanti</i> (Bourcier, 1842)	II/B	=	<i>Chaetocercus mulsant</i> (Bourcier, 1843)	Spelling and date correction

⁸ Species lumps (indicated by the symbol "<") refer to taxa recognised as separate in Dickinson (2003) but that have been grouped together under another name in Dickinson & Remsen (2013).

⁹ The symbol "=" is used to indicate taxonomic or nomenclature changes between Dickinson (2003) and Dickinson & Remsen (2013) that do not involve a change in the scope of the taxon in question.

¹⁰ Species splits (indicated by the symbol ">") refer to cases where one taxon as recognised in Dickinson (2003) has been split into various taxa in Dickinson & Remsen (2013).

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Chlorostilbon aureoventris</i> (d'Orbigny & Lafresnaye, 1838)	II/B	=	<i>Chlorostilbon lucidus</i> (Shaw, 1812)	Replacement name. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Chlorostilbon mellisugus</i> (Linnaeus, 1758)	II/B	>	<i>Chlorostilbon mellisugus</i> (Linnaeus, 1758) (including ssp. <i>mellisugus</i> , <i>caribaeus</i> , <i>duidae</i> , <i>napensis</i> , <i>peruanus</i> , <i>phoeopygus</i> , <i>subfurcatus</i>)	
			<i>Chlorostilbon gibsoni</i> (Fraser, 1840) (including ssp. <i>gibsoni</i> , <i>chrysogaster</i> , <i>nitens</i>)	Species split
			<i>Chlorostilbon melanorhynchus</i> Gould, 1860	Species split
<i>Chlorostilbon notatus</i> (Reich, 1795)	II/B	=	<i>Chlorestes notata</i> (Reich, 1793)	Generic change and date correction
<i>Coeligena bonapartei</i> (Boissonneau, 1840)	II/B	>	<i>Coeligena bonapartei</i> (Boissonneau, 1840) (including ssp. <i>bonapartei</i> , <i>consita</i> , <i>eos</i>)	
			<i>Coeligena orina</i> Wetmore, 1953	Species split
<i>Damophila julie</i> (Bourcier, 1842)	II/B	=	<i>Juliamyia julie</i> (Bourcier, 1843)	Generic change and date correction
<i>Eriocnemis alinae</i> (Bourcier, 1842)	II/B	=	<i>Eriocnemis aline</i> (Bourcier, 1843)	Spelling and date correction
N/A	II/B		<i>Eriocnemis isabellae</i> Corté-Diago, Ortega, Mazariegos-Hurtado & Weller, 2007	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Heliangelus amethysticollis</i> (d'Orbigny & Lafresnaye, 1838)	II/B	>	<i>Heliangelus amethysticollis</i> (d'Orbigny & Lafresnaye, 1838) (including ssp. <i>amethysticollis</i> , <i>clarisse</i> , <i>decolor</i> , <i>laticlavus</i> , <i>violiceps</i> & <i>apurimacensis</i> Weller, 2009)	
			<i>Heliangelus spencei</i> (Bourcier, 1847)	Species split
<i>Hylocharis grayi</i> (DeLattre & Bourcier, 1846)	II/B	>	<i>Hylocharis grayi</i> (DeLattre & Bourcier, 1846)	
			<i>Hylocharis humboldtii</i> (Bourcier & Mulsant, 1852)	Species split
<i>Phaethornis ruber</i> x <i>P. rufurumii</i>	II/B	=	<i>Phaethornis aethopygus</i> J. T. Zimmer, 1950	Species revalidated. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Sappho sparganura</i> (Shaw, 1812)	II/B	=	<i>Sappho sparganurus</i> (Shaw, 1812)	Spelling correction
<i>Sephanoides sephanoides</i> (Lesson, 1827)	II/B	=	<i>Sephanoides sephanoides</i> (Lesson & Garnot, 1827)	See H&M 3 corrigendum 2.1 for spelling as 'sephanoides'. Author correction.
<i>Stellula calliope</i> (Gould, 1847)	II/B	=	<i>Selasphorus calliope</i> (Gould, 1847)	Generic change
<i>Thalurania fannyi</i> (DeLattre & Bourcier, 1846)	II/B	<	<i>Thalurania colombica</i> (Bourcier, 1843)	Species lump
<i>Threnetes niger</i> (Linnaeus, 1758)	II/B	>	<i>Threnetes niger</i> (Linnaeus, 1766) (including ssp. <i>niger</i> , <i>loehkeni</i>)	

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
			<i>Threnetes leucurus</i> (Linnaeus, 1766) (including ssp. <i>leucurus</i> , <i>cervinicauda</i> , <i>medianus</i> , <i>rufigastra</i>)	Species split
<i>Urosticte benjamini</i> (Bourcier, 1851)	II/B	>	<i>Urosticte benjamini</i> (Bourcier, 1851)	
			<i>Urosticte ruficrissa</i> Lawrence, 1864	Species split
CICONIIFORMES				
PHOENICOPTERIDAE				
<i>Phoenicopterus ruber</i> Linnaeus, 1758	II/A	>	<i>Phoenicopterus ruber</i> Linnaeus, 1758	
			<i>Phoenicopterus roseus</i> Pallas, 1811	Species split
COLUMBIFORMES				
COLUMBIDAE				
<i>Claravis godefrida</i> (Temminck, 1811)	-/A	=	<i>Claravis geoffroyi</i> (Temminck, 1811)	Replacement name
<i>Nesoenas mayeri</i> (Prévost, 1843)	III/C	=	<i>Streptopelia mayeri</i> (Prévost, 1843)	Generic change
CORACIIFORMES				
BUCEROTIDAE				
<i>Aceros cassidix</i> (Temminck, 1823)	II/B	=	<i>Rhyticeros cassidix</i> (Temminck, 1823)	Generic change
<i>Aceros corrugatus</i> (Temminck, 1832)	II/B	=	<i>Rhyticeros corrugatus</i> (Temminck, 1832)	Generic change
<i>Aceros leucocephalus</i> (Vieillot, 1816)	II/B	=	<i>Rhyticeros leucocephalus</i> (Vieillot, 1816)	Generic change
<i>Aceros waldeni</i> (Sharpe, 1877)	II/B	=	<i>Rhyticeros waldeni</i> (Sharpe, 1877)	Generic change
<i>Anorrhinus tickelli</i> (Blyth, 1855)	II/B	>	<i>Ptilolaemus tickelli</i> (Blyth, 1855)	Generic change
			<i>Ptilolaemus austeni</i> (Blyth, 1855)	Generic change & species split
CUCULIFORMES				
MUSOPHAGIDAE				
<i>Tauraco porphyreolophus</i> (Vigors, 1831)	II/B	=	<i>Gallirex porphyreolophus</i> (Vigors, 1831)	Generic change. Current listing of <i>Tauraco</i> spp. cannot be changed to <i>Tauraco</i> spp. and <i>Gallirex</i> spp. because other species of <i>Gallirex</i> are not CITES listed.
FALCONIFORMES				
ACCIPITRIDAE				
<i>Accipiter cirrhocephalus</i> (Vieillot, 1817)	II/B	=	<i>Accipiter cirrocephalus</i> (Vieillot, 1817)	Spelling correction

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Accipiter francesii</i> A. Smith, 1834	II/B	=	<i>Accipiter francesiae</i> A. Smith, 1834	Spelling correction
<i>Accipiter novaehollandiae</i> (J. F. Gmelin, 1788)	II/B	>	<i>Accipiter novaehollandiae</i> (J. F. Gmelin, 1788)	
			<i>Accipiter hiogaster</i> (S. Müller, 1841) (including ssp. <i>hiogaster</i> , <i>albiventris</i> , <i>bougainvillei</i> , <i>dampieri</i> , <i>griseogularis</i> , <i>lavongai</i> , <i>leucosomus</i> , <i>lihirensis</i> , <i>malaitae</i> , <i>manusi</i> , <i>matthiae</i> , <i>misoriensis</i> , <i>misulae</i> , <i>mortyi</i> , <i>obiensis</i> , <i>pallidiceps</i> , <i>pallidimas</i> , <i>polionotus</i> , <i>pulchellus</i> , <i>rubianae</i> , <i>rufoschistaceus</i> , <i>sylvestris</i>)	Species split
<i>Accipiter toussenelii</i> (J. & E. Verreaux, 1855)	II/B	<	<i>Accipiter tachiro</i> (Daudin, 1800)	Species lump
<i>Asturina nitida</i> (Latham, 1790)	II/B	>	<i>Buteo nitidus</i> (Latham, 1790) (including ssp. <i>nitidus</i> , <i>blakei</i> (syn. <i>costaricensis</i>), <i>pallidus</i>)	Generic change
			<i>Buteo plagiatus</i> (Schlegel, 1862)	Generic change and species split
<i>Aquila clanga</i> Pallas, 1811	II/A	=	<i>Clanga clanga</i> (Pallas, 1811)	Generic change
<i>Aquila pomarina</i> C. L. Brehm, 1831	II/A	>	<i>Clanga pomarina</i> (C. L. Brehm, 1831)	Generic change
			<i>Clanga hastata</i> (Lesson, 1831)	Generic change and species split
<i>Buteo albicaudatus</i> Vieillot, 1816	II/B	=	<i>Geranoaetus albicaudatus</i> (Vieillot, 1816)	Generic change
<i>Buteo buteo</i> (Linnaeus, 1758)	II/A	>	<i>Buteo buteo</i> (Linnaeus, 1758) (including ssp. <i>buteo</i> , <i>harterti</i> , <i>insularum</i> , <i>menetriesi</i> , <i>pojana</i> , <i>vulpinus</i>)	
			<i>Buteo japonicus</i> Temminck & Schlegel, 1844 (including ssp. <i>japonicus</i> , <i>oshiroi</i> , <i>toyoshimai</i>)	Species split
			<i>Buteo refectus</i> Portenko, 1935	Species split
<i>Buteo buteo bannermani</i> Swann, 1919	II/A	=	<i>Buteo rufinus bannermani</i> Swann, 1919	Subspecies transfer
<i>Buteo leucorrhous</i> (Quoy & Gaimard, 1824)	II/B	=	<i>Parabuteo leucorrhous</i> (Quoy & Gaimard, 1824)	Generic change
<i>Buteo magnirostris</i> (J. F. Gmelin, 1788)	II/B	=	<i>Rupornis magnirostris</i> (J. F. Gmelin, 1788)	Generic change
<i>Buteo poecilochrous</i> J. H. Gurney, 1879	II/B	<	<i>Geranoaetus polyosoma</i> (Quoy & Gaimard, 1824)	Generic change & species lump
<i>Buteo polyosoma</i> (Quoy & Gaimard, 1824)	II/B	=	<i>Geranoaetus polyosoma</i> (Quoy & Gaimard, 1824)	Generic change
N/A	II/B		<i>Buteo socotraensis</i> Porter & Kirwan, 2010	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
		<	<i>Buteo rufinus</i> (Cretzschmar, 1829)	Species lump
<i>Buteogallus anthracinus</i> (Deppe, 1830)	II/B	>	<i>Buteogallus anthracinus</i> (Deppe, 1830) (including ssp. <i>anthracinus</i> , <i>utilis</i>)	
			<i>Buteogallus gundlachi</i> (Cabanis, 1855)	Species split
<i>Buteogallus subtilis</i> (Thayer & Bangs, 1905)	II/B	<	<i>Buteogallus anthracinus</i> (Deppe, 1830)	Species lump

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Circus maillardi</i> J. Verreaux, 1862	II/B	>	<i>Circus maillardi</i> J. Verreaux, 1862	
			<i>Circus macroscelus</i> A. Newton, 1863	Species split
<i>Gyps rueppellii</i> (A. E. Brehm, 1852)	II/B	=	<i>Gyps rueppelli</i> (A. E. Brehm, 1852)	Spelling correction. See H&M 3 corrigendum 2.1R for spelling as 'rueppellii'
<i>Harpyhaliaetus coronatus</i> (Vieillot, 1817)	II/B	=	<i>Buteogallus coronatus</i> (Vieillot, 1817)	Generic change
<i>Harpyhaliaetus solitarius</i> (Tschudi, 1844)	II/B	=	<i>Buteogallus solitarius</i> (Tschudi, 1844)	Generic change
<i>Hieraaetus fasciatus</i> (Vieillot, 1822)	II/A	=	<i>Aquila fasciata</i> Vieillot, 1822	Generic change
<i>Hieraaetus kienerii</i> (I. Geoffroy Saint-Hilaire, 1835)	II/B	=	<i>Lophotriorchis kienerii</i> (de Sparre, 1835)	Generic change and author correction
<i>Hieraaetus morphnoides</i> (Gould, 1841)	II/B	>	<i>Hieraaetus morphnoides</i> (Gould, 1841)	
			<i>Hieraaetus weiskei</i> (Reichenow, 1900)	Species split
<i>Hieraaetus spilogaster</i> (Bonaparte, 1850)	II/B	=	<i>Aquila spilogaster</i> (Bonaparte, 1850)	Generic change
<i>Ichthyophaga humilis</i> (Müller & Schlegel, 1841)	II/B	=	<i>Ichthyophaga humilis</i> (Müller & Schlegel, 1841)	Spelling correction
<i>Ichthyophaga ichthyaetus</i> (Horsfield, 1821)	II/B	=	<i>Ichthyophaga ichthyaetus</i> (Horsfield, 1821)	Spelling correction
<i>Ictinaetus malayensis</i> (Temminck, 1822)	II/B	=	<i>Ictinaetus malaiensis</i> (Temminck, 1822)	Spelling correction
<i>Leptodon cayanensis</i> (Latham, 1790)	II/B	>	<i>Leptodon cayanensis</i> (Latham, 1790) (including ssp. <i>cayanensis</i> , <i>monachus</i>)	
			<i>Leptodon forbesi</i> (Swann, 1922)	Species split
<i>Leucopternis albicollis</i> (Latham, 1790)	II/B	=	<i>Pseudastur albicollis</i> (Latham, 1790)	Generic change
<i>Leucopternis lacernulatus</i> (Temminck, 1827)	II/B	=	<i>Buteogallus lacernulatus</i> (Temminck, 1827)	Generic change
<i>Leucopternis occidentalis</i> Salvin, 1876	II/A	=	<i>Pseudastur occidentalis</i> (Salvin, 1876)	Generic change
<i>Leucopternis plumbeus</i> Salvin, 1872	II/B	=	<i>Cryptoleucopteryx plumbea</i> (Salvin, 1872)	Generic change
<i>Leucopternis polionotus</i> (Kaup, 1847)	II/B	=	<i>Pseudastur polionotus</i> (Kaup, 1847)	Generic change
<i>Leucopternis princeps</i> P. L. Sclater, 1865	II/B	=	<i>Morphnarchus princeps</i> (P. L. Sclater, 1865)	Generic change
<i>Leucopternis schistaceus</i> (Sundevall, 1851)	II/B	=	<i>Buteogallus schistaceus</i> (Sundevall, 1850)	Generic change and date correction
<i>Oroaetus isidori</i> (Des Murs, 1845)	II/B	=	<i>Spizaetus isidori</i> (Des Murs, 1845)	Generic change
<i>Pernis celebensis</i> Wallace, 1868	II/B	>	<i>Pernis celebensis</i> Wallace, 1868	
			<i>Pernis steerei</i> W. L. Sclater, 1919 (including ssp. <i>steerei</i> , <i>winkleri</i>)	Species split
<i>Rostrhamus hamatus</i> (Temminck, 1821)	II/B	=	<i>Helicolestes hamatus</i> (Temminck, 1821)	Generic change

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Spizaetus africanus</i> (Cassin, 1865)	II/B	=	<i>Aquila africana</i> (Cassin, 1865)	Generic change
<i>Spizaetus alboniger</i> (Blyth, 1845)	II/B	=	<i>Nisaetus alboniger</i> Blyth, 1845	Generic change
<i>Spizaetus bartelsi</i> Stresemann, 1924	II/B	=	<i>Nisaetus bartelsi</i> (Stresemann, 1924)	Generic change
<i>Spizaetus cirrhatus</i> (J. F. Gmelin, 1788)	II/B	=	<i>Nisaetus cirrhatus</i> (J. F. Gmelin, 1788)	Generic change
<i>Spizaetus lanceolatus</i> Temminck & Schlegel, 1844	II/B	=	<i>Nisaetus lanceolatus</i> (Temminck & Schlegel, 1844)	Generic change
<i>Spizaetus nanus</i> Wallace, 1868	II/B	=	<i>Nisaetus nanus</i> (Wallace, 1868)	Generic change
<i>Spizaetus nipalensis</i> (Hodgson, 1836)	II/B	=	<i>Nisaetus nipalensis</i> Hodgson, 1836	Generic change
<i>Spizaetus philippensis</i> Gould, 1863	II/B	>	<i>Nisaetus philippensis</i> (Gould, 1863)	Generic change
			<i>Nisaetus pinskeri</i> (Preleuthner & Gamauf, 1998)	Generic change and species split
<i>Spizastur melanoleucus</i> (Vieillot, 1816)	II/B	=	<i>Spizaetus melanoleucus</i> (Vieillot, 1816)	Generic change
<i>Torgos tracheliotus</i> (J. R. Forster, 1791)	II/B	=	<i>Torgos tracheliotos</i> (J. R. Forster, 1796)	Spelling and date correction
FALCONIDAE				
<i>Falco pelegrinoides</i> Temminck, 1829	I/A	<	<i>Falco peregrinus</i> Tunstall, 1771	Species lump
N/A	II/B		<i>Micrastur mintoni</i> Whittaker, 2003	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Milvago chimango</i> (Vieillot, 1816)	II/B	=	<i>Phalcoboenus chimango</i> (Vieillot, 1816)	Generic change
GALLIFORMES				
PHASIANIDAE				
<i>Lophura hatinhensis</i> Vo Quy, 1975	-/B	<	<i>Lophura edwardsi</i> (Oustalet, 1896)	Species lump
<i>Polyplectron bicalcaratum</i> (Linnaeus, 1758)	II/B	>	<i>Polyplectron bicalcaratum</i> (Linnaeus, 1758) (now considered monotypic)	
			<i>Polyplectron katsumatae</i> Rothschild, 1906	Species split
GRUIFORMES				
GRUIDAE				
<i>Anthropoides paradiseus</i> (A. A. H. Lichtenstein, 1793)	II/B	=	<i>Grus paradisea</i> (A. A. H. Lichtenstein, 1793)	Generic change
<i>Anthropoides virgo</i> (Linnaeus, 1758)	II/B	=	<i>Grus virgo</i> (Linnaeus, 1758)	Generic change
<i>Bugeranus carunculatus</i> (J. F. Gmelin, 1789)	II/B	=	<i>Grus carunculata</i> (J. F. Gmelin, 1789)	Generic change
<i>Grus antigone</i> (Linnaeus, 1758)	II/B	=	<i>Antigone antigone</i> (Linnaeus, 1758)	Generic change

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Grus canadensis</i> (Linnaeus, 1758)	II/A	=	<i>Antigone canadensis</i> (Linnaeus, 1758)	Generic change
<i>Grus leucogeranus</i> Pallas, 1773	I/A	=	<i>Leucogeranus leucogeranus</i> (Pallas, 1773)	Generic change
<i>Grus rubicunda</i> (Perry, 1810)	II/B	=	<i>Antigone rubicunda</i> (Perry, 1810)	Generic change
<i>Grus vipio</i> Pallas, 1811	I/A	=	<i>Antigone vipio</i> (Pallas, 1811)	Generic change
OTIDIDAE				
<i>Eupodotis humilis</i> (Blyth, 1856)	II/B	=	<i>Heterotetrax humilis</i> (Blyth, 1855)	Generic change and date correction
<i>Eupodotis rueppellii</i> (Wahlberg, 1856)	II/B	=	<i>Heterotetrax rueppellii</i> (Wahlberg, 1856)	Generic change and spelling correction
<i>Eupodotis vigorsii</i> (A. Smith, 1831)	II/B	=	<i>Heterotetrax vigorsii</i> (A. Smith, 1831)	Generic change
<i>Neotis denhami</i> (Children, 1826)	II/B	=	<i>Ardeotis denhami</i> (Children & Vigors, 1826)	Generic change and author correction
<i>Neotis heuglinii</i> (Hartlaub, 1859)	II/B	=	<i>Ardeotis heuglinii</i> (Hartlaub, 1859)	Generic change
<i>Neotis ludwigii</i> (Rüppell, 1837)	II/B	=	<i>Ardeotis ludwigii</i> (Rüppell, 1837)	Generic change
<i>Neotis nuba</i> (Cretzschmar, 1826)	II/B	=	<i>Ardeotis nuba</i> (Cretzschmar, 1826)	Generic change
RALLIDAE				
<i>Gallirallus sylvestris</i> (P. L. Sclater, 1870)	I/A	=	<i>Hypotaenidia sylvestris</i> (P. L. Sclater, 1870)	Generic change
PICIFORMES				
RAMPHASTIDAE				
<i>Baillonioides bailloni</i> (Vieillot, 1819)	III/C	=	<i>Pteroglossus bailloni</i> (Vieillot, 1819)	Generic change
PSITTACIFORMES				
CACATUIDAE				
<i>Cacatua goffini</i> (Finsch, 1863)	I/A	=	<i>Cacatua goffiniana</i> Roselaar & Michels, 2004	Replacement name. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Calyptorhynchus baudinii</i> Lear, 1832	II/B	=	<i>Zanda baudinii</i> (Lear, 1832)	Generic change
<i>Calyptorhynchus funereus</i> (Shaw, 1794)	II/B	=	<i>Zanda funerea</i> (Shaw, 1794)	Generic change
<i>Calyptorhynchus latirostris</i> Camaby, 1948	II/B	=	<i>Zanda latirostris</i> (Camaby, 1948)	Generic change

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
LORIIDAE				
<i>Eos rubra</i> (J. F. Gmelin, 1788)	II/B	=	<i>Eos bornea</i> (Linnaeus, 1758)	Replacement name
PSITTACIDAE				
<i>Amazona mercenaria</i> (Tschudi, 1844)	II/B	=	<i>Amazona mercenarius</i> (Tschudi, 1844)	Spelling correction
<i>Amazona xanthops</i> (Spix, 1824)	II/B	=	<i>Alipiopsitta xanthops</i> (von Spix, 1824)	Generic change and author correction
<i>Aratinga acuticaudata</i> (Vieillot, 1818)	II/B	=	<i>Psittacara acuticaudatus</i> (Vieillot, 1818)	Generic change
<i>Aratinga aurea</i> (J. F. Gmelin, 1788)	II/B	=	<i>Eupsittula aurea</i> (J. F. Gmelin, 1788)	Generic change
<i>Aratinga cactorum</i> (Kuhl, 1820)	II/B	=	<i>Eupsittula cactorum</i> (Kuhl, 1820)	Generic change
<i>Aratinga canicularis</i> (Linnaeus, 1758)	II/B	=	<i>Eupsittula canicularis</i> (Linnaeus, 1758)	Generic change
<i>Aratinga chloroptera</i> (Souancé, 1856)	II/B	=	<i>Psittacara chloropterus</i> (Souancé, 1856)	Generic change
<i>Aratinga erythrogenys</i> (Lesson, 1844)	II/B	=	<i>Psittacara erythrogenys</i> (Lesson, 1844)	Generic change
<i>Aratinga euops</i> (Wagler, 1832)	II/B	=	<i>Psittacara euops</i> (Wagler, 1832)	Generic change
<i>Aratinga finschi</i> (Salvin, 1871)	II/B	=	<i>Psittacara finschi</i> (Salvin, 1871)	Generic change
<i>Aratinga holochlora</i> (P. L. Sclater, 1859)	II/B	=	<i>Psittacara holochlorus</i> (P. L. Sclater, 1859)	Generic change
<i>Aratinga leucophthalma</i> (Stadius Müller, 1776)	II/B	=	<i>Psittacara leucophthalmus</i> (Stadius Müller, 1776)	Generic change
<i>Aratinga mitrata</i> (Tschudi, 1844)	II/B	=	<i>Psittacara mitratus</i> (Tschudi, 1844)	Generic change
<i>Aratinga nana</i> (Vigors, 1830)	II/B	=	<i>Eupsittula nana</i> (Vigors, 1830)	Generic change
<i>Aratinga pertinax</i> (Linnaeus, 1758)	II/B	=	<i>Eupsittula pertinax</i> (Linnaeus, 1758)	Generic change
<i>Aratinga solstitialis</i> (Linnaeus, 1758)	II/B	>	<i>Aratinga solstitialis</i> (Linnaeus, 1758)	
			<i>Aratinga maculata</i> (Stadius Müller, 1776)	Species split. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Aratinga strenua</i> (Ridgway, 1915)	II/B	=	<i>Psittacara strenuus</i> (Ridgway, 1915)	Generic change
<i>Aratinga wagleri</i> (G. R. Gray, 1845)	II/B	=	<i>Psittacara wagleri</i> (G. R. Gray, 1845)	Generic change

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Coracopsis nigra</i> (Linnaeus, 1758)	II/B	>	<i>Coracopsis nigra</i> (Linnaeus, 1758) (including ssp. <i>nigra</i> , <i>libs</i> , <i>sibilans</i>)	
			<i>Coracopsis barklyi</i> E. Newton, 1867	Species split
<i>Cyclopsitta gulelmitertii</i> (Schlegel, 1866)	II/B	>	<i>Cyclopsitta gulelmitertii</i> (Schlegel, 1866)	
			<i>Cyclopsitta melanogenia</i> (von Rosenberg, 1866) (including ssp. <i>melanogenia</i> , <i>fuscifrons</i> , <i>suavissima</i>)	Species split
			<i>Cyclopsitta nigrifrons</i> (Reichenow, 1891) (including ssp. <i>nigrifrons</i> , <i>amabilis</i> , <i>ramuensis</i>)	Species split
<i>Eunymphicus cornutus</i> (J. F. Gmelin, 1788)	I/A	>	<i>Eunymphicus cornutus</i> (J. F. Gmelin, 1788)	
			<i>Eunymphicus uvaeensis</i> (E. L. & E. L. C. Layard, 1882)	Species split
<i>Forpus sclateri</i> (G. R. Gray, 1859)	II/B	=	<i>Forpus modestus</i> (Cabanis, 1849)	Replacement name. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Guarouba guarouba</i> (J. F. Gmelin, 1788)	I/A	=	<i>Guaruba guarouba</i> (J. F. Gmelin, 1788)	Spelling correction
<i>Loriculus aurantiifrons</i> Schlegel, 1871	II/B	>	<i>Loriculus aurantiifrons</i> Schlegel, 1871 (including ssp. <i>aurantiifrons</i> , <i>batavorum</i> , <i>meekei</i>)	
			<i>Loriculus tener</i> P. L. Sclater, 1877	Species split
<i>Loriculus amabilis</i> Wallace, 1862	II/B	>	<i>Loriculus amabilis</i> Wallace, 1862	
			<i>Loriculus sclateri</i> Wallace, 1863 (including ssp. <i>sclateri</i> , <i>ruber</i>)	Species split
<i>Nandayus nenday</i> (Vieillot, 1823)	II/B	=	<i>Aratinga nenday</i> (Vieillot, 1823)	Generic change
<i>Orthopsittaca manilata</i> (Boddaert, 1783)	II/B	=	<i>Orthopsittaca manilatus</i> (Boddaert, 1783)	Spelling correction
<i>Pezoporus wallicus</i> (Kerr, 1792)	I/A	>	<i>Pezoporus wallicus</i> (Kerr, 1792) (including ssp. <i>wallicus</i> , <i>leachi</i>)	
			<i>Pezoporus flaviventris</i> North, 1911	Species split
N/A	II/B		<i>Pionopsitta aurantiocephala</i> Gaban-Lima, Raposo & Höfling, 2002	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
		=	<i>Pyrilia aurantiocephala</i> (Gaban-Lima, Raposo & Höfling, 2002)	Generic change
<i>Pionopsitta barrabandi</i> (Kuhl, 1820)	II/B	=	<i>Pyrilia barrabandi</i> (Kuhl, 1820)	Generic change
<i>Pionopsitta caica</i> (Latham, 1790)	II/B	=	<i>Pyrilia caica</i> (Latham, 1790)	Generic change
<i>Pionopsitta haematotis</i> (Sclater & Salvin, 1860)	II/B	=	<i>Pyrilia haematotis</i> (Sclater & Salvin, 1860)	Generic change
<i>Pionopsitta pulchra</i> Berlepsch, 1897	II/B	=	<i>Pyrilia pulchra</i> (von Berlepsch, 1897)	Generic change and author correction

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Pionopsitta pyrilia</i> (Bonaparte, 1853)	II/B	=	<i>Pyrilia pyrilia</i> (Bonaparte, 1853)	Generic change
<i>Pionopsitta vulturina</i> (Kuhl, 1820)	II/B	=	<i>Pyrilia vulturina</i> (Kuhl, 1820)	Generic change
<i>Prioniturus montanus</i> Ogilvie-Grant, 1895	II/B	>	<i>Prioniturus montanus</i> Ogilvie-Grant, 1895	
			<i>Prioniturus waterstradti</i> Rothschild, 1904 (including ssp. <i>waterstradti</i> , <i>malindangensis</i>)	Species split
<i>Prosopeia personata</i> (G. R. Gray, 1848)	II/B	=	<i>Pyrrhulopsis personatus</i> (G. R. Gray, 1848)	Generic change
<i>Prosopeia splendens</i> (Peale, 1848)	II/B	=	<i>Pyrrhulopsis splendens</i> (Peale, 1848)	Generic change
<i>Prosopeia tabuensis</i> (J. F. Gmelin, 1788)	II/B	=	<i>Pyrrhulopsis tabuensis</i> (J. F. Gmelin, 1788)	Generic change
<i>Psephotus chryspterygius</i> Gould, 1858	I/A	=	<i>Psephotellus chryspterygius</i> (Gould, 1857)	Generic change and date correction
<i>Psephotus dissimilis</i> Collett, 1898	I/A	=	<i>Psephotellus dissimilis</i> (Collett, 1898)	Generic change
<i>Psephotus pulcherrimus</i> (Gould, 1845)	I/A	=	<i>Psephotellus pulcherrimus</i> (Gould, 1845)	Generic change
<i>Psephotus varius</i> A. H. Clark, 1910	II/B	=	<i>Psephotellus varius</i> (A. H. Clark, 1910)	Generic change
<i>Psittacula calthorpa</i> (Blyth, 1849)	II/B	=	<i>Psittacula calthrapae</i> (Blyth, 1849)	Spelling correction
<i>Psittacula echo</i> (A. & E. Newton, 1876)	I/A	<	<i>Psittacula eques</i> (Boddaert, 1783)	Species lump
<i>Pyrrhura caeruleiceps</i> Todd, 1947	II/B	<	<i>Pyrrhura picta</i> (Stadius Müller, 1776)	Species lump
<i>Pyrrhura eisenmanni</i> Delgado, 1985	II/B	<	<i>Pyrrhura picta</i> (Stadius Müller, 1776)	Species lump
<i>Pyrrhura leucotis</i> (Kuhl, 1820)	II/B	>	<i>Pyrrhura leucotis</i> (Kuhl, 1820) (including ssp. <i>leucotis</i> , <i>emma</i>)	
			<i>Pyrrhura griseipectus</i> Salvadori, 1900	Species split
N/A	II/B		<i>Pyrrhura parvifrons</i> Arndt, 2008	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
		<	<i>Pyrrhura roseifrons</i> (G. R. Gray, 1859)	Species lump
<i>Pyrrhura picta</i> (Stadius Müller, 1776)	II/B	>	<i>Pyrrhura picta</i> (Stadius Müller, 1776) (including ssp. <i>picta</i> , <i>caeruleiceps</i> , <i>eisenmanni</i> , <i>subandina</i>)	
			<i>Pyrrhura amazonum</i> Hellmayr, 1906 (including ssp. <i>amazonum</i> , <i>lucida</i> Arndt, 2008, <i>snethlageae</i> Joseph & Bates, 2002)	Species split
			<i>Pyrrhura lucianii</i> (Deville, 1851)	Species split
			<i>Pyrrhura roseifrons</i> (G. R. Gray, 1859) (including ssp. <i>roseifrons</i> , <i>dilutissima</i> Arndt, 2008, <i>parvifrons</i> Arndt, 2008, <i>peruviana</i>)	Species split
<i>Pyrrhura subandina</i> Todd, 1917	II/B	<	<i>Pyrrhura picta</i> (Stadius Müller, 1776)	Species lump
RHEIFORMES				

Dickinson (2003)	CITES Appendix/ EU Annex	Dickinson & Remsen (2013)	Notes
RHEIDAE			
<i>Pterocnemia pennata</i> (d'Orbigny, 1834)	I/A	= <i>Rhea pennata</i> d'Orbigny, 1834	Generic change
STRIGIFORMES			
STRIGIDAE			
<i>Bubo bubo</i> (Linnaeus, 1758)	II/A	> <i>Bubo bubo</i> (Linnaeus, 1758) (including ssp. <i>bubo</i> , <i>borissowi</i> , <i>hemachalanus</i> , <i>hispanus</i> , <i>interpositus</i> , <i>jakutensis</i> , <i>kiautschensis</i> , <i>nikolskii</i> , <i>omissus</i> , <i>ruthenus</i> , <i>sibiricus</i> , <i>tarimensis</i> , <i>tibetanus</i> , <i>turcomanus</i> , <i>ussuriensis</i> , <i>yenisseensis</i>)	
		<i>Bubo bengalensis</i> (Franklin, 1831)	Species split
<i>Bubo cinerascens</i> Guérin-Méneville, 1843	II/B	< <i>Bubo africanus</i> (Temminck, 1821)	Species lump
<i>Bubo vosseleri</i> Reichenow, 1908	II/B	< <i>Bubo poensis</i> Fraser, 1854	Species lump
<i>Glaucidium castanonotum</i> (Blyth, 1852)	II/B	= <i>Glaucidium castanonotum</i> (Blyth, 1852)	Spelling correction
N/A	II/B	<i>Glaucidium mooreorum</i> Silva, Coelho & Gonzaga, 2003	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Gymnoglaux lawrencii</i> Sclater & Salvin, 1868	II/B	= <i>Margarobyas lawrencii</i> (Sclater & Salvin, 1868)	Generic change
<i>Ninox affinis isolata</i> E. C. S. Baker, 1926	II/B	= <i>Ninox scutulata isolata</i> E. C. S. Baker, 1926	Subspecies transfer
<i>Ninox affinis rexpimenti</i> Abdulali, 1979	II/B	= <i>Ninox scutulata rexpimenti</i> Abdulali, 1979	Subspecies transfer
<i>Ninox boobook</i> (Latham, 1802)	II/B	> <i>Ninox boobook</i> (Latham, 1802) (including ssp. <i>boobook</i> , <i>cinnamomina</i> , <i>fusca</i> , <i>halmaturina</i> , <i>moae</i> , <i>ocellata</i> , <i>plesseni</i> , <i>pusilla</i> , <i>remigialis</i> , <i>rotiensis</i> ,)	
		<i>Ninox lurida</i> De Vis, 1887	Species split
N/A	II/B	<i>Ninox burhani</i> Indrawan & Somadikarta, 2004	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
N/A		<i>Ninox leventisi</i> Rasmussen, Allen, Collar, DeMeulemeester, Hutchinson, Jakosalem, Kennedy, Lambert & Paguntalan, 2012	New species
<i>Ninox novaeseelandiae</i> (J. F. Gmelin, 1788)	II/B	> <i>Ninox novaeseelandiae</i> (J. F. Gmelin, 1788) (including ssp. <i>novaeseelandiae</i> , <i>albaria</i> , <i>undulata</i>)	
		<i>Ninox leucopsis</i> (Gould, 1838)	Species split
<i>Ninox philippensis</i> Bonaparte, 1855	II/B	> <i>Ninox philippensis</i> Bonaparte, 1855 (including ssp. <i>philippensis</i> [syn. <i>proxima</i>], <i>centralis</i> , <i>ticaoensis</i>)	
		<i>Ninox mindorensis</i> Ogilvie-Grant, 1896	Species split
		<i>Ninox reyi</i> Oustalet, 1880	Species split

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
			<i>Ninox spilocephala</i> Tweeddale, 1879	Species split
			<i>Ninox spilonotus</i> Bourns & Worcester, 1894 (including ssp. <i>spilonotus</i> , <i>fisheri</i> Rasmussen, Allen, Collar, DeMeulemeester, Hutchinson, Jakosalem, Kennedy, Lambert & Paguntalan, 2012)	Species split
N/A			<i>Ninox rumseyi</i> Rasmussen, Allen, Collar, DeMeulemeester, Hutchinson, Jakosalem, Kennedy, Lambert & Paguntalan, 2012	New species
<i>Ninox scutulata</i> (Raffles, 1822)	II/B	>	<i>Ninox scutulata</i> (Raffles, 1822) (including ssp. <i>scutulata</i> , <i>borneensis</i> , <i>burmanica</i> , <i>hirsuta</i> , <i>isolata</i> , <i>javanensis</i> , <i>lugubris</i> , <i>obscura</i> , <i>palawanensis</i> , <i>rexpimenti</i>)	
			<i>Ninox japonica</i> (Temminck & Schlegel, 1845) (including ssp. <i>japonica</i> , <i>florensis</i> , <i>totogo</i>)	Species split
			<i>Ninox randi</i> Deignan, 1951	Species split
<i>Ninox squamipila</i> (Bonaparte, 1850)	II/B	>	<i>Ninox squamipila</i> (Bonaparte, 1850) (including ssp. <i>squamipila</i> , <i>hantu</i>)	
			<i>Ninox forbesi</i> P. L. Sclater, 1883	Species split
			<i>Ninox hypogramma</i> (G. R. Gray, 1861)	Species split
<i>Ninox supercilii</i> (Vieillot, 1817)	II/B	=	<i>Athene supercilii</i> (Vieillot, 1817)	Generic change
<i>Nyctea scandiaca</i> (Linnaeus, 1758)	II/A	=	<i>Bubo scandiacus</i> (Linnaeus, 1758)	Generic change
<i>Otus albogularis</i> (Cassin, 1849)	II/B	=	<i>Megascops albogularis</i> (Cassin, 1849)	Generic change
<i>Otus asio</i> (Linnaeus, 1758)	II/B	=	<i>Megascops asio</i> (Linnaeus, 1758)	Generic change
<i>Otus atricapilla</i> (Temminck, 1822)	II/B	=	<i>Megascops atricapilla</i> (Temminck, 1822)	Generic change
<i>Otus barbarus</i> (Sclater & Salvin, 1868)	II/B	=	<i>Megascops barbarus</i> (Sclater & Salvin, 1868)	Generic change
<i>Otus clarkii</i> L. & E. H. Kelso, 1935	II/B	=	<i>Megascops clarkii</i> (L. & E. H. Kelso, 1935)	Generic change
<i>Otus choliba</i> (Vieillot, 1817)	II/B	=	<i>Megascops choliba</i> (Vieillot, 1817)	Generic change
<i>Otus colombianus</i> Traylor, 1952	II/B	=	<i>Megascops colombianus</i> (Traylor, 1952)	Generic change
<i>Otus cooperi</i> (Ridgway, 1878)	II/B	=	<i>Megascops cooperi</i> (Ridgway, 1878)	Generic change
<i>Otus guatemalae</i> (Sharpe, 1875)	II/B	=	<i>Megascops guatemalae</i> (Sharpe, 1875)	Generic change
<i>Otus hoyi</i> C. König & Straneck, 1989	II/B	=	<i>Megascops hoyi</i> (C. König & Straneck, 1989)	Generic change
<i>Otus ingens</i> (Salvin, 1897)	II/B	=	<i>Megascops ingens</i> (Salvin, 1897)	Generic change
<i>Otus kennicottii</i> (Elliot, 1867)	II/B	=	<i>Megascops kennicottii</i> (Elliot, 1867)	Generic change
<i>Otus koepckeae</i> Hekstra, 1982	II/B	=	<i>Megascops koepckeae</i> (Hekstra, 1982)	Generic change
<i>Otus madagascariensis</i> (A. Grandidier, 1867)	II/B	<	<i>Otus rutilus</i> (Pucheran, 1849)	Species lump
<i>Otus magicus siaoensis</i> (Schlegel, 1873)	II/B	=	<i>Otus manadensis siaoensis</i> (Schlegel, 1873)	Subspecies transfer

Dickinson (2003)	CITES Appendix/ EU Annex		Dickinson & Remsen (2013)	Notes
<i>Otus marshalli</i> Weske & Terborgh, 1981	II/B	=	<i>Megascops marshalli</i> (Weske & Terborgh, 1981)	Generic change
<i>Otus nudipes</i> (Daudin, 1800)	II/B	=	<i>Megascops nudipes</i> (Daudin, 1800)	Generic change
<i>Otus petersoni</i> Fitzpatrick & O'Neill, 1986	II/B	=	<i>Megascops petersoni</i> (Fitzpatrick & O'Neill, 1986)	Generic change
<i>Otus roboratus</i> Bangs & Noble, 1918	II/B	=	<i>Megascops roboratus</i> (Bangs & Noble, 1918)	Generic change
<i>Otus sanctaecatarinae</i> (Salvin, 1897)	II/B	=	<i>Megascops sanctaecatarinae</i> (Salvin, 1897)	Generic change
<i>Otus seductus</i> R. T. Moore, 1941	II/B	=	<i>Megascops seductus</i> (R. T. Moore, 1941)	Generic change
N/A	II/B		<i>Otus thilohoffmanni</i> Warakagoda & Rasmussen, 2004	New species. Already adopted at CoP16 referring to original publication, see Res. Conf. 12.11 (rev. CoP16).
<i>Otus trichopsis</i> (Wagler, 1832)	II/B	=	<i>Megascops trichopsis</i> (Wagler, 1832)	Generic change
<i>Otus watsonii</i> (Cassin, 1848)	II/B	=	<i>Megascops watsonii</i> (Cassin, 1849)	Generic change and date correction
<i>Pseudoscops clamator</i> (Vieillot, 1808)	II/B	=	<i>Asio clamator</i> (Vieillot, 1808)	Generic change
<i>Strix albitarsis</i> (Bonaparte, 1850)	II/B	=	<i>Ciccaba albitarsis</i> (Bonaparte, 1850)	Generic change
<i>Strix virgata</i> (Cassin, 1849)	II/B	=	<i>Ciccaba virgata</i> (Cassin, 1849)	Generic change
TYTONIDAE				
<i>Phodilus prigoginei</i> Schouteden, 1952	II/B	=	<i>Tyto prigoginei</i> (Schouteden, 1952)	Generic change
<i>Tyto alba</i> (Scopoli, 1769)	II/A	>	<i>Tyto alba</i> (Scopoli, 1769) (including ssp. <i>alba</i> , <i>bargei</i> , <i>bondi</i> , <i>contempta</i> , <i>detorta</i> , <i>erlangeri</i> , <i>ernesti</i> , <i>furcata</i> , <i>gracilirostris</i> , <i>guatemalae</i> , <i>guttata</i> , <i>hypermetra</i> , <i>insularis</i> , <i>javanica</i> , <i>nigrescens</i> , <i>niveicauda</i> , <i>poensis</i> , <i>pratincola</i> , <i>punctatissima</i> , <i>schmitzi</i> , <i>stertens</i> , <i>thomensis</i> , <i>tuidara</i> , <i>zottae</i>)	
			<i>Tyto delicatula</i> (Gould, 1837) (including ssp. <i>delicatula</i> , <i>crassirostris</i> , <i>interposita</i> , <i>meekei</i> , <i>sumbaensis</i>)	Species split
			<i>Tyto deroepstorffi</i> (Hume, 1875)	Species split
<i>Tyto capensis</i> (A. Smith, 1834)	II/B	>	<i>Tyto capensis</i> (A. Smith, 1834) (including ssp. <i>capensis</i> , <i>cameroonensis</i>)	
			<i>Tyto longimembris</i> (Jerdon, 1839) (including ssp. <i>longimembris</i> , <i>amauronota</i> , <i>chinensis</i> [syn. <i>pithecops</i>], <i>papuensis</i>)	Species split
<i>Tyto novaehollandiae</i> (Stephens, 1826)	II/B	>	<i>Tyto novaehollandiae</i> (Stephens, 1826) (including ssp. <i>novaehollandiae</i> , <i>calabyi</i> , <i>galei</i> , <i>kimberli</i> , <i>melvillensis</i>)	
			<i>Tyto castanops</i> (Gould, 1837)	Species split

Species identified until February 2014**within the family Chamaeleonidae and the genus *Phelsuma*****1. Chamaeleonidae**

Archaius tigris (Kuhl, 1820)

Bradypodion atromontanum Branch, Tolley & Tilbury, 2006

Bradypodion caeruleogula Raw & Brothers, 2008

Bradypodion caffer (Boettger, 1889)

Bradypodion damaranum (Boulenger, 1887)

Bradypodion dracomontanum Raw, 1976

Bradypodion gutturale (Smith, 1849)

Bradypodion kentanicum (Hewitt, 1935)

Bradypodion melanocephalum (Gray, 1865)

Bradypodion nemorale Raw, 1978

Bradypodion ngomeense Tilbury & Tolley, 2009

Bradypodion occidentale (Hewitt, 1935)

Bradypodion pumilum (Gmelin, 1789)

Bradypodion setaroi Raw, 1976

Bradypodion taeniabronchum (Smith, 1831)

Bradypodion thamnobates Raw, 1976

Bradypodion transvaalense (Fitzsimons, 1930)

Bradypodion ventrale (Gray, 1845)

Brookesia ambreensis Raxworthy & Nussbaum, 1995

Brookesia antakarana Raxworthy & Nussbaum, 1995

Brookesia bekolosy Raxworthy & Nussbaum, 1995

Brookesia betschi Brygoo, Blanc & Domergue, 1974

Brookesia bonisi Ramanantsoa, 1980

Brookesia brunoi Crottini, Miralles, Glaw, Harris, Lima & Vences, 2012

Brookesia brygooi Raxworthy & Nussbaum, 1995

Brookesia confidens Glaw, Köhler, Townsend & Vences, 2012

Brookesia decaryi Angel, 1938

Brookesia dentata Mocquard, 1900

Brookesia desperata Glaw, Köhler, Townsend & Vences, 2012

Brookesia ebenau (Boettger, 1880)

Brookesia exarmata Schimmenti & Jesu, 1996

Brookesia griveaudi Brygoo, Blanc & Domergue, 1974

Brookesia karchei Brygoo, Blanc & Domergue, 1970

Brookesia lambertoni Brygoo & Domergue, 1970

Brookesia lineata Raxworthy & Nussbaum, 1995

Brookesia micra Glaw, Köhler, Townsend & Vences, 2012

Brookesia minima Boettger, 1893

Brookesia perarmata (Angel, 1933)

Brookesia peyrierasi Brygoo, Blanc & Domergue, 1974

Brookesia ramanantsoai Brygoo & Domergue, 1975

Brookesia stumpffi Boettger, 1894

Brookesia superciliaris (Kuhl, 1820)

Brookesia therezieni Brygoo & Domergue, 1969

Brookesia thieli Brygoo & Domergue, 1969

Brookesia tristis Glaw, Köhler, Townsend & Vences, 2012

Brookesia tuberculata Mocquard, 1894

Brookesia vadoni Brygoo & Domergue, 1968

Brookesia valerieae Raxworthy, 1991

Calumma amber Raxworthy & Nussbaum, 2006

Calumma ambreense (Ramanantsoa, 1974)

Calumma andringitraense (Brygoo, Blanc & Domergue, 1972)

Calumma boettgeri (Boulenger, 1888)

Calumma brevicorne (Günther, 1879)

Calumma capuroni (Brygoo, Blanc & Domergue, 1972)

Calumma crypticum Raxworthy & Nussbaum, 2006

Calumma cucullatum (Gray, 1831)

Calumma fallax (Mocquard, 1900)

Calumma furcifer (Vaillant & Grandidier, 1880)

Calumma gallus (Günther, 1877)

Calumma gastrotaenia (Boulenger, 1888)

Calumma glawi Böhme, 1997

Calumma globifer (Günther, 1879)

Calumma guibei (Hillenius, 1959)

Calumma guillaumeti (Brygoo, Blanc & Domergue, 1974)

Calumma hafahafa Raxworthy & Nussbaum, 2006

Calumma hilleniusi (Brygoo, Blanc & Domergue, 1973)

Calumma jevy Raxworthy & Nussbaum, 2006

Calumma linotum (Müller, 1924)

Calumma malthe (Günther, 1879)

Calumma marojezense (Brygoo, Blanc & Domergue, 1970)

Calumma nasutum (Duméril & Bibron, 1836)

Calumma oshaughnessyi (Günther, 1881)

Calumma parsonii (Cuvier, 1824)

Calumma peltierorum Raxworthy & Nussbaum, 2006

Calumma peyrierasi (Brygoo & Domergue, 1974)

Calumma tarzan Gehring, Pabijan, Ratsoavina, Köhler, Vences & Glaw, 2010

Calumma tsaratananense (Brygoo & Domergue, 1968)

Calumma tsycorne Raxworthy & Nussbaum, 2006

Calumma vatosoa Andreone, Mattioli, Jesu & Randrianirina, 2001

Calumma vencesi Andreone, Mattioli, Jesu & Randrianirina, 2001

Calumma vohibola Gehring, Ratsavina, Vences & Glaw, 2011

Chamaeleo africanus Laurenti, 1768

Chamaeleo anchietae Bocage, 1872

Chamaeleo arabicus Matschie, 1893

Chamaeleo calcaricarens Böhme, 1985

Chamaeleo calyptratus Duméril & Duméril, 1851

Chamaeleo chamaeleon (Linnaeus, 1758)

Chamaeleo dilepis Leach, 1819

Chamaeleo gracilis Hallowell, 1844

Chamaeleo laevigatus Gray, 1863

Chamaeleo monachus Gray, 1865

Chamaeleo namaquensis Smith, 1831

Chamaeleo necasi Ullrich, Krause & Böhme, 2007

Chamaeleo senegalensis Daudin, 1802

Chamaeleo zeylanicus Laurenti, 1768

Furcifer angeli (Brygoo & Domergue, 1968)

Furcifer antimena (Grandidier, 1872)

Furcifer balteatus (Duméril & Bibron, 1851)

Furcifer belalandaensis (Brygoo & Domergue, 1970)

Furcifer bifidus (Brongniart, 1800)

Furcifer campani (Grandidier, 1872)

Furcifer cephalolepis (Günther, 1880)

Furcifer labordi (Grandidier, 1872)

Furcifer lateralis (Gray, 1831)

Furcifer major (Brygoo, 1971)

Furcifer minor (Günther, 1879)

Furcifer nicosiai Jesu, Mattioli & Schimmenti, 1999

Furcifer oustaleti (Mocquard, 1894)

Furcifer pardalis (Cuvier, 1829)

Furcifer petteri (Brygoo & Domergue, 1966)

Furcifer polleni (Peters 1874)

Furcifer rhinoceratus (Gray, 1843)

Furcifer timoni Glaw, Köhler & Vences, 2009

Furcifer tuzetae (Brygoo, Bourgat & Domergue, 1972)

Furcifer verrucosus (Cuvier, 1829)

Furcifer viridis Florio, Ingram, Rakotondravony, Louis & Raxworthy, 2012

Furcifer willsii (Günther, 1890)

Kinyongia adolffriderici (Sternfeld, 1912)

Kinyongia asheorum Necas, Sindaco, Koreny, Kopecna, Malonza & Modry, 2009

Kinyongia boehmei (Lutzmann & Necas, 2002)

Kinyongia carpenteri (Parker, 1929)

Kinyongia excubitor (Barbour, 1911)

Kinyongia fischeri (Reichenow, 1887)

Kinyongia gyrolepis Greenbaum, Tolley, Joma & Kusamba, 2012

Kinyongia magomberae Menegon, Tolley, Jones, Rovero, Marshall & Tilbury, 2009

Kinyongia matschiei (Werner, 1895)

Kinyongia multituberculata (Nieden, 1913)

Kinyongia oxyrhina (Klaver & Böhme, 1988)

Kinyongia tavetana (Steindachner, 1891)

Kinyongia tenue (Matschie, 1892)

Kinyongia uluguruensis (Loveridge, 1957)

Kinyongia uthmoelleri (Müller, 1938)

Kinyongia vanheygeni Necas, 2009

Kinyongia vosseleri (Nieden, 1913)

Kinyongia xenorhina (Boulenger, 1901)

Nadzikambia baylissi Branch & Tolley, 2010

Nadzikambia mlanjensis (Broadley, 1965)

Palleon lolontany (Raxworthy & Nussbaum, 1995)

Palleon nasus (Boulenger, 1887)

Rhampholeon acuminatus Mariaux & Tilbury, 2006

Rhampholeon beraduccii Mariaux & Tilbury, 2006

Rhampholeon boulengeri Steindachner, 1911

Rhampholeon chapmanorum Tilbury, 1992

Rhampholeon gorongosae Broadley, 1971

Rhampholeon marshalli Boulenger, 1906

Rhampholeon moyeri Menegon, Salvidio & Tilbury, 2002

Rhampholeon nchisiensis (Loveridge, 1953)

Rhampholeon platyceps Günther, 1893

Rhampholeon spectrum (Buchholz, 1874)

Rhampholeon spinosus (Matschie, 1892)

Rhampholeon temporalis (Matschie, 1892)

Rhampholeon uluguruensis Tilbury & Emmrich, 1996

Rhampholeon viridis Mariaux & Tilbury, 2006

Rieppeleon brachyurus (Günther, 1893)

Rieppeleon brevicaudatus (Matschie, 1892)

Rieppeleon kerstenii (Peters, 1868)

Rieppeleon robecchii (Boulenger, 1891)

Trioceros affinis (Rüppell, 1845)

Trioceros balebicornutus (Tilbury, 1998)

Trioceros bitaeniatus (Fischer, 1884)

Trioceros camerunensis (Müller, 1909)

Trioceros chapini (De Witte, 1964)

Trioceros conirostratus (Tilbury, 1998)

Trioceros cristatus (Stutchbury, 1837)

Trioceros deremensis (Matschie, 1892)

Trioceros eisentrauti (Mertens, 1968)

Trioceros ellioti (Günther, 1895)

Trioceros feae (Boulenger, 1906)

Trioceros fuelleborni (Tornier, 1900)

Trioceros goetzei (Tornier, 1899)

Trioceros hanangensis Krause & Böhme, 2010

Trioceros harennae (Largen, 1995)

Trioceros hoehnelii (Steindachner, 1891)

Trioceros incornutus (Loveridge, 1932)

Trioceros ituriensis (Schmidt, 1919)

Trioceros jacksonii (Boulenger, 1896)

Trioceros johnstoni (Boulenger, 1901)

Trioceros kinangopensis Stipala, Lutzmann, Malonza, Wilkinson, Godley, Nyamache & Evans, 2012

Trioceros kinetensis (Schmidt, 1943)

Trioceros laterispinis (Loveridge, 1932)

Trioceros marsabitensis (Tilbury, 1991)

Trioceros melleri (Gray, 1865)

Trioceros montium (Buchholz, 1874)

Trioceros narraioica (Necas, Modry & Slapeta, 2003)

Trioceros ntunte (Necas, Modry & Slapeta, 2005)

Trioceros nyirit Stipala, Lutzmann, Malonza, Borghesio, Wilkinson, Godley & Evans, 2011

Trioceros oweni (Gray, 1831)

Trioceros perreti (Klaver & Böhme, 1992)

Trioceros pfefferi (Tornier, 1900)

Trioceros quadricornis (Tornier, 1899)

Trioceros rudis (Boulenger, 1906)

Trioceros schoutedeni (Laurent, 1952)

Trioceros schubotzi (Sternfeld, 1912)

Trioceros serratus (Mertens, 1922)

Trioceros sternfeldi (Rand, 1963)

Trioceros tempeli (Tornier, 1899)

Trioceros tremperi (Necas, 1994)

Trioceros wernerii (Tornier, 1899)

Trioceros wiedersheimi (Nieden, 1910)

2. *Phelsuma* spp.

Phelsuma abbotti Stejneger, 1893

Phelsuma andamanense Blyth, 1861

Phelsuma antanosy Raxworthy & Nussbaum, 1993

Phelsuma astriata Tornier, 1901

Phelsuma barbouri Loveridge, 1942

Phelsuma berghofi Krüger, 1996

Phelsuma borai Glaw, Köhler & Vences, 2009

Phelsuma borbonica Mertens, 1966

Phelsuma breviceps Boettger, 1894

Phelsuma cepediana (Milbert, 1812)

Phelsuma comorensis Boettger, 1913

Phelsuma dorsivittata Mertens, 1964

Phelsuma dubia (Boettger, 1881)

Phelsuma edwardnewtoni Vinson & Vinson, 1969

Phelsuma flavigularis Mertens, 1962

Phelsuma gigas Lienard, 1842

Phelsuma gouldi Crottini, Gehring, Glaw, Harris, Lima & Vences, 2011

Phelsuma grandis Gray, 1870

Phelsuma guentheri Boulenger, 1885
Phelsuma guimbeaui Mertens, 1963
Phelsuma guttata Kaudern, 1922
Phelsuma hielscheri Rösler, 2001
Phelsuma hoeschi Berghof & Trautmann, 2009
Phelsuma inexpectata Mertens, 1966
Phelsuma kely Schönecker, Bach & Glaw, 2004
Phelsuma klemmeri Seipp, 1991
Phelsuma kochi Mertens, 1954
Phelsuma laticauda (Boettger, 1880)
Phelsuma lineata Gray, 1842
Phelsuma madagascariensis Gray, 1831
Phelsuma malamakibo Nussbaum, Raxworthy, Raselimanana & Ramanamanjato, 2000
Phelsuma masohoala Raxworthy & Nussbaum, 1994
Phelsuma modesta Mertens, 1970
Phelsuma mutabilis (Grandidier, 1869)
Phelsuma nigristriata Meier, 1984
Phelsuma ornata Gray, 1825
Phelsuma parkeri Loveridge, 1941
Phelsuma parva Meier, 1983
Phelsuma pasteuri Meier, 1984
Phelsuma pronki Seipp, 1994
Phelsuma pusilla Mertens, 1964
Phelsuma quadriocellata (Peters, 1883)
Phelsuma ravenala Raxworthy, Ingram, Rabibisoa & Pearson, 2007
Phelsuma robertmertensi Meier, 1980
Phelsuma roesleri Glaw, Gehring, Köhler, Franzen & Vences, 2010
Phelsuma rosagularis Vinson & Vinson, 1963
Phelsuma seippi Meier, 1987
Phelsuma serraticauda Mertens, 1963

Phelsuma standingi Methuen & Hewitt, 1913

Phelsuma sundbergi Rendahl, 1939

Phelsuma v-nigra Boettger, 1913

Phelsuma vanheygeni Lerner, 2004

UNEP-WCMC **technical report**

Reptile taxonomy

New species and other proposed taxonomic changes relating to reptile species listed in the EU wildlife trade regulations (which includes all CITES listed species)



Reptile taxonomy: New species and other proposed taxonomic changes relating to reptile species listed in the EU wildlife trade regulations (including CITES listed species)

Prepared for

The European Commission, Directorate General Environment, Directorate E - Global & Regional Challenges, LIFE ENV.E.2. – Global Sustainability, Trade & Multilateral Agreements, Brussels, Belgium

Prepared

January 2014

Copyright

European Commission 2014

Citation

UNEP-WCMC. 2014. Reptile taxonomy: New species and other proposed taxonomic changes relating to reptile species listed in the EU wildlife trade regulations (including CITES listed species). UNEP-WCMC, Cambridge.



The UNEP World Conservation Monitoring Centre (UNEP-WCMC) is the specialist biodiversity assessment of the United Nations Environment Programme, the world's foremost intergovernmental environmental organization. The Centre has been in operation for over 30 years, combining scientific research with policy advice and the development of decision tools.

We are able to provide objective, scientifically rigorous products and services to help decision-makers recognize the value of biodiversity and apply this knowledge to all that they do. To do this, we collate and verify data on biodiversity and ecosystem services that we analyze and interpret in comprehensive assessments, making the results available in appropriate forms for national and international level decision-makers and businesses. To ensure that our work is both sustainable and equitable we seek to build the capacity of partners where needed, so that they can provide the same services at national and regional scales.

The contents of this report do not necessarily reflect the views or policies of UNEP, contributory organisations or editors. The designations employed and the presentations do not imply the expressions of any opinion whatsoever on the part of UNEP, the European Commission or contributory organisations, editors or publishers concerning the legal status of any country, territory, city area or its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of a commercial entity or product in this publication does not imply endorsement by UNEP.

UNEP World Conservation Monitoring Centre (UNEP-WCMC)

219 Huntingdon Road,
Cambridge CB3 0L, UK
Tel: +44 1223 277314
www.unep-wcmc.org

UNEP promotes environmentally sound practices globally and in its own activities. Printing on paper from environmentally sustainable forests and recycled fibre is encouraged.

Introduction

This report provides an overview of recent reptile taxonomic and nomenclature changes proposed in the scientific literature, to inform nomenclature discussions by the CITES Animals Committee.

At CITES CoP 16, a number of new CITES standard references were adopted for fauna (see <http://www.cites.org/eng/res/12/12-11R16.php>). These are also the standard references determining nomenclature in the EU Wildlife Trade Regulations (most recently Commission Regulation (EC) 750/2013). Since these references were compiled, a number of new species have been described and a number of other taxa have been subject to taxonomic revisions (splits and lumps) that may have implications for the names used and taxa included in CITES and in the EU Wildlife Trade Regulations.

The table below contains a list of taxonomic changes identified in recent publications relating to reptiles (except for *Chamaeleonidae* spp. and *Phelsuma* spp.) listed in CITES and in the EU Wildlife Trade Regulations that were not taken into account in the decisions adopted at CITES CoP 16.

With regard to generic changes, it is noted that Hoser (2013b) and Stanley *et al.* (2011) contain conflicting proposals to change the genus of six *Cordylus* species (*C. breyeri*, *C. langi*, *C. mossambicus*, *C. regius*, *C. vandami* and *C. warreni*) and Hoser (2012a) and Reynolds *et al.*, (2014) contain conflicting proposals to change the genus of four *Morelia* species (*M. amethystina*, *M. clastolepis*, *M. nauta* and *M. tracyae*).

Taxon	CITES Appendix/ EU Annex		Proposed change	References	Notes
SAURIA					
CORDYLIDAE					
<i>Cordylus barbertonensis</i> (van Dam, 1921)	II/B	< ¹¹	<i>Smaug warreni</i> (Boulenger, 1908)	Stanley <i>et al.</i> (2011)	Species lump
<i>Cordylus breyeri</i> (Van Dam, 1921)	II/B	= ¹²	<i>Cottonsaurus breyeri</i> (Van Dam, 1921)	Hoser (2013b)	Generic change
<i>Cordylus breyeri</i> (Van Dam, 1921)	II/B	=	<i>Smaug breyeri</i> (Van Dam, 1921)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus campbelli</i> (FitzSimons, 1938)	II/B	=	<i>Namazonurus campbelli</i> (FitzSimons, 1938)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus capensis</i> A. Smith, 1838	II/B	=	<i>Hemicordylus capensis</i> (Smith, 1838)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus cataphractus</i> Boie, 1828	II/B	=	<i>Ouroborus cataphractus</i> (Boie, 1828)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus coeruleopunctatus</i> (Hewitt & Methuen, 1913)	II/B	=	<i>Ninurta coeruleopunctatus</i> (Hewitt & Methuen, 1913)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus depressus</i> (FitzSimons, 1930)	II/B	<	<i>Smaug warreni</i> (Boulenger, 1908)	Stanley <i>et al.</i> (2011)	Species lump
<i>Cordylus fasciatus</i> A. Smith, 1838	II/B	<	<i>Pseudocordylus microlepidotus</i> (Cuvier, 1829)	Stanley <i>et al.</i> (2011)	Species lump
<i>Cordylus giganteus</i> A. Smith, 1844	II/B	=	<i>Smaug giganteus</i> (Smith, 1844)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus jordani</i> (Parker, 1936)	II/B	=	<i>Karusaurus jordani</i> (Parker, 1936)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus langi</i> (Loveridge, 1944)	II/B	=	<i>Ninsaurus langi</i> (Loveridge, 1944)	Hoser (2013b)	Generic change
<i>Cordylus langi</i> (Loveridge, 1944)	II/B	=	<i>Pseudocordylus langi</i> Loveridge, 1944	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus lawrenci</i> (FitzSimons, 1939)	II/B	=	<i>Namazonurus lawrenci</i> (FitzSimons, 1939)	Stanley <i>et al.</i> (2011)	Generic change
N/A	II/B		<i>Cordylus marunguensis</i> Greenbaum, Stanley, Kusamba, Moninga, Goldberg & Bursey, 2012	Greenbaum <i>et al.</i> (2012)	New species
<i>Cordylus melanotus</i> A. Smith, 1838	II/B	=	<i>Pseudocordylus melanotus</i> (Smith, 1838)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus microlepidotus</i> Cuvier, 1829	II/B	=	<i>Pseudocordylus microlepidotus</i> (Cuvier, 1829)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus mossambicus</i> FitzSimons, 1958	II/B	=	<i>Cottonsaurus mossambicus</i> (FitzSimons, 1958)	Hoser (2013b)	Generic change
<i>Cordylus mossambicus</i> FitzSimons, 1958	II/B	=	<i>Smaug mossambicus</i> (FitzSimons, 1958)	Stanley <i>et al.</i> (2011)	Generic change

¹¹ Species lumps (indicated by the symbol "<") refer to taxa currently recognised as separate but that have been grouped together under another name in the associated reference.

¹² The symbol "=" is used to indicate taxonomic or nomenclature changes that do not involve a change in the scope of the taxon in question.

Taxon	CITES Appendix/ EU Annex	Proposed change	References	Notes
<i>Cordylus namaquensis</i> (Methuen & Hewitt, 1914)	II/B	= <i>Namazonurus namaquensis</i> (Methuen & Hewitt, 1914)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus nebulosus</i> (Mouton & Van Wyk, 1995)	II/B	= <i>Hemidactylus nebulosus</i> (Mouton & Van Wyk, 1995)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus peersi</i> (Hewitt, 1932)	II/B	= <i>Namazonurus peersi</i> (Hewitt, 1932)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus polyzonus</i> A. Smith, 1838	II/B	= <i>Karusaurus polyzonus</i> (Smith, 1838)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus pustulatus</i> (Peters, 1862)	II/B	= <i>Namazonurus pustulatus</i> (Peters, 1862)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus regius</i> Broadley, 1962	II/B	= <i>Cottosaurus regius</i> (Broadley, 1962)	Hoser (2013b)	Generic change
<i>Cordylus regius</i> Broadley, 1962	II/B	= <i>Smaug regius</i> (Broadley, 1962)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus spinosus</i> FitzSimons, 1947	II/B	= <i>Pseudocordylus spinosus</i> FitzSimons, 1947	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus subviridis</i> A. Smith, 1838	II/B	= <i>Pseudocordylus subviridis</i> (Smith, 1838)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus tasmani</i> Broadley, 1971	II/B	< <i>Cordylus cordylus</i> (Linnaeus, 1758),	Stanley <i>et al.</i> (2011)	Species lump
<i>Cordylus transvaalensis</i> (FitzSimons, 1943)	II/B	= <i>Pseudocordylus transvaalensis</i> FitzSimons, 1943	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus vandami</i> FitzSimons, 1930	II/B	= <i>Cottosaurus vandami</i> (FitzSimons, 1930)	Hoser (2013b)	Generic change
<i>Cordylus vandami</i> FitzSimons, 1930	II/B	= <i>Smaug vandami</i> (FitzSimons, 1930)	Stanley <i>et al.</i> (2011)	Generic change
<i>Cordylus warreni</i> (Boulenger, 1908)	II/B	= <i>Cottosaurus warreni</i> (Boulenger, 1908)	Hoser (2013b)	Generic change
<i>Cordylus warreni</i> (Boulenger, 1908)	II/B	= <i>Smaug warreni</i> (Boulenger, 1908)	Stanley <i>et al.</i> (2011)	Generic change
GEKKONIDAE				
<i>Hoplodactylus chrysoireticus</i> Robb, 1980	III/C (NZ)	= <i>Woodworthia chrysoireticus</i> (Robb, 1980)	Nielsen <i>et al.</i> (2011)	Generic change
<i>Hoplodactylus cryptozoicus</i> Jewell & Leschen, 2004	III/C (NZ)	= <i>Mokopirirakau cryptozoicus</i> (Jewell & Leschen, 2004)	Nielsen <i>et al.</i> (2011)	Generic change
<i>Hoplodactylus granulatus</i> (Gray, 1845)	III/C (NZ)	= <i>Mokopirirakau granulatus</i> (Gray, 1845)	Nielsen <i>et al.</i> (2011)	Generic change
<i>Hoplodactylus kahutarae</i> Whitaker, 1895	III/C (NZ)	= <i>Mokopirirakau kahutarae</i> (Whitaker, 1895)	Nielsen <i>et al.</i> (2011)	Generic change
<i>Hoplodactylus maculatus</i> (Gray, 1845)	III/C (NZ)	= <i>Woodworthia maculatus</i> (Gray, 1845)	Nielsen <i>et al.</i> (2011)	Generic change
<i>Hoplodactylus nebulosus</i> (McCann, 1955)	III/C (NZ)	= <i>Mokopirirakau nebulosus</i> (McCann, 1955)	Nielsen <i>et al.</i> (2011)	Generic change

Taxon	CITES Appendix/ EU Annex		Proposed change	References	Notes
<i>Hoplodactylus pacificus</i> (Gray, 1842)	III/C (NZ)	> ¹³	<i>Dactylocnemis pacificus</i> (Gray, 1842)	Nielsen <i>et al.</i> (2011)	Generic change
			<i>Woodworthia brunneus</i> (Cope, 1869)	Nielsen <i>et al.</i> (2011)	Species split
<i>Hoplodactylus rakiurae</i> Thomas, 1981	III/C (NZ)	=	<i>Tukutuku rakiurae</i> (Thomas, 1981)	Nielsen <i>et al.</i> (2011)	Generic change
<i>Hoplodactylus stephensi</i> Robb, 1980	III/C (NZ)	=	<i>Toropuku stephensi</i> (Robb, 1980)	Nielsen <i>et al.</i> (2011)	Generic change
TEIIDAE					
<i>Tupinambis duseni</i> Lönnberg, 1896	II/B	=	<i>Salvator duseni</i> Lönnberg, 1910	Harvey <i>et al.</i> (2012)	Generic change
<i>Tupinambis merianae</i> (Duméril & Bibron, 1839)	II/B	=	<i>Salvator merianae</i> (Duméril & Bibron, 1839)	Harvey <i>et al.</i> (2012)	Generic change
<i>Tupinambis rufescens</i> (Günther, 1871)	II/B	=	<i>Salvator rufescens</i> (Günther, 1871)	Harvey <i>et al.</i> (2012)	Generic change
SERPENTES					
BOIDAE					
N/A	I/A		<i>Acrantophis sloppi</i> Hoser, 2013a	Hoser (2013a)	New species
<i>Boa constrictor</i> Linnaeus, 1758	II/B	>	<i>Boa imperator</i> Daidin, 1803	Reynolds <i>et al.</i> (2014)	Species split
<i>Charina bottae</i> (Blainville, 1835)	II/B	>	<i>Charina umbratica</i> Klauber, 1943	Reynolds <i>et al.</i> (2014)	Species split
<i>Chilabothrus angulifer</i> Bibron, 1840	II/B	=	<i>Chilabothrus angulifer</i> (Bibron, 1843)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Corallus annulatus</i> (Cope, 1875)	II/B	>	<i>Corallus blombergi</i> (Rendahl & Vestergren, 1941)	Henderson <i>et al.</i> (2013)	Species split
<i>Epicrates chrysogaster</i> (Cope, 1871)	II/B	=	<i>Chilabothrus chrysogaster</i> (Cope, 1871)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Epicrates exsul</i> Netting & Goin, 1944	II/B	=	<i>Chilabothrus exsul</i> (Netting & Goin, 1944)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Epicrates fordii</i> (Günther, 1861)	II/B	=	<i>Chilabothrus fordii</i> (Günther, 1861)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Epicrates gracilis</i> (Fischer, 1888)	II/B	=	<i>Chilabothrus gracilis</i> (Fischer, 1888)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Epicrates inornatus</i> (Reinhardt, 1843)	I/A	=	<i>Chilabothrus inornatus</i> (Reinhardt, 1843)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Epicrates monensis</i> Zenneck, 1898	I/A	=	<i>Chilabothrus monensis</i> (Zenneck, 1898)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Epicrates striatus</i> (Fischer, 1856)	II/B	>	<i>Chilabothrus striatus</i> (Fischer, 1856)	Reynolds <i>et al.</i> (2013)	Generic change

¹³ Species splits (indicated by the symbol ">") refer to cases where a currently recognised taxon has been split into various taxa in the associated reference

Taxon	CITES Appendix/ EU Annex	Proposed change	References	Notes
		<i>Chilabothrus strigilatus</i> (Cope, 1862)	Reynolds <i>et al.</i> (2013)	Species split
<i>Epicrates subflavus</i> Stejneger, 1901	I/A	= <i>Chilabothrus subflavus</i> (Stejneger, 1901)	Reynolds <i>et al.</i> (2013)	Generic change
<i>Eryx jayakari</i> Boulenger, 1888	II/B	= <i>Pseudogongylophis jayakari</i> (Boulenger, 1888)	Hoser (2013a)	Generic change
<i>Gongylophis colubrinus</i> (Linnaeus, 1758)	II/B	= <i>Eryx colubrinus</i> (Linnaeus, 1758)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Gongylophis conicus</i> (Schneider, 1801)	II/B	= <i>Eryx conicus</i> (Schneider, 1801)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Gongylophis muelleri</i> Boulenger, 1892	II/B	= <i>Eryx muelleri</i> (Boulenger, 1892)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Sanzinia madagascariensis</i> (Duméril & Bibron, 1844)	I/A	> <i>Sanzinia voluntary</i> Vences & Glaw, 2004	Reynolds <i>et al.</i> (2014)	Species split
COLUBRIDAE				
<i>Xenochrophis piscator</i> (Schneider, 1799)	III/C (IN)	> <i>Xenochrophis schnurrenbergeri</i> Kramer, 1977	Purkayastha <i>et al.</i> (2010), Vogel & David (2006, 2012)	Species split
		<i>Xenochrophis tytleri</i> (Blyth, 1863)	Vogel & David (2006, 2012)	Species split
ELAPIDAE				
<i>Micrurus diastema</i> (Duméril, Bibron & Duméril, 1854)	III/C (HN)	= <i>Hoserelapidea diastema</i> (Duméril, Bibron & Duméril, 1854)	Hoser (2012b)	Generic change
<i>Micrurus nigrocinctus</i> (Girard, 1854)	III/C (HN)	> <i>Hoserelapidea nigrocinctus</i> (Girard, 1855)	Hoser (2012b)	Generic change
		<i>Micrurus ruatanus</i> (Günther, 1895)	McCranie 2011, Townsend <i>et al.</i> (2013), Uetz (2013)	Species split
PYTHONIDAE				
<i>Apodora papuana</i> (Peters & Doria, 1878)	II/B	= <i>Liasis papuanus</i> Peters & Doria, 1878	Reynolds <i>et al.</i> (2014)	Generic change
<i>Antaresia stimsoni</i> (L. A. Smith, 1985)	II/B	= <i>Antaresia saxacola</i> Wells & Wellington, 1985	Hoser (2012a)	Name considered to be valid and to have date priority over <i>Antaresia stimsoni</i>
N/A	II/B	<i>Australiasis funki</i> Hoser, 2012	Hoser (2012a)	New species
<i>Leiopython albertisii</i> (Peters & Doria, 1878)	II/B	= <i>Bothrochilus albertisii</i> (Peters & Doria, 1878)	Reynolds <i>et al.</i> (2014)	Generic change

Taxon	CITES Appendix/ EU Annex	Proposed change	References	Notes
<i>Leiopython bennettorum</i> Hoser, 2000	II/B	= <i>Bothrochilus bennettorum</i> (Hoser, 2000)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Leiopython biakensis</i> Schleip, 2008	II/B	= <i>Bothrochilus biakensis</i> (Schleip, 2008)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Leiopython fredparkeri</i> Schleip, 2008	II/B	= <i>Bothrochilus fredparkeri</i> (Schleip, 2008)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Leiopython hoserae</i> Hoser, 2000	II/B	= <i>Bothrochilus hoserae</i> (Hoser, 2000)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Leiopython huonensis</i> Schleip, 2008	II/B	= <i>Bothrochilus huonensis</i> (Schleip, 2008)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia amethystina</i> (Schneider, 1801)	II/B	> <i>Australiasis amethystina</i> (Schneider, 1801)	Hoser (2012a)	Generic change
		<i>Australiasis clarki</i> (Barbour, 1914)	Hoser (2012a)	Species split
		<i>Australiasis duceboracensis</i> (Günther, 1879)	Hoser (2012a)	Species split
<i>Morelia amethystina</i> (Schneider, 1801)	II/B	= <i>Simalia amethystina</i> (Schneider, 1801)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia boeleni</i> (Brongersma, 1953)	II/B	= <i>Simalia boeleni</i> (Brongersma, 1953)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia carinata</i> (L. A. Smith, 1981)	II/B	= <i>Jackypython carinata</i> Hoser, 2009	Hoser (2009), Hoser (2012a)	Generic change
<i>Morelia clastolepis</i> Harvey, Barker, Ammerman & Chippindale, 2000	II/B	= <i>Australiasis clastolepis</i> (Harvey, Barker, Ammerman & Chippindale, 2000)	Hoser (2012a)	Generic change
<i>Morelia clastolepis</i> Harvey, Barker, Ammerman & Chippindale, 2000	II/B	= <i>Simalia clastolepis</i> (Harvey, Barker, Ammerman & Chippindale, 2000)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia kinghorni</i> (Stull, 1933)	II/B	= <i>Simalia kinghorni</i> (Stull, 1933)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia nauta</i> Harvey, Barker, Ammerman & Chippindale, 2000	II/B	= <i>Australiasis nauta</i> (Harvey, Barker, Ammerman & Chippindale, 2000)	Hoser (2012a)	Generic change
<i>Morelia nauta</i> Harvey, Barker, Ammerman & Chippindale, 2000	II/B	= <i>Simalia nauta</i> (Harvey, Barker, Ammerman & Chippindale, 2000)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia oenpelliensis</i> (Gow, 1977)	II/B	= <i>Simalia oenpelliensis</i> (Gow, 1977)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia tracyae</i> Harvey, Barker, Ammerman & Chippindale, 2000	II/B	= <i>Australiasis tracyae</i> (Harvey, Barker, Ammerman & Chippindale, 2000)	Hoser (2012a)	Generic change
<i>Morelia tracyae</i> Harvey, Barker, Ammerman & Chippindale, 2000	II/B	= <i>Simalia tracyae</i> (Harvey, Barker, Ammerman & Chippindale, 2000)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Morelia viridis</i> (Schlegel, 1872)	II/B	> <i>Chondropython viridis</i> (Schlegel, 1872)	Hoser (2012a)	Generic change

Taxon	CITES Appendix/ EU Annex		Proposed change	References	Notes
			<i>Chondropython azureus</i> Meyer, 1874	Hoser (2012a)	Species split
N/A	II/B		<i>Morelia wellsi</i> Hoser, 2012	Hoser (2012a)	New species
<i>Python reticulatus</i> (Schneider, 1801)	II/B	=	<i>Malayopython reticulatus</i> (Schneider, 1801)	Reynolds <i>et al.</i> (2014)	Generic change
<i>Python timoriensis</i> (Peters, 1876)	II/B	=	<i>Malayaopython timoriensis</i> (Peters, 1876)	Reynolds <i>et al.</i> (2014)	Generic change
TROPIDOPHIIDAE					
N/A	II/B		<i>Tropidophis grapiuna</i> Curcio, Sales Nunes, Suzart Argolo, Skuk & Rodrigues, 2012	Curcio <i>et al.</i> (2012)	New species
N/A	II/B		<i>Tropidophis preciosus</i> Curcio, Sales Nunes, Suzart Argolo, Skuk & Rodrigues, 2012	Curcio <i>et al.</i> (2012)	New species
TESTUDINES					
EMYDIDAE					
<i>Chrysemys picta</i> (Schneider, 1783)	-/B	>	<i>Chrysemys dorsalis</i> Agassiz, 1857	Turtle Taxonomy Working Group (2012)	Species split
GEOEMYDIDAE					
<i>Cuora galbinifrons</i> Bourret, 1939	II/B	>	<i>Cuora bourreti</i> Obst & Reimann, 1994	Turtle Taxonomy Working Group (2012)	Species split
			<i>Cuora picturata</i> Lehr, Fritz & Obst, 1998	Turtle Taxonomy Working Group (2012)	Species split
N/A	II/B		<i>Cyclemys enigmatica</i> Fritz, Guicking, Auer, Sommer, Wink & Hundsdörfer, 2009	Fritz <i>et al.</i> (2008), Turtle Taxonomy Working Group (2012)	New species
N/A	II/B		<i>Cyclemys fusca</i> Fritz, Guicking, Auer, Sommer, Wink & Hundsdörfer, 2009	Fritz <i>et al.</i> (2008), Turtle Taxonomy Working Group (2012)	New species
N/A	II/B		<i>Cyclemys gemeli</i> Fritz, Guicking, Auer, Sommer, Wink & Hundsdörfer, 2009	Fritz <i>et al.</i> (2008), Turtle Taxonomy Working Group (2012)	New species
<i>Cyclemys oldhamii</i> Gray, 1863	II/B	<	<i>Cyclemys shanensis</i> Annandale, 1918	Turtle Taxonomy Working Group (2012)	Species lump

Taxon	CITES Appendix/ EU Annex		Proposed change	References	Notes
<i>Mauremys reevesii</i> (Gray, 1831)	III/C (CN)	<	<i>Mauremys megaloccephala</i> (Fang, 1934)	Turtle Taxonomy Working Group (2012)	Species lump
TESTUDINIDAE					
<i>Chelonoidis nigra</i> (Quoy & Gaimard, 1824)	I/A	>	<i>Chelonoidis abingdonii</i> (Günther, 1877)	Turtle Taxonomy Working Group (2012)	Species split. Extinct since 2012
			<i>Chelonoidis becki</i> (Rothschild, 1901)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Chelonoidis chathamensis</i> (Van Denburgh, 1907)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Chelonoidis darwini</i> (Van Denburgh, 1907)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Chelonoidis duncanensis</i> (Garman in Pritchard, 1966)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Chelonoidis hoodensis</i> (Van Denburgh, 1907)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Chelonoidis phantastica</i> (Van Denburgh, 1907)	Turtle Taxonomy Working Group (2012)	Species split. Extinct since ca. 1960
			<i>Chelonoidis porteri</i> (Rothschild, 1903)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Chelonoidis vicina</i> (Günther, 1875)	Turtle Taxonomy Working Group (2012)	Species split
<i>Kinixys belliana</i> (Gray, 1831)	II/B	>	<i>Kinixys nogueyi</i> (Lataste, 1886)	Turtle Taxonomy Working Group (2012)	Species split
			<i>Kinixys zombensis</i> Hewitt, 1931	Turtle Taxonomy Working Group (2012)	Species split

References

- Curcio, F. F., Sales Nunes, P. M., Suzart Argolo, A. J., Skuk, G. & Rodrigues, M. T. 2012. Taxonomy of the South American dwarf boas of the genus *Tropidophis* Bibron, 1840, with the description of two new species from the Atlantic forest (Serpentes: Tropidophiidae). – *Herpetological Monographs* 26 (1): 80-121.
- Fritz, U., Guicking, D., Auer, M., Sommer, R. S., Wink, M. & Hundsdörfer, A. K. 2008. Diversity of the Southeast Asian leaf turtle genus *Cyclemys*: how many leaves on its tree of life? – *Zoologica Scripta* 37: 367-390.
- Greenbaum, E., Stanley, E. L., Kusamba, C., Moninga, W. M., Goldberg, S. R. & Cha 2012. A new species of *Cordylus* (Squamata: Cordylidae) from the Marungu Plateau of south-eastern Democratic Republic of the Congo. – *African Journal of Herpetology* 61 (1): 14-39.
- Harvey, M. B., Ugueto, G. N. & Gutberlet, R. L. Jr. 2012. Review of teiid morphology with a revised taxonomy and phylogeny of the Teiidae (Lepidosauria: Squamata). – *Zootaxa* 3459: 1–156.
- Henderson, R. W., Pauers, M. J. & Colston, T. J. 2013. On the congruence of morphology, trophic ecology, and phylogeny in Neotropical treeboas (Squamata: Boidae: *Corallus*). – *Biological Journal of the Linnean Society* 109: 466–475.
- Hoser, R. 2009. Creationism and contrived science: a review of recent python systematics papers and the resolution of issues of taxonomy and nomenclature. – *Australasian Journal of Herpetology* 2: 1-34 .
- Hoser, R. T. 2012a. An updated review of the pythons including resolution of issues of taxonomy and nomenclature. – *Australasian Journal of Herpetology* 10: 2-32.
- Hoser, R. T. 2012b. A reclassification of the rattlesnakes; species formerly exclusively referred to the genera *Crotalus* and *Sistrurus* and a division of the elapid genus *Micrurus*. – *Australasian Journal of Herpetology* 11: 2-24.
- Hoser, R. T. 2013a. Tidying up the taxonomy of the extant Booidea, including the erection and naming of two new families, the description of *Acrantophis sloppi* sp. nov., a new species of ground boa from Madagascar and *Candoia aspera iansimpsoni*, subsp. nov., a new subspecies of boa from Papua New Guinea. – *Australasian Journal of Herpetology* 16: 3-8.
- Hoser, R. T. 2013b. A revised taxonomy for the lizard families Gerrhosauridae and Cordylidae. – *Australasian Journal of Herpetology* 21: 3-32.
- McCranie, J. R. 2011. *The snakes of Honduras - systematics, distribution and conservation*. Thomson-Shore Inc., Michigan.
- Nielsen, S. V., Bauer, A. M., Jackman, T. R., Hitchmough, R. A. & Daugherty, C. H. 2011. New Zealand geckos (Diplodactylidae): Cryptic diversity in a post-Gondwanan lineage with trans-Tasman affinities. – *Molecular Phylogenetics and Evolution* 59 (1): 1-22.
- Purkayastha, J., Das, M., Sengupta, S. & Dutta, S. K. 2010. Notes on *Xenochrophis schnurrenbergeri* Kramer, 1977 (Serpentes: Colubridae) from Assam, India with some comments on its morphology and distribution. – *Herpetology Notes* 3: 175-180.
- Reynolds, R. G., Niemiller, M. L., Hedges, S. B., Dornburg, A., Puente-Rolón, A. R. & Revell, L. J. 2013. Molecular phylogeny and historical biogeography of West Indian boid snakes (*Chilabothrus*). – *Molecular Phylogenetics and Evolution* 68: 461-470.
- Reynolds, R. G., Niemiller, M. L. & Revell, L. J. 2014. Toward a tree-of life for the boas and pythons: multilocus species-level phylogeny with unprecedented taxon sampling. – *Molecular Phylogenetics and Evolution* 71: 201-213.
- Stanley, E. L., Bauer, A. M., Jackman, T. R., Branch, W. R. & P. le F. N. 2011. Between a rock and a hard polytomy: rapid radiation in the rupicolous girdled lizards (Squamata: Cordylidae). – *Molecular Phylogenetics and Evolution* 58(1): 53-70.
- Townsend, J. H., McCranie, J. R. & Stafford, P. 2013. *Micrurus ruatanus*. In: IUCN 2013. *IUCN Red List of Threatened Species*. Version 2013.2. <www.iucnredlist.org>. Downloaded on 31 December 2013.
- Turtle Taxonomy Working Group [van Dijk, P. P., Iverson, J. B., Shaffer, H. B., Bour, R. & Rhodin, A. G. J.] 2012. Turtles of the world, 2012 update: Annotated checklist of taxonomy, synonymy, distribution, and conservation status. In: Rhodin, A. G. J., Pritchard, P. C. H., van Dijk, P. P., Saumure, R. A., Buhmann, K. A., Iverson, J. B., and Mittermeier, R. A. (Eds.). *Conservation biology of freshwater turtles and tortoises: A compilation project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. - *Chelonian Research Monographs* 5: 243-328.
- Uetz, P. (ed.) 2013. *The reptile database*. <http://www.reptile-database.org>. Downloaded on 31 December 2013.
- Vogel, G. & David, P. 2006. On the taxonomy of the *Xenochrophis piscator* complex (Serpentes, Natricidae). – Pp. 231-246 in M. Vences, J. Kohler, T. Ziegler & W. Böhme, eds. *Herpetologia Bonnensis II*. Proceedings of the 13th Congress of the Societas Europaea Herpetologica.
- Vogel, G. & David, P. 2012. A revision of the species group of *Xenochrophis piscator* (Schneider, 1799) (Squamata: Natricidae). – *Zootaxa* 3473: 1-60.

Species information extracted from:
FROST, D.R. (2014):
"Amphibian Species of the World, on online Reference"
V. 6, downloaded February 18 2014

Copyright © 1998-2011, Darrel Frost and The American Museum of Natural History.
 All Rights Reserved.

ANURA

BUFONIDAE

Altiphrynoides Dubois, 1987

Altiphrynoides malcolmi (Grandison, 1978)

- *Nectophrynoides malcolmi* [Grandison, 1978, Monit. Zool. Ital., N.S., Suppl., 11](#): 124. Holotype: BMNH 1975.1961, by original designation. Type locality: "6-8 km SE Goba, road to Maslo, Balé Province, Ethiopia, 06° 59' N-40° 01' E, elevation 3200 m".
- *Altiphrynoides malcolmi* — [Dubois, 1987 "1986", Alytes, 5](#): 27.

Distribution: Bale Mountains at altitudes of 3200-4000 m, Bale Province, Ethiopia.

Comment: See [Largen, 2001, Tropical Zool., 14](#): 326, for comments on distribution. See photograph, map, description of geographic range and habitat, and conservation status in [Stuart, Hoffmann, Chanson, Cox, Berridge, Ramani, and Young, 2008, Threatened Amph. World](#): 156. See account (as *Altiphrynoides malcolmi*), photograph, and map for Ethiopia by [Largen and Spawls, 2010, Amph. Rept. Ethiopia Eritrea](#): 100-101.

DENDROBATIDAE

Hyloxalus Jiménez de la Espada, 1870

Hyloxalus azureiventris (Kneller and Henle, 1985)

- *Phyllobates azureiventris* Kneller and Henle, 1985, Salamandra, 21: 62. Holotype: ZFMK 41507, by original designation. Type locality: "km 26, Carretera Tarapoto--Yurimaguas, Departamento San Martín, Peru, ca. 700 m NN".
- *Dendrobates azureiventris* — Myers and Burrowes, 1987, Am. Mus. Novit., 2899: 1-17.
- *Epipedobates azureiventris* — Myers, 1987, Pap. Avulsos Zool., São Paulo, 36: 303.
- *Phyllobates (Pseudendrobates) azureiventris* — Bauer, 1988, Het Paludarium, Netherlands, November: 6.
- *Cryptophyllobates azureiventris* — Lötters, Jungfer, and Widmer, 2000, Jahresheft. Ges. Naturkd. Württemberg, 156: 236.
- *Ameerega azureiventris* — Frost, Grant, Faivovich, Bain, Haas, Haddad, de Sá, Channing, Wilkinson, Donnellan, Raxworthy, Campbell, Blotto, Moler, Drewes, Nussbaum, Lynch, Green, and Wheeler, 2006, Bull. Am. Mus. Nat. Hist., 297: 130. by implication.

- *Hyloxalus azureiventris* — Grant, Frost, Caldwell, Gagliardo, Haddad, Kok, Means, Noonan, Schargel, and Wheeler, 2006, *Bull. Am. Mus. Nat. Hist.*, 299: 168.

Distribution: Lower eastern versant of the Andes in the upper Amazon basin of the Department of San Martín, Peru.

Comment: Schulte, 1999, *Pfeilgifffrösche*: 245-253, provided an account (as *Epipedobates azureiventris*). See account by Lötters, Jungfer, Henkel, and Schmidt, 2007, *Poison Frogs*: 317-318. See photograph, map, description of geographic range and habitat, and conservation status (as *Cryptophyllobates azureiventris*) in Stuart, Hoffmann, Chanson, Cox, Berridge, Ramani, and Young, 2008, *Threatened Amph. World*: 227.

CAUDATA

HONOBIIDAE

Hynobius Tschudi, 1838

Hynobius amjiensis Gu, 1992

- *Hynobius amjiensis* [Gu, 1992 "1991", in Qian et al. \(eds.\), Animal Sci. Res.](#): 39. Holotype: HTC 90301, by original designation. Type locality: "Mount Longwang natural reserve, Anji County, Zhejiang Province; altitude 1300 m", China.
- *Hynobius* (*Hynobius*) *amjiensis* — [Dubois and Raffaëlli, 2012, Alytes](#), 28: 77-161.

Distribution: Known only from the Longwangshan Nature Reserve, Anji County, Zhejiang Province, China (30° 23.68'N, 119° 27.32'E), in a marshy meadow of about 7000 square meters, at the top of Longwangshan, ca. 1300 m elevation.

Comment: Not assigned to species group by [Thorn, 1968, Salamand. Eur. Asie Afr. Nord](#): 37. See accounts by [Ye, Fei, and Hu, 1993, Rare and Economic Amph. China](#): 29, and [Thorn and Raffaëlli, 2000, Salamand. Ancien Monde](#): 90. See [Fei, 1999, Atlas Amph. China](#), for figure, brief account, and distribution map. In the *Hynobius leechii* group of [Fei and Ye, 2005, in Fei et al. \(eds.\), Illust. Key Chinese Amph.](#): 30 (who only noted Chinese species). [Fu, Hayes, Liu, and Zeng, 2003, Acta Zool. Sinica](#), 49: 585-591, provided a discussion of molecular data in support of the species distinction of this species from *Hynobius yiwuensis*. [Fei, Hu, Ye, and Huang, 2006, Fauna Sinica, Amph. 1](#): 154-157, provided an account. See brief account by [Raffaëlli, 2007, Les Urodèles du Monde](#): 44. See illustration, map, description of geographic range and habitat, and conservation status in [Stuart, Hoffmann, Chanson, Cox, Berridge, Ramani, and Young, 2008, Threatened Amph. World](#): 549. [Fei, Ye, and Jiang, 2010, Colored Atlas of Chinese Amph.](#): 39, provided a brief account, photographs and illustration of specimens as well as a habitat shot. [Fei, Ye, and Jiang, 2012, Colored Atlas Chinese Amph. Distr.](#): 32-33, provided an account, photographs, and map for China.

**Additional species lacking from the
current nomenclature fish reference
either because species newly listed at and since CoP16
or overlooked by error prior to CoP16**

extracted from
ESCHMEYER, W.N. & FRICKE, R. (eds.):
Catalog of Fishes, electronic version (5 February 2014)

Downloaded February 9 2014

Copyright © W.N. Eschmeyer and California Academy of Sciences. All Rights reserved.

Footnotes added by the Nomenclature Specialist of the Animals Committee

ELASMOBRANCHII

CARCHARHINIFORMES

Carcharhinidae

Carcharhinus longimanus (POEY, 1861)

budkeri*, *Pterolamiops Fourmanoir [P.] 1961:76 [Mémoires de l'Institut Scientifique de Madagascar. Série F. Océanographie v. 4; ref. 9345] Mozambique Channel. Syntypes: (1) 12 miles west of Kalakazoro Island (1) off Majunga. •Synonym of *Carcharhinus longimanus* (Poey 1861) -- (Garrick 1982:150 [ref. 5454], Compagno 1984:484 [ref. 6846], Voigt & Weber 2011:77 [ref. 31424], Castro 2011:438 [ref. 31457]). **Current status:** Synonym of *Carcharhinus longimanus* (Poey 1861). Carcharhinidae. Habitat: marine.

insularum*, *Carcharias Snyder [J. O.] 1904:513, Pl. 1 (fig. 1) [Bulletin of the U. S. Fish Commission v. 22 [1902]; ref. 4149] Off Diamond Head, Oahu Island, Hawaiian Islands, Albatross station 3815, depth 228-312 fathoms. Holotype: USNM 50859. Paratypes: SU 12788-89 (1, 1). Type catalog: Böhlke 1953:9 [ref. 12291], Howe & Springer 1993:8 [ref. 21812]. •Synonym of *Carcharhinus longimanus* (Poey 1861) -- (Garrick 1982:150 [ref. 5454], Compagno 1984:484 [ref. 6846], Mundy 2005:90 [ref. 28379], Voigt & Weber 2011:77 [ref. 31424], Castro 2011:438 [ref. 31457]). **Current status:** Synonym of *Carcharhinus longimanus* (Poey 1861). Carcharhinidae. Habitat: marine.

longimanus*, *Squalus Poey [F.] 1861:338, Pl. 19 (figs. 9-10) [Memorias sobre la historia natural de la Isla de Cuba v. 2; ref. 3499] Cuba, western Atlantic. Holotype (unique): No types known. As *Prionodon longimanus* on plate. On Official List (Opinion 723). Authors point out that *Squalus (Carcharias) maou* Lesson 1830 probably is the same as this species. •Valid as *Carcharhinus longimanus* (Poey 1861) -- (Compagno 1973:24 [ref. 7163], Garrick 1982:150 [ref. 5454], Eschmeyer & Herald 1983:39 [ref. 9277], Compagno 1984:484 [ref. 6846], Branstetter in Whitehead et al. 1984:107 [ref. 13675], Dor 1984:6 [ref. 29757], Nakaya in Masuda et al. 1984:6 [ref. 6441], Bass et al. 1986:74 [ref. 5638], Robins & Ray 1986:26 [ref. 23100], Allen & Swainston 1988:22 [ref. 25477], Scott & Scott 1988:24 [ref. 25518], Paxton et al. 1989:78 [ref. 12442], McAllister 1990:32 [ref. 14674], Springer 1990:106 [ref. 19320], Randall et al. 1990:20 [ref. 15987], Boschung 1992:21 [ref. 23239], Cervigón 1992:173 [ref. 23827], Allen & Robertson 1994:22 [ref. 22193], Gomon et al. 1994:122 [ref. 22532], Last & Stevens 1994:247 [ref. 23873], Goren & Dor 1994:2 [ref. 25356], Compagno et al. in Fischer et al. 1995:677 [ref. 22829], Randall 1995:32 [ref. 22896], Santos et al. 1997:8 [ref. 23531], Allen 1997:42 [ref. 23977], De La Cruz Agüero et al. 1997:28 [ref. 24545], Grove & Lavenberg 1997:81 [ref. 24023], Arruda 1997:18 [ref. 24952], Randall et al. 1997:20 [ref. 25919], Compagno & Niem 1998:1341 [ref. 23787], McEachran & Fechhelm 1998:81 [ref. 23897], Chirichigno F. & Vélez D. 1998:53 [ref. 24555], Myers 1999:34 [ref. 23965], Fricke 1999:20 [ref. 24106], Francis et al. 1999:574 [ref. 24249], Morón et al. 1999:147 [ref. 24253], Smith-Vaniz et al. 1999:115 [ref. 25013], Lessa et al. 1999:353 [ref. 25267], Compagno 1999:483 [ref. 25589], Nakabo 2000:137 [ref. 25086], Compagno in Randall & Lim 2000:580 [ref. 25122], Laboute & Grandperrin 2000:92 [ref. 25191], Grace 2001:18 [ref. 25694], Hutchins 2001:14 [ref. 25847], Soto 2001:64, 79 [ref. 26637], Bilecenoglu et al. 2002:173 [ref. 26753], Nakabo 2002:137 [ref. 26001], Compagno 2003:484 [ref. 26984], Gadig & Gomes in Menezes et al. 2003:23 [ref. 27192], Manilo & Bogorodsky 2003:S92 [ref. 27377], Myers & Donaldson 2003:609 [ref. 27495], Randall et al. 2004:5 [ref. 27624], Espinosa Pérez et al. 2004:56 [ref. 27705], Bonfil & Abdallah 2004:33 [ref. 27735], Nelson et al. 2004:53 [ref. 27807], Mundy 2005:90 [ref. 28379], Compagno et al. 2005:39 [ref. 29145], Hoese et al. 2006:101 [ref. 29001], Randall 2007:30 [ref. 30952], White 2008:69 [ref. 30617], Fricke et al. 2009:9 [ref. 30213], George 2009:34 [ref. 30539], McCosker & Rosenblatt 2010:187 [ref. 30957], Fricke et al. 2011:345 [ref. 31242], Voigt & Weber 2011:77 [ref. 31424], Castro 2011:438 [ref. 31457], White 2012:3 [ref. 31843], Page et al. 2013:51 [ref. 32708], Wirtz et al. 2013:115 [ref. 32972], Larson et al. 2013:10 [ref. 32988], Ebert et al. 2013:332 [ref. 33045]). **Current status:** Valid as *Carcharhinus longimanus* (Poey 1861). Carcharhinidae. Distribution: Circumglobal in tropical and subtropical seas (including Red Sea), straying into temperate waters including North Sea. Habitat: marine.

magnipinnis*, *Pterolamiops Smith [J. L. B.] 1958:132, Pl. 1; Fig. 1C [Ichthyological Bulletin, Department of Ichthyology, Rhodes University No. 10; ref. 12027] Off Port Elizabeth, 7 miles from shore, 30°05'S, 25°18'E,

South Africa, western Indian Ocean, depth 50 fathoms. Holotype (unique): SAIAB [formerly RUSI] 126. •Synonym of *Carcharhinus longimanus* (Poey 1861) -- (Garrick 1982:150 [ref. 5454], Compagno 1984:484 [ref. 6846], Bass et al. 1986:74 [ref. 5638], Voigt & Weber 2011:77 [ref. 31424], Castro 2011:438 [ref. 31457]). **Current status:** Synonym of *Carcharhinus longimanus* (Poey 1861). Carcharhinidae. Habitat: marine.

maou, *Squalus (Carcharias)* Lesson [R. P.] 1831:91, Poissons Pl. 1 [Voyage autour du monde Zool. v. 2 (pt 1); ref. 2776] French Polynesia, 18°S, 144°W. Syntypes: (2) whereabouts unknown. See Garrick 1982:151 [ref. 5454] for comments on use of this name. We suggest continuing current usage of *Carcharhinus longimanus* (Poey 1861). •In the synonymy of the younger *Carcharhinus longimanus* (Poey 1861) -- (Garrick 1982:150 [ref. 5454], Branstetter in Whitehead et al. 1984:107 [ref. 13675], Compagno 1984:484 [ref. 6846], Bass et al. 1986:74 [ref. 5638], Randall et al. 1990:20 [ref. 15987], Allen & Robertson 1994:22 [ref. 22193], Compagno & Niem 1998:1341 [ref. 23787], Lessa et al. 1999:354 [ref. 25267], Compagno 2003:484 [ref. 26984], Espinosa Pérez et al. 2004:56 [ref. 27705], Bonfil & Abdallah 2004:33 [ref. 27735], Voigt & Weber 2011:77 [ref. 31424], Castro 2011:438 [ref. 31457]). **Current status:** Synonym of *Carcharhinus longimanus* (Poey 1861). Carcharhinidae. Habitat: marine.

Sphyrnidae

***Sphyrna gilberti* QUATTRO, DRIGGERS, GRADY, ULRICH & ROBERTS, 2013¹⁴**

gilberti, Sphyrna Quattro [J. M.], Driggers III [W. B.], Grady [J. M.], Ulrich [G. F.] & Roberts [M. A.] 2013:171, Figs. 5-6, 7A [Zootaxa 3702 (no. 2); ref. 32875] Bulls head, South Carolina, USA. Holotype: UF 183577. Paratypes: UF 183578 (2), 183579 (2). •Valid as *Sphyrna gilberti* Quattro, Driggers III, Grady, Ulrich & Roberts 2013. **Current status:** Valid as *Sphyrna gilberti* Quattro, Driggers III, Grady, Ulrich & Roberts 2013. Sphyrnidae. Distribution: Southeastern USA, possibly elsewhere. Habitat: marine.

***Sphyrna lewini* (GRIFFITH & SMITH, 1834)**

diplana, Sphyrna Springer [S.] 1941:46, Figs. 1-2 [Proceedings of the Florida Academy of Science v. 5; ref. 10183] Off Englewood, Florida, U.S.A., Gulf of Mexico, western Atlantic. Holotype: USNM 108451. Paratypes: USNM 108452 (1, head only), 110296-97 (2, jaws only). Type catalog: Howe & Springer 1993:6 [ref. 21812]. •Synonym of *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1967:38 [ref. 21135], Gilbert 1973:33 [ref. 7164], Dor 1984:11 [ref. 29757], Compagno 1984:545 [ref. 6846], Springer 1990:109 [ref. 19320], Gomon et al. 1994:139 [ref. 22532], Compagno et al. in Fischer et al. 1995:718 [ref. 22829], Compagno 1998:1364 [ref. 23785], Castro-Aguirre et al. 1999:54 [ref. 24550], Soto 2001:85 [ref. 26637], Compagno 2003:500 [ref. 26984], Espinosa Pérez et al. 2004:70 [ref. 27705], Bonfil & Abdallah 2004:36 [ref. 27735], Castro 2011:509 [ref. 31457], Quattro et al. 2013:176 [ref. 32875]). **Current status:** Synonym of *Sphyrna lewini* (Griffith & Smith 1834). Sphyrnidae. Habitat: brackish, marine.

erythraea, Zygaena Klunzinger [C. B.] (ex Ehrenberg) 1871:666 [Verhandlungen der K.-K. zoologisch-botanischen Gesellschaft in Wien v. 21; ref. 2622] Massawa, Eritrea, Red Sea. Type catalog: Paepke & Schmidt 1988:170 [ref. 21041] dated to 1899 with ZMB 7814 as holotype. Regarded as not available; appeared above as name in synonymy; also as name in synonymy in Hemprich & Ehrenberg 1899: 8, Pl. 6 (fig. 2) [ref. 4977]. Subsequent publication in an available way not researched. •Synonym of *Sphyrna lewini*

¹⁴ currently covered by CITES by the listing of *Sphyrna lewini*

(Griffith & Smith 1834) -- (Gilbert 1967:38 [ref. 21135], Gilbert 1973:33 [ref. 7164], Dor 1984:11 [ref. 29757], Compagno 1984:545 [ref. 6846]). **Current status:** Synonym of *Sphyrna lewini* (Griffith & Smith 1834). Sphyrnidae.

indica, Zygaena van Hasselt [J. C.] 1823:315 [Algemeene Konst- en Letter-bode I Deel (no. 20); ref. 4513] Vizagapatam, India; Java, Indonesia, Java Sea, eastern Indian Ocean. No types known. Based on figure in Russel, 1823, I., p. 8, pl. XII (see Alfred 1961:81, Pl. 3 [ref. 20553]). Also in van Hasselt 1824:90 [ref. 5104]. Should be suppressed and the use of the name *lewini* continued. •Available name -- (Kottelat 1987:369 [ref. 5962]). •Synonym of [later] *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1967:37 [ref. 21135] with question, Compagno 1984:545 [ref. 6846], Last et al. 2010:134 [ref. 32461]). **Current status:** Synonym of *Sphyrna lewini* (Griffith & Smith 1834). Sphyrnidae. Habitat: marine.

leeuwenii, Cestracion Day [F.] 1865:271 [The fishes of Malabar; ref. 1074] Malabar coast, India, Arabian Sea, western Indian Ocean. Unexplained new spelling for *Zygaena lewini* Griffith & Smith 1834. •Synonym of *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1967:38 [ref. 21135], Gilbert 1973:33 [ref. 7164], Compagno 1984:545 [ref. 6846]). **Current status:** Synonym of *Sphyrna lewini* (Griffith & Smith 1834). Sphyrnidae. Habitat: brackish, marine.

lewini, Zygaena Griffith [E.] & Smith [C. H.] 1834:640, Pl. 50 [The class Pisces, arranged by the Baron Cuvier; ref. 1908] [South coast of New Holland] southern Australia. No types known. Name available from Pl. 50. Based on a drawing by Mr. Lewin. Misspelled *leeuwenii* by Rochebrune 1883:44 [ref. 18639]. •Valid as *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1967:37 [ref. 21135], Gilbert 1973:33 [ref. 7164], Kyushin et al. 1982:23 [ref. 19754], Quéro in Whitehead et al. 1984:123 [ref. 13675], Compagno 1984:545 [ref. 6846], Dor 1984:11 [ref. 29757], Nakaya in Masuda et al. 1984:7 [ref. 6441], Nakaya in Okamura & Kitajima 1984:51, 300 [ref. 8057], Bass 1986:97 [ref. 5635], Robins & Ray 1986:30 [ref. 23100], Allen & Swainston 1988:24 [ref. 25477], Paxton et al. 1989:87 [ref. 12442], Springer 1990:109 [ref. 19320], Randall et al. 1990:23 [ref. 15987], Talwar & Jhingran 1991:28 [ref. 20764], Boschung 1992:24 [ref. 23239], Cervigón 1992:186 [ref. 23827], Allen & Robertson 1994:25 [ref. 22193], Gomon et al. 1994:139 [ref. 22532], Last & Stevens 1994:272 [ref. 23873], Goren & Dor 1994:4 [ref. 25356], Compagno et al. in Fischer et al. 1995:718 [ref. 22829], Randall 1995:38 [ref. 22896], Mohsin & Ambak 1996:67 [ref. 27969], Allen 1997:44 [ref. 23977], Chen et al. 1997:5 [ref. 26476], Cheng & Zhou 1997:35 [ref. 26385] with author as Griffith, Grove & Lavenberg 1997:96 [ref. 24023], Murdy et al. 1997:27 [ref. 23144], Larson & Williams 1997:343 [ref. 23967], Arruda 1997:19 [ref. 24952], Randall et al. 1997:23 [ref. 25919], De la Cruz-Agüero & Cota-Gómez 1998:356 [ref. 23520], Compagno 1998:1364 [ref. 23785], McEachran & Fechhelm 1998:93 [ref. 23897], Chirichigno F. & Vélez D. 1998:40 [ref. 24555], Myers 1999:36 [ref. 23965], Fricke 1999:24 [ref. 24106], Aguilera 1998:45 [ref. 24221], Cervigón & Alcalá 1999:97 [ref. 24490], Castro-Aguirre et al. 1999:53 [ref. 24550], Mishra & Srinivasan 1999:234 [ref. 24754], Smith-Vaniz et al. 1999:117 [ref. 25013], Afonso et al. 1999:68 [ref. 25466], Johnson 1999:720 [ref. 25471], Compagno 1999:484 [ref. 25589], Nakabo 2000:140 [ref. 25086], Compagno in Randall & Lim 2000:580 [ref. 25122], Laboute & Grandperrin 2000:97 [ref. 25191], Lea & Rosenblatt 2000:119 [ref. 25206], Thomson et al. 2000:282 [ref. 25640], Bijukumar & Sushama 2000:184 [ref. 25703], Randall & Earle 2000:5 [ref. 25806], Shane 2001:162 [ref. 25756], Hutchins 2001:14 [ref. 25847], Rocha & Rosa 2001:989 [ref. 25909], Soto 2001:65 [ref. 26637], Camargo & Isaac 2001:139 [ref. 27639] as Cuvier & Griffith & Smith 1834, Bilecenoglu et al. 2002:173 [ref. 26753], Nakabo 2002:140 [ref. 26001], Youn 2002:56, 475 [ref. 26218], Allen & Adrim 2003:22 [ref. 26830], Collette et al. 2003:98 [ref. 26784], Compagno 2003:500 [ref. 26984], Gadig & Gomes in Menezes et al. 2003:24 [ref. 27192], Manilo & Bogorodsky 2003:S92 [ref. 27377], Myers & Donaldson 2003:609 [ref. 27495], Smith et al. 2003:6 [ref. 27621], Randall et al. 2004:5 [ref. 27624], Espinosa Pérez et al. 2004:70 [ref. 27705], Bonfil & Abdallah 2004:36 [ref. 27735], Nelson et al. 2004:54 [ref. 27807], Heemstra & Heemstra 2004:68 [ref. 28072], Randall 2005:16 [ref. 28239], Mundy 2005:92 [ref. 28379], Randall et al. 2005:131 [ref. 28745], Compagno et al. 2005:46 [ref. 29145], Hoese et al. 2006:110 [ref. 29001], Wirtz et al. 2007:24 [ref. 30263], Randall 2007:37 [ref. 30952], Khalaf & Zajonz 2007:423 [ref. 31739], White 2008:73 [ref. 30617], McCosker & Rosenblatt 2010:187 [ref. 30957], Motomura et al. 2010:69 [ref. 31256], Last et al. 2010:136 [ref. 32461], Fricke et al. 2011:346 [ref. 31242], Castro 2011:509 [ref. 31457], Moore et al. 2012:11 [ref. 31771], Weigmann 2012:8 [ref. 31968], Allen & Erdmann 2012:59 [ref. 31980], Yamashita et al. 2012:125 [ref. 32408], Fricke et al. 2013:250 [ref. 32706], Page et al. 2013:52 [ref. 32708], Quattro et al. 2013:159 [ref. 32875], Wirtz et al. 2013:116 [ref. 32972] needs confirmation, Larson et al. 2013:13 [ref. 32988], Ebert et al. 2013:333 [ref. 33045]). **Current status:** Valid as *Sphyrna lewini* (Griffith & Smith 1834). Sphyrnidae. Distribution: Circumglobal in tropical seas (including western Mediterranean Sea, Red Sea). Habitat: brackish, marine.

oceanica, Cestracion Garman [S.] 1913:158 [Memoirs of the Museum of Comparative Zoology v. 36; ref. 1545] Society Islands, French Polynesia, South Pacific. Syntypes: MCZ 460-S (3), USNM 153587 [ex MCZ 460] (1). Type catalog: Howe & Springer 1993:10 [ref. 21812], Hartel & Dingerkus 1997:xl [ref. 23119]. •Synonym of *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1967:38 [ref. 21135], Gilbert 1973:33 [ref. 7164], Dor 1984:11 [ref. 29757], Compagno 1984:545 [ref. 6846], Castro-Aguirre et al. 1999:54 [ref. 24550]). **Current status:** Synonym of *Sphyrna lewini* (Griffith & Smith 1834). Sphyrnidae. Habitat: brackish, marine.

***Sphyrna mokarran* (RÜPPELL, 1837)**

chieraghini, Sphyrna Nardo [G. D.] 1847:col. 112 [Sinonimia moderna delle specie registrate nell' opera intitolata: ...; ref. 17994] Not available, name only based on *Squalus tiburo* of Chieraghini (manuscript). •In the synonymy of *Sphyrna mokarran* (Rüppell 1837) -- (Gilbert 1967:26 [ref. 21135]). **Current status:** Synonym of *Sphyrna mokarran* (Rüppell 1837). Sphyrnidae.

dissimilis, Zygaena Murray [J. A.] 1887:103, Pl. [Journal of the Bombay Natural History Society v. 2 (no. 2); ref. 17806] Karachi, Pakistan. Published in 3 places, earliest not determined. As above, in Ann. Mag. Nat. Hist. [see ref. 17806], and in Murray 1887 [ref. 17805]. •Synonym of *Sphyrna mokarran* (Rüppell 1837) -- (Gilbert 1967:26 [ref. 21135], Compagno 1984:548 [ref. 6846]). **Current status:** Synonym of *Sphyrna mokarran* (Rüppell 1837). Sphyrnidae. Habitat: brackish, marine.

ligo, Sphyrna Fraser-Brunner [A.] 1950:214, Fig. 1 [Records of the Australian Museum v. 22 (no. 3); ref. 12952] Clarence River, New South Wales, Australia. Holotype: BMNH 1890.9.23.231 [ex Imperial Inst.] (embryo). •Synonym of *Sphyrna mokarran* (Rüppell 1837) -- (Gilbert 1967:26 [ref. 21135], Compagno 1984:548 [ref. 6846], Paxton et al. 1989:87 [ref. 12442], Hoese et al. 2006:111 [ref. 29001]). **Current status:** Synonym of *Sphyrna mokarran* (Rüppell 1837). Sphyrnidae. Habitat: brackish, marine.

mokarran, Zygaena Rüppell [W. P. E. S.] 1837:66, Pl. 17 (fig. 3) [Neue Wirbelthiere zu der Fauna von Abyssinien gehörig. Fische des Rothen Meeres; ref. 3844] Massawa, Eritrea, Red Sea. Lectotype: SMF 3590 (stuffed). Type catalog and lectotype designation: Klauswitz 1960:293 [ref. 21330]. •Valid as *Sphyrna mokarran* (Rüppell 1837) -- (Gilbert 1967:26 [ref. 21135], Quérou in Whitehead et al. 1984:124 [ref. 13675], Compagno 1984:548 [ref. 6846], Dor 1984:11 [ref. 29757], Nakaya in Masuda et al. 1984:7 [ref. 6441], Bass 1986:97 [ref. 5635], Robins & Ray 1986:30 [ref. 23100], Allen & Swainston 1988:24 [ref. 25477], Paxton et al. 1989:87 [ref. 12442], Springer 1990:109 [ref. 19320], Randall et al. 1990:23 [ref. 15987], Boschung 1992:24 [ref. 23239], Cervigón 1992:187 [ref. 23827], Baranes & Golani 1993:301 [ref. 22372], Allen & Robertson 1994:25 [ref. 22193], Last & Stevens 1994:274 [ref. 23873], Goren & Dor 1994:4 [ref. 25356], Compagno et al. in Fischer et al. 1995:720 [ref. 22829], Randall 1995:38 [ref. 22896], Mohsin & Ambak 1996:68 [ref. 27969], Allen 1997:44 [ref. 23977], Grove & Lavenberg 1997:99 [ref. 24023] [but see McCosker 1998:809 [ref. 24025]], Compagno 1998:1365 [ref. 23785], McEachran & Feckhelm 1998:94 [ref. 23897], Aguilera 1998:45 [ref. 24221], Chirichigno F. & Vélez D. 1998:40 [ref. 24555], Myers 1999:36 [ref. 23965], Fricke 1999:24 [ref. 24106], Cervigón & Alcalá 1999:100 [ref. 24490], Castro-Aguirre et al. 1999:53 [ref. 24550], Smith-Vaniz et al. 1999:118 [ref. 25013], Johnson 1999:720 [ref. 25471], Compagno 1999:484 [ref. 25589], Nakabo 2000:140 [ref. 25086], Compagno in Randall & Lim 2000:580 [ref. 25122], Laboute & Grandperrin 2000:97 [ref. 25191], Schmitter-Soto et al. 2000:146 [ref. 27754], Hutchins 2001:14 [ref. 25847], Soto 2001:65, 85 [ref. 26637], Nakabo 2002:140 [ref. 26001], Allen & Adrim 2003:22 [ref. 26830], Compagno 2003:502 [ref. 26984], Gadig & Gomes in Menezes et al. 2003:24 [ref. 27192], Manilo & Bogorodsky 2003:S92 [ref. 27377], Espinosa Pérez et al. 2004:71 [ref. 27705], Bonfil & Abdallah 2004:36 [ref. 27735], Nelson et al. 2004:54 [ref. 27807], Heemstra & Heemstra 2004:68 [ref. 28072], Randall 2005:16 [ref. 28239], Mundy 2005:93 [ref. 28379], Randall et al. 2005:131 [ref. 28745], Compagno et al. 2005:46 [ref. 29145], Hoese et al. 2006:111 [ref. 29001], Randall 2007:38 [ref. 30952], Fricke et al. 2009:10 [ref. 30213], McCosker & Rosenblatt 2010:187 [ref. 30957], Last et al. 2010:138 [ref. 32461], Fricke et al. 2011:346 [ref. 31242], Castro 2011:516 [ref. 31457], Moore et al. 2012:11 [ref. 31771], Allen & Erdmann 2012:60 [ref. 31980], Fricke et al. 2013:250 [ref. 32706], Page et al. 2013:52 [ref. 32708], Wirtz et al. 2013:116 [ref. 32972], Larson et al. 2013:13 [ref. 32988], Ebert et al. 2013:340 [ref. 33045]). **Current status:** Valid as

Sphyrna mokarran (Rüppell 1837). Sphyrnidae. Distribution: Circumglobal in tropical through warm temperate seas (including Red Sea). Habitat: brackish, marine.

***Sphyrna zygaena* (LINNAEUS, 1758)**

carolinensis, Squalus (Cestrorhinus) Blainville [H. de] 1816:121 [Bulletin de la Société Philomathique de Paris v. 8; ref. 306] Not available, name only. •In the synonymy of *Sphyrna zygaena* (Linnaeus 1758) or *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1973:33 [ref. 7164]). •In the synonymy of *Sphyrna zygaena* (Linnaeus 1758) -- (Compagno 1984:553 [ref. 6846]). **Current status:** Synonym of *Sphyrna zygaena* (Linnaeus 1758). Sphyrnidae.

malleus, Zygaena Valenciennes [A.] 1822:223, Pl. 1 (fig. 1) [Mémoires du Muséum National d'Histoire Naturelle, Paris v. 9; ref. 17862] France; Brazil, southwestern Atlantic; Mediterranean Sea. No types known. •Synonym of *Sphyrna zygaena* (Linnaeus 1758) -- (Gilbert 1967:31 [ref. 21135], Gilbert 1973:32 [ref. 7164] as Cuvier 1816, Springer 1990:110 [ref. 19320], Soto 2001:66 [ref. 26637], Bilecenoglu et al. 2002:15 [ref. 26753] with authorship as Cuvier 1817, Mundy 2005:93 [ref. 28379]). **Current status:** Synonym of *Sphyrna zygaena* (Linnaeus 1758). Sphyrnidae. Habitat: brackish, marine.

pictus, Squalis (Cestrorhinus) Blainville [H. de] 1816:121 [Bulletin de la Société Philomathique de Paris v. 8; ref. 306] Not available, name only. •In the synonymy of *Sphyrna zygaena* (Linnaeus 1758) or *Sphyrna lewini* (Griffith & Smith 1834) -- (Gilbert 1973:33 [ref. 7164]). •In the synonymy of *Sphyrna zygaena* (Linnaeus 1758) -- (Compagno 1984:553 [ref. 6846]). **Current status:** Synonym of *Sphyrna zygaena* (Linnaeus 1758). Sphyrnidae.

subarcuata, Zygaena Storer [D. H.] 1848:71 [Proceedings of the Boston Society of Natural History v. 3 (1848-1851); ref. 18844] Harbor at Provincetown, Massachusetts, U.S.A., Cape Cod Bay, western North Atlantic. Holotype (unique): MCZ. Non-types: ?MCZ 1416-S (1), 89507-08 (1, 1). •Synonym of *Sphyrna zygaena* (Linnaeus 1758) -- (Gilbert 1967:31 [ref. 21135], Gilbert 1973:32 [ref. 7164], Compagno 1984:553 [ref. 6846], Castro 2011:529 [ref. 31457]). **Current status:** Synonym of *Sphyrna zygaena* (Linnaeus 1758). Sphyrnidae. Habitat: brackish, marine.

vulgaris, Zygaena Cloquet [H.] 1830:621 (v. 60) [Dictionnaire des sciences Naturelles; ref. 852] Mediterranean Sea, Arabian Sea, and Red Sea. whereabouts unknown. Based on *Squalus zygaena* Linnaeus 1758, probably to avoid strickland tautonymy. •Synonym of *Sphyrna zygaena* (Linnaeus 1758) -- (Gilbert 1973:32 [ref. 7164], Compagno 1984:553 [ref. 6846]). **Current status:** Synonym of *Sphyrna zygaena* (Linnaeus 1758). Sphyrnidae. Habitat: brackish, marine.

zygaena, Squalus Linnaeus [C.] 1758:234 [Systema Naturae, Ed. X v. 1; ref. 2787] Mediterranean Sea and Atlantic [original: "Europa, America"]; localities include Spain; Marseille, France; Rome, Italy; Lesbos Island, Greece; Syria, Mediterranean Sea. Syntypes: NRM 88 (1). Type catalog: Fernholm & Wheeler 1983:208 [ref. 20707]. Originally based on multiple species and at least 24 pre-Linnaean sources including Artedi 1738:68 [ref. 30578] and Artedi 1738:96 [ref. 30349] as *Squalus capite latissimo transverso mallei instar*, Rondelet 1554:389 [ref. 30354] as *Zygæna*, Salviani 1558:128 [ref. 30229] as *Libella*, Willughby 1686:55 [ref. 30164] and Ray 1713:20 [ref. 30231] as *Zygæna*. Spelled *zygena* by Bonnaterre 1788:9 [ref. 4940], and *zigoena* by Cabrera, Pérez & Haenseler 1817:11 [ref. 17319], see Graells 1887:185 [ref. 30438]. Spelled *zygoena* by Chevey 1932:6 [ref. 23060] as *Cestracion*. Name spelled *Squalus zigaena* by Sonnini 1803:74 [ref. 30464] and Berthelot 1890:115 [ref. 30346]. •Valid as *Sphyrna zygaena* (Linnaeus 1758) -- (Gilbert 1967:31 [ref. 21135], Gilbert 1973:32 [ref. 7164], Eschmeyer & Herald 1983:43 [ref. 9277], Quérou in Whitehead et al. 1984:125 [ref. 13675], Compagno 1984:553 [ref. 6846], Nakaya in Masuda et al. 1984:7 [ref. 6441], Bass 1986:97 [ref. 5635], Robins & Ray 1986:30 [ref. 23100], Allen & Swainston 1988:24 [ref. 25477], Scott & Scott 1988:29 [ref. 25518], Paxton et al. 1989:87 [ref. 12442], Pequeño 1989:13 [ref. 14125], Paulin et al. 1989:26 [ref. 24556], McAllister 1990:32 [ref. 14674], Springer 1990:110 [ref. 19320], Cervigón 1992:188 [ref. 23827], Francis 1993:157 [ref. 25479], Francis & Randall 1993:128 [ref. 20996], Kuitert 1993:11 [ref. 23929], Gomon et al. 1994:140 [ref. 22532], Last & Stevens 1994:275 [ref. 23873], Compagno et al. in

Fischer et al. 1995:722 [ref. 22829], Mohsin & Ambak 1996:69 [ref. 27969], Murdy et al. 1997:28 [ref. 23144], Santos et al. 1997:9 [ref. 23531], Allen 1997:44 [ref. 23977], Cheng & Zhou 1997:35 [ref. 26385], Grove & Lavenberg 1997:100 [ref. 24023], Arruda 1997:19 [ref. 24952], Compagno 1998:1366 [ref. 23785], Chirichigno F. & Vélez D. 1998:39 [ref. 24555], Sokolovskaya et al. 1998:7 [ref. 24670], Fricke 1999:25 [ref. 24106], Morón et al. 1999:148 [ref. 24253], Smith-Vaniz et al. 1999:118 [ref. 25013], Capapé et al. 1999:50 [ref. 25276], Compagno 1999:484 [ref. 25589], Nakabo 2000:140 [ref. 25086], Compagno in Randall & Lim 2000:580 [ref. 25122], Lea & Rosenblatt 2000:119 [ref. 25206], Hutchins 2001:14 [ref. 25847], Wang et al. 2001:34 [ref. 26566], Soto 2001:65, 85 [ref. 26637], Bilecenoglu et al. 2002:15 [ref. 26753], Branstetter in Collette & Klein-MacPhee 2002:46 [ref. 26158], Nakabo 2002:140 [ref. 26001], Youn 2002:56, 475 [ref. 26218], López et al. 2002:62 [ref. 26808], Compagno 2003:505 [ref. 26984], Gadig & Gomes in Menezes et al. 2003:24 [ref. 27192], Espinosa Pérez et al. 2004:73 [ref. 27705], Nelson et al. 2004:54 [ref. 27807], Heemstra & Heemstra 2004:68 [ref. 28072], Mundy 2005:93 [ref. 28379], Compagno et al. 2005:46 [ref. 29145], Hoese et al. 2006:111 [ref. 29001], Fricke et al. 2007:15 [ref. 29533], Vasil'eva 2007:16 [ref. 30517], Randall 2007:39 [ref. 30952], White 2008:74 [ref. 30617], George 2009:55 [ref. 30539], McCosker & Rosenblatt 2010:187 [ref. 30957], Castro 2011:529 [ref. 31457], Yamashita et al. 2012:125 [ref. 32408], Page et al. 2013:52 [ref. 32708], Wirtz et al. 2013:116 [ref. 32972], Ebert et al. 2013:340 [ref. 33045]). **Current status:** Valid as *Sphyrna zygaena* (Linnaeus 1758). Sphyrnidae. Distribution: Cosmopolitan in warm temperate seas, occasionally in tropical seas (including Mediterranean Sea, Black Sea, Hawaiian Islands). Habitat: brackish, marine.

RAJIFORMES

Mobulidae

Manta alfredi (KREFFT, 1868)

alfredi, *Deratoptera* Krefft [J. L. G.] 1868:3, 9, Fig. [The Illustrated Sydney News v. 5 (11 July 1868); ref. 5074] Watson's Bay, at entrance to Sydney Harbour, New South Wales, Australia. Holotype (unique): AMS I.1731(stuffed and painted over). Original description reproduced in Whitley 1936:176 [ref. 6075] and with correction of type locality. Spelling of the original genus should have been *Ceratoptera*, *Deratoptera* regarded as a typesetting error. Authorship has been attributed to Krefft (by Whitley 1936:176 and unquestioned), the then curator of the Museum; no author is given with the article. •Valid as *Manta alfredi*, but may be *Manta birostris* (Walbaum 1792) -- (Paxton et al. 1989:51 [ref. 12442], Compagno 1999:498 [ref. 25589]). •Synonym of *Manta birostris* (Donndorff 1798 [Walbaum 1792]) -- (Last & Stevens 1994:460 [ref. 23873], Allen et al. 2006:207 [ref. 29002]). •See Grove & Lavenberg 1997:130 [ref. 24023]. •Valid as *Manta alfredi* (Krefft 1868) -- (Marshall et al. 2009:13 [ref. 30599], Kitchen-Wheeler 2010:351 [ref. 30962], Fricke et al. 2011:348 [ref. 31242], Wirtz et al. 2013:116 [ref. 32972], Larson et al. 2013:22 [ref. 32988], Ebert et al. 2013:368 [ref. 33045]). **Current status:** Valid as *Manta alfredi* (Krefft 1868). Myliobatidae: Mobulinae. Distribution: Circumglobal in tropical through subtropical seas (including Red Sea and Hawaiian Islands). Habitat: marine.

fowleri, *Manta* Whitley [G. P.] 1936:182 [Australian Zoologist v. 8 (pt 3); ref. 6075] Tabuaeran [= Fanning Island], Line Islands, central Pacific. Holotype (unique): whereabouts unknown. Based on illustrations and a description of *Manta birostris* by Fowler 1927:3, Pl. 1 (figs. D-G) [ref. 15785]. •Synonym of *Manta alfredi* (Krefft 1868) -- (Marshall et al. 2009:13 [ref. 30599]). **Current status:** Synonym of *Manta alfredi* (Krefft 1868). Myliobatidae: Mobulinae. Habitat: marine.

pakoka, *Manta* Whitley [G. P.] 1936:183 [Australian Zoologist v. 8 (pt 3); ref. 6075] Near Hat Island (Teuaa), Ua Huka, Marquesas Islands. No types known. Based on a *Manta* with the Polynesian vernacular name *Pakoka* by Pinchot, "To the South Seas" 1930:406, 408, 421, fig. •Synonym of *Manta alfredi* (Krefft 1868) -- (Marshall et al. 2009:13 [ref. 30599]). **Current status:** Synonym of *Manta alfredi* (Krefft 1868). Myliobatidae: Mobulinae. Habitat: marine.

***Manta birostris* (WALBAUM, 1792)**

americana*, *Manta Bancroft [E. N.] 1829:454 [Zoological Journal, London v. 4 (no. 16) (art. 55); ref. 5051] American Seas. Apparently an alternate name for *Cephalopterus manta* Bancroft 1829 proposed in the same paper (p. 453). •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

birostris*, *Raja Walbaum [J. J.] 1792:535 [Petri Artedi sueci genera piscium Part 3; ref. 4572] No locality stated. No types known. Some authors (e.g., Whitley 1936:180 [ref. 6075]) date to Donndorff 1798:876 because they regarded Walbaum's treatment as non-binominal, but the style of this section suggests that the second word "birostris" was not italicized through an oversight (see index) and binominal nomenclature was intended. Earlier authors, such as Jordan & Evermann 1896:92 [ref. 2443] credited the name to Walbaum. See remarks under *Cephalopterus giorna* Lesueur 1824. •Valid as *Manta birostris* (Walbaum 1792) [sometimes with author as Donndorff 1790 or 1798] -- (Eschmeyer & Herald 1983:57 [ref. 9277], Nakaya in Masuda et al. 1984:16 [ref. 6441], Robins & Ray 1986:44 [ref. 23100], Allen & Swainston 1988:28 [ref. 25477], Scott & Scott 1988:59 [ref. 25518], Winterbottom et al. 1989:5 [ref. 13251], McEachran & Séret 1990:73 [ref. 19318], Nishida 1990:92 [ref. 19783], Randall et al. 1990:31 [ref. 15987], Boschung 1992:28 [ref. 23239], Cervigón 1992:201 [ref. 23827], Allen & Robertson 1994:37 [ref. 22193], Last & Stevens 1994:459 [ref. 23873], Randall 1995:49 [ref. 22896], Acero P. & Franke 1995:17 [ref. 22546], McEachran & Notarbartolo-di-Sciara in Fischer et al. 1995:764 [ref. 22829], Castro-Aguirre & Espinosa Pérez 1996:62 [ref. 22793], Murdy et al. 1997:49 [ref. 23144], Santos et al. 1997:16 [ref. 23531], Allen 1997:48 [ref. 23977], Grove & Lavenberg 1997:129 [ref. 24023], Arruda 1997:24 [ref. 24952], Randall et al. 1997:31 [ref. 25919], McEachran & Fechhelm 1998:194 [ref. 23897], Chirichigno F. & Vélez D. 1998:73 [ref. 24555], Myers 1999:40 [ref. 23965], Fricke 1999:33 [ref. 24106], Aguilera 1998:46 [ref. 24221], Cervigón & Alcalá 1999:206 [ref. 24490], Compagno & Last 1999:1527 [ref. 24637], Smith-Vaniz et al. 1999:124 [ref. 25013], Johnson 1999:718 [ref. 25471], Compagno 1999:498 [ref. 25589], Menni & Stehmann 2000:94 [ref. 24909], Nakabo 2000:186 [ref. 25086], Compagno in Randall & Lim 2000:583 [ref. 25122], Thomson et al. 2000:284 [ref. 25640], Randall & Earle 2000:5 [ref. 25806], Allen 2000:95 [ref. 25868], Iwatsuki et al. 2000:96 [ref. 26368], Schmitter-Soto et al. 2000:147 [ref. 27754], Hutchins 2001:16 [ref. 25847], McEachran in Collette & Klein-MacPhee 2002:80 [ref. 26158], Mecklenburg et al. 2002:108 [ref. 25968], Nakabo 2002:186 [ref. 26001], Allen & Adrim 2003:22 [ref. 26830] with author and date as Donndorf 1798, Gadig & Gomes in Menezes et al. 2003:31 [ref. 27192] with author and date as Donndorff 1798, McEachran & Carvalho 2003:588 [ref. 26985], Milessi & Oddone 2003:126 [ref. 27316] with author and date as Donndorf 1798, Manilo & Bogorodsky 2003:S94 [ref. 27377], Duffy & Abbott 2003:715 [ref. 27479], Myers & Donaldson 2003:610 [ref. 27495], Lobel & Lobel 2004:67 [ref. 27576], Randall et al. 2004:5 [ref. 27624], Bonfil & Abdallah 2004:55 [ref. 27735], Nelson et al. 2004:57 [ref. 27807], Heemstra & Heemstra 2004:88 [ref. 28072], Randall 2005:23 [ref. 28239], Mundy 2005:107 [ref. 28379], Compagno et al. 2005:80 [ref. 29145], Allen et al. 2006:207 [ref. 29002], Randall 2007:47 [ref. 30952], Fricke et al. 2009:12 [ref. 30213], Kimura 2009:16 [ref. 30426], Marshall et al. 2009:4 [ref. 30599], McCosker & Rosenblatt 2010:188 [ref. 30957], Kitchen-Wheeler 2010:351 [ref. 30962], Allen & Erdmann 2012:67 [ref. 31980], Page et al. 2013:57 [ref. 32708], Wirtz et al. 2013:116 [ref. 32972], Ebert et al. 2013:369 [ref. 33045]). **Current status:** Valid as *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Distribution: Circumglobal in tropical through warm temperate water (including Red Sea). Habitat: marine.

ehrenbergii*, *Ceratoptera Müller [J.] & Henle [F. G. J.] 1841:187 [Systematische Beschreibung der Plagiostomen; ref. 3069] Red Sea. Syntypes: ZMB 4708 (1), 22621 [ex Anat.-zool. Mus. 8731] (1) Type catalog: Paepke & Schmidt 1988:180 [ref. 21041]. •Valid as *Manta ehrenbergii* (Müller & Henle 1841) -- (Dor 1984:21 [ref. 29757], Goren & Dor 1994:6 [ref. 25356]). •Synonym of *Manta birostris* (Walbaum 1792) -- (Marshall et al. 2009:4 [ref. 30599]). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Distribution: Red Sea endemic [if valid]. Habitat: marine.

elliotti*, *Diabolicthys Holmes [F. S.] 1856:45 (39?) [Proceedings of the Elliott Society of Natural History v. 1; ref. 2194] Charleston, South Carolina, U.S.A. No types known. •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

fimbriata, Raja Lacepède [B. G. E.] 1802:671, 677, Pl. 16 (fig. 3) [Histoire naturelle des poissons (Lacepède) v. 4; ref. 4929] North Atlantic. No types known. Type catalog: Séret & McEachran 1986:34 [ref. 9312]. Also appeared in Sonnini 1802:76, 299 [ref. 30461] with author as Lacepède. •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

giorna, Cephalopterus Lesueur [C. A.] 1824:115 [Journal of the Academy of Natural Sciences, Philadelphia v. 4 (pt 1); ref. 17520] Georgia, U.S.A. No types known. Not *Raja giorna* Lacepède 1803. Marshall et al. 2009:13 [ref. 30599] tentatively recognize a valid species distinct from (but occurring with) *birostris*; it occurs in the Atlantic. A neotype selection would be needed to fix the name of this species as *giorna* Lesueur 1824 if the species is shown to be distinct from *birostris*. •Synonym of *Manta birostris* (Walbaum 1792) -- (authors). •Treated as a synonym of *Manta* sp. cf. *birostris* (Walbaum 1792) -- (Marshall et al. 2009:22 [ref. 30599]). **Current status:** Uncertain as *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Distribution: Atlantic, including the Caribbean. Habitat: marine.

hamiltoni, Brachioptilon Hamilton [F.] & Newman [E.] in Newman 1849:2358 [The Zoologist: a Popular Miscellany of Natural History. v. 7 (for 1849); ref. 3169] Gulf of California, Mexico. No types known. Description based on account by Hamilton, Newman provided name; therefore we treat authorship as Hamilton & Newman. •May be a synonym of *Manta birostris* (Walbaum 1792) -- (Compagno 1999:498 [ref. 25589]). •See Chirichigno F. & Vélez D. 1998:73 [ref. 24555]. •Synonym of *Manta birostris* (Walbaum 1792) - (Mecklenburg et al. 2002:108 [ref 25968], Marshall et al. 2009:4 [ref. 30599]). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

johnii, Ceratoptera Müller [J.] & Henle [F. G. J.] 1841:186, [Pl. 59 (right)] [Systematische Beschreibung der Plagiostomen; ref. 3069] Jamaica. Holotype (unique): RUSM uncat. (whereabouts unknown). •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

manatia, Raja Bloch [M. E.] & Schneider [J. G.] (ex Lacepède) 1801:364 [M. E. Blochii, Systema Ichthyologiae; ref. 471] Tropical America. No types known. •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

manta, Cephalopterus Bancroft [E. N.] 1829:453 [Zoological Journal, London v. 4 (no. 16) (art. 55); ref. 5051] Kingston, Jamaica. No types known. •Synonym of *Manta birostris* (Walbaum 1792) -- (Allen et al. 2006:207 [ref. 29002]). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

marinus, Raja diabolus Bloch [M. E.] & Schneider [J. G.] 1801:368 [M. E. Blochii, Systema Ichthyologiae; ref. 471] India. No types known. Not the same as *Raja diabolus* Shaw 1904. •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

orissa, Ceratoptera Lloyd [R. E.] 1908:176, Fig. 1, Pl. 5 (figs. 1-3) [Records of the Indian Museum (Calcutta) v. 2 (pt 2); ref. 14327] Puri, Bay of Bengal, Orissa coast, India. Holotype: ZSI F72968/1. Type catalog: Menon & Yazdani 1968:97 [ref. 20743]. •Synonym of *Manta birostris* -- (pers. comm., W. White, 11 April 2011). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

pinchoti, Manta Whitley [G. P.] 1936:182 [Australian Zoologist v. 8 (pt 3); ref. 6075] Near Hat Island, Ua Huka, Marquesas Islands. Holotype: USNM 89721 (parts). Paratypes: USNM 89722 (parts), 143796 (pieces), 143797 (1). Based on descriptions of *Manta birostris* by Fowler 1932:2 [ref. 1412]. Myliobatidae: Mobulinae. Habitat: marine.

raya, Manta Baer [G. A.] 1899:112 [Bulletin du Muséum National d'Histoire Naturelle (Série 1) v. 5 (no. 3); ref. 15635] Zorritos, about 40 kilometers south of Tombez, 4 kilometers from Grau, Peru. No types known. •Probably a synonym of *Manta birostris* (Walbaum 1792). **Current status:** Uncertain as *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

stelligera, Cephaloptera Günther [A.] (ex Ehrenberg) 1870:498 [Catalogue of the fishes in the British Museum v. 8; ref. 1995] Red Sea. Syntypes: ZMB 4701 (1), 22621 [ex 8731] (1). Type catalog: Paepke & Schmidt 1988:179 [ref. 21041] with ZMB 47801 and 22621 as syntypes. In footnote as name on unpublished plate under *Ceratoptera ehrenergii*; later appeared in Hemprich & Ehrenberg 1899:7, Pl. 2 (figs. 1-9) and second pl. 10 [ref. 4977] but treated in synonymy by editor Hilgendorf so not available. •In the synonymy of *Manta ehrenbergii* (Müller & Henle 1841) -- (Dor 1984:21 [ref. 29757]). •In the synonymy of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae.

vampyrus, Cephalopterus Mitchill [S. L.] 1824:23, Pl. 2 (fig. 1) [Annals of the Lyceum of Natural History of New York v. 1; ref. 17777] Near entrance to Delaware Bay, U.S.A. No types known. •Synonym of *Manta birostris* (Walbaum 1792). **Current status:** Synonym of *Manta birostris* (Walbaum 1792). Myliobatidae: Mobulinae. Habitat: marine.

OSTEOGLOSSIFORMES

Osteoglossidae

***Scleropages inscriptus* ROBERTS, 2012**

(currently regarded as part of *Scleropages formosus*, see Notification 2012/43)

inscriptus, Scleropages Roberts [T. R.] 2012:115, Figs. 1-2 [aqua, International Journal of Ichthyology v. 18 (no. 2); ref. 31952] Supposedly from Tananthayi district, Tananthayi River basin, obtained dead from aquarium fish vendor at Meik. Holotype: THNHM-F-01521. Paratypes: THNHM-F-01522 (1). •Valid as *Scleropages inscriptus* Roberts 2012 -- (Kottelat 2013:31 [ref. 32989]). **Current status:** Valid as *Scleropages inscriptus* Roberts 2012. Osteoglossidae. Distribution: Myanmar. Habitat: freshwater.

SARCOPTERYGII

CERATODONTIFORMES

Ceratodontidae

***Neoceratodus forsteri* (KREFFT, 1870)**

blanchardi, Neoceratodus Castelnau [F. L.] 1876:133 [Journal de Zoologie v. 5; ref. 760] Fitzroy River, Queensland, Australia. Holotype (unique): MNHN A-0419. Type catalog: Bertin 1940:246 [ref. 293]. •Synonym of *Neoceratodus forsteri* (Krefft 1870) -- (Paxton et al. 1989:102 [ref. 12442], Allen & Cross 2006:213 [ref. 29003], Kemp 1997:720 [ref. 31059]). **Current status:** Synonym of *Neoceratodus forsteri* (Krefft 1870). Neoceratodontidae. Habitat: freshwater.

forsteri, Ceratodus Krefft [J. L. G.] 1870:65, col. 5, Fig. 1-3 [Sydney Morning Herald [newspaper] 18 Jan. 1870; ref. 18433] Wide-Bay District, Queensland, Australia. No types known. Description also appeared in Krefft 1870:221 [ref. 18434]. •Valid as *Neoceratodus forsteri* (Krefft 1870) -- (Paxton et al. 1989:102 [ref. 12442], Kemp 1995 [ref. 22139], Unmack 2001:1060 [ref. 25797], Allen et al. 2002:55 [ref. 25930], Allen & Cross 2006:212 [ref. 29003], Kemp 1997:720 [ref. 31059]). **Current status:** Valid as *Neoceratodus forsteri* (Krefft 1870). Neoceratodontidae. Distribution: Australia. Habitat: freshwater.

mirolepis, *Ceratodus* Günther [A.] 1871:222 [1] [Annals and Magazine of Natural History (Series 4) v. 7 (no. 39); ref. 19154] Mary River, Queensland, Australia. Possible type or Günther specimen: AMS A.14079 (1). Also appeared in Günther 1871:377 [ref. 19155] and in more detail in Günther 1872 [ref. 19153]. •Synonym of *Neoceratodus forsteri* (Krefft 1870) -- (Paxton et al. 1989:102 [ref. 12442], Allen & Cross 2006:212 [ref. 29003], Kemp 1997:720 [ref. 31059]). **Current status:** Synonym of *Neoceratodus forsteri* (Krefft 1870). Neoceratodontidae. Habitat: freshwater.

COELACANTHIFORMES

Latimeriidae

Latimeria chalumnae SMITH, 1939

anjouanae, *Malania* Smith [J. L. B.] 1953:100 [Nature (London) v. 171 (no. 4342); ref. 4088] Comoro Islands, western Indian Ocean. Holotype (unique): SAIAB [formerly RUSI] 600. Specimen parts: USNM 163126 (2 scales). Possibly appeared first in Time Magazine, v. 61 (no. 2) dated 12 Jan. 1953; Nature dated 17 Jan. 1953. •Synonym of *Latimeria chalumnae* Smith 1939. **Current status:** Synonym of *Latimeria chalumnae* Smith 1939. Latimeriidae. Habitat: marine.

chalumnae, *Latimeria* Smith [J. L. B.] 1939:455 [Nature (London) v. 143; ref. 4068] Western Indian Ocean: west of East London, Cape Colony, South Africa, depth 40 fathoms. Holotype: East London Mus. Specimen parts: USNM 112258 (scales). •Valid as *Latimeria chalumnae* Smith 1939 -- (Heemstra in Smith & Heemstra 1995:151 [ref. 21953], Bruton 1995:104 [ref. 22132], Heemstra et al. 1996:150 [ref. 25940], Springer 1999:453 [ref. 24816], Holder et al. 1999:12616 [ref. 25259], Heemstra & Heemstra 2004:93 [ref. 28072]). **Current status:** Valid as *Latimeria chalumnae* Smith 1939. Latimeriidae. Distribution: Comoros, South Africa, Kenya, Madagascar and Mozambique. Habitat: marine.

Latimeria menadoensis POUYAUD, WIRJOATMODJO, RACHMATIKA, TJAKRAWIDJAJA, HADIATY & HADIE, 1999

menadoensis, *Latimeria* Pouyaud [L.], Wirjoatmodjo [S.], Rachmatika [I.], Tjakrawidjaja [A.], Hadiaty [R.] & Hadie [W.] 1999:266, Fig. [Comptes rendus de l'Académie des Sciences. Série 3, Sciences de la vie = Life Sciences Paris (Elsevier) No. 322; ref. 23788] Menadotua Island, Sulawesi, Indonesia. Holotype (unique): LBN [Mus. Bogoriense Zool., Cibinog]. •Valid as *Latimeria menadoensis* Pouyaud, Wirjoatmodjo, Rachmatika, Tjakrawidjaja, Hadiaty & Hadie 1999 -- (Holder et al. 1999:12616 [ref. 25259], Jewett 2001:3969 [ref. 26320]). **Current status:** Valid as *Latimeria menadoensis* Pouyaud, Wirjoatmodjo, Rachmatika, Tjakrawidjaja, Hadiaty & Hadie 1999. Latimeriidae. Distribution: Indonesia. Habitat: marine.

Differences between the current CITES Appendices and species currently planned to be included in the non-passerine volume of the HBW and BirdLife International Illustrated Checklist of the Birds of the World

CITES-RELEVANT SPLITS

Family	Parent species	Relevant CITES listings
Struthionidae	Struthio camelus	I/NC
Rheidae	Rhea pennata	I/II
Cracidae	Pauxi pauxi	III
Anatidae	Sarkidiornis melanotos	II
Falconidae	Falco chicquera	II
Accipitridae	Pernis celebensis	II
Accipitridae	Circus spilonotus	II
Accipitridae	Circus cyaneus	II
Accipitridae	Accipiter tachiro	II
Accipitridae	Accipiter novaehollandiae	II
Accipitridae	Accipiter striatus	II
Accipitridae	Buteo nitidus	II
Accipitridae	Buteo oreophilus	II
Accipitridae	Nisaetus philippensis	II
Otididae	Chlamydotis undulata	I
Columbidae	Goura scheepmakeri	II
Psittacidae	Trichoglossus haematodus	II
Psittacidae	Trichoglossus flavoviridis	II
Psittacidae	Chamosyna papou	II
Psittacidae	Psittacula picta	II
Psittacidae	Psittinus cyanurus	II
Psittacidae	Geoffroyus heteroclitus	II
Psittacidae	Prioniturus discurus	II
Psittacidae	Coracopsis nigra	II
Psittacidae	Diopsittaca nobilis	II
Psittacidae	Aratinga wagleri	II
Psittacidae	Aratinga nana	II
Psittacidae	Pyrrhura picta	II
Psittacidae	Pyrrhura melanura	II
Psittacidae	Myiopsitta monachus	II
Psittacidae	Forpus xanthopterygius	II
Psittacidae	Pionites leucogaster	II
Psittacidae	Pionus menstruus	II
Psittacidae	Pionus tumultuosus	II
Psittacidae	Amazona autumnalis	II
Psittacidae	Amazona festiva	II
Psittacidae	Amazona farinosa	II
Psittacidae	Cyclopsitta guliemitertii	II
Psittacidae	Cyclopsitta diophthalma	I/II
Psittacidae	Psittaculirostris desmarestii	II
Tytonidae	Tyto tenebrosa	II
Tytonidae	Tyto aurantia	II
Tytonidae	Phodilus badius	II
Strigidae	Megascops guatemalae	II
Strigidae	Otus manadensis	II
Strigidae	Otus senegalensis	II
Strigidae	Otus magicus	II
Strigidae	Otus bakkamoena	II
Strigidae	Otus megalotis	II
Strigidae	Otus leucotis	II
Strigidae	Bubo virginianus	II
Strigidae	Bubo africanus	II

Strigidae	<i>Strix aluco</i>	II
Strigidae	<i>Glaucidium gnoma</i>	II
Strigidae	<i>Glaucidium brasilianum</i>	II
Strigidae	<i>Ninox novaeseelandiae</i>	I/II
Strigidae	<i>Ninox scutulata</i>	II
Strigidae	<i>Ninox philippensis</i>	II
Strigidae	<i>Ninox squamipila</i>	II
Strigidae	<i>Ninox jacquiniti</i>	II
Trochilidae	<i>Phaethornis longirostris</i>	II
Trochilidae	<i>Phaethornis bourcierii</i>	II
Trochilidae	<i>Phaethornis griseogularis</i>	II
Trochilidae	<i>Anthracothorax dominicus</i>	II
Trochilidae	<i>Stephanoxis lalandi</i>	II
Trochilidae	<i>Lophornis chalybeus</i>	II
Trochilidae	<i>Cyananthus latirostris</i>	II
Trochilidae	<i>Amazilia lactea</i>	II
Trochilidae	<i>Amazilia viridigaster</i>	II
Trochilidae	<i>Amazilia viridifrons</i>	II
Trochilidae	<i>Lampornis castaneoventris</i>	II
Trochilidae	<i>Heliodoxa schreibersii</i>	II
Trochilidae	<i>Urochroa bougueri</i>	II
Trochilidae	<i>Oreotrochilus estella</i>	II
Trochilidae	<i>Coeligena torquata</i>	II
Trochilidae	<i>Coeligena bonapartei</i>	II
Trochilidae	<i>Coeligena violifer</i>	II
Trochilidae	<i>Heliangelus amethysticollis</i>	II
Trochilidae	<i>Eriocnemis luciani</i>	II
Trochilidae	<i>Oreonympha nobilis</i>	II
Trochilidae	<i>Oxypogon guerinii</i>	II
Trochilidae	<i>Schistes geoffroyi</i>	II
Bucerotidae	<i>Buceros hydrocorax</i>	II
Ramphastidae	<i>Ramphastos vitellinus</i>	II

CITES-RELEVANT LUMPS

Family	Species A	Species B
Phasianidae	<i>Lophura edwardsi</i>	<i>Lophura hatinhensis</i>
Falconidae	<i>Falco peregrinus</i>	<i>Falco pelegrinoides</i>
Psittacidae	<i>Cyanoramphus novaezelandiae</i>	<i>Cyanoramphus cookii</i>
Psittacidae	<i>Aratinga holochlora</i>	<i>Aratinga brevipes</i>
Psittacidae	<i>Pyrrhura leucotis</i>	<i>Pyrrhura griseipectus</i>
Tytonidae	<i>Tyto novaehollandiae</i>	<i>Tyto manusi</i>
Strigidae	<i>Bubo poensis</i>	<i>Bubo vosseleri</i>
Trochilidae	<i>Campylopterus curvipennis</i>	<i>Campylopterus excellens</i>
Trochilidae	<i>Chlorostilbon mellisugus</i>	<i>Chlorostilbon melanorhynchus</i>
Trochilidae	<i>Chlorostilbon poortmani</i>	<i>Chlorostilbon alicae</i>
Trochilidae	<i>Thalurania colombica</i>	<i>Thalurania fannyi</i>
Bucerotidae	<i>Penelopides affinis</i>	<i>Penelopides samarensis</i>

New nominate

Struthio camelus
Rhea pennata
Pauxi pauxi
Sarkidiornis melanotos
Falco chicquera
Pernis celebensis
Circus spilonotus
Circus cyaneus
Accipiter tachiro
Accipiter novaehollandiae
Accipiter striatus
Buteo nitidus
Buteo oreophilus
Nisaetus philippensis

Split A

Struthio molybdophanes
Rhea tarapacensis
Pauxi koepckeae
Sarkidiornis sylvicola
Falco ruficollis
Pernis steerei
Circus spilothorax
Circus hudsonius
Accipiter toussenelii
Accipiter hiogaster
Accipiter chionogaster
Buteo plagiatus
Buteo trizonatus
Nisaetus pinskeri

Chlamydotis undulata

Chlamydotis macqueenii

Goura scheepmakeri

Goura sclaterii

Trichoglossus haematodus

Trichoglossus moluccanus

Trichoglossus flavoviridis

Trichoglossus meyeri

Charmosyna papou

Charmosyna stellae

Psittacella picta

Psittacella lorentzi

Psittinus cyanurus

Psittinus abbotti

Geoffroyus heteroclitus

Geoffroyus hyacinthinus

Prioniturus discurus

Prioniturus mindorensis

Coracopsis nigra

Coracopsis barklyi

Diopsittaca nobilis

Diopsittaca cumanensis

Psittacara wagleri

Psittacara frontatus

Eupsittula nana

Eupsittula astec

Pyrrhura picta

Pyrrhura snethlageae

Pyrrhura melanura

Pyrrhura pacifica

Myiopsitta monachus

Myiopsitta luchi

Forpus xanthopterygius

Forpus spengeli

Pionites leucogaster

Pionites xanthurus

Pionus menstruus

Pionus reichenowi

Pionus tumultuosus

Pionus seniloides

Amazona autumnalis

Amazona lilacina

Amazona festiva

Amazona bodini

Amazona farinosa

Amazona guatemalae

Cyclopsitta guliemitertii

Cyclopsitta nigrifrons

Cyclopsitta diophthalma

Cyclopsitta coxeni

Psittaculirostris desmarestii

Psittaculirostris godmani

Tyto tenebricosa

Tyto multipunctata

Tyto aurantia

Tyto almae

Phodilus badius

Phodilus assimilis

Megascops guatemalae

Megascops vermiculatus

Otus manadensis

Otus mendeni

Otus senegalensis

Otus feae

Otus magicus

Otus jolandae

Otus bakkamoena

Otus lettia

Otus megalotis

Otus nigrorum

Ptilopsis leucotis

Ptilopsis granti

Bubo virginianus

Bubo magellanicus

Bubo africanus

Bubo cinerascens

<i>Strix aluco</i>	<i>Strix nivicola</i>
<i>Glaucidium gnoma</i>	<i>Glaucidium californicum</i>
<i>Glaucidium brasilianum</i>	<i>Glaucidium tucumanum</i>
<i>Ninox novaeseelandiae</i>	<i>Ninox boobook</i>
<i>Ninox scutulata</i>	<i>Ninox japonica</i>
<i>Ninox philippensis</i>	<i>Ninox spilocephala</i>
<i>Ninox squamipila</i>	<i>Ninox hypogramma</i>
<i>Ninox jacquinoti</i>	<i>Ninox granti</i>
<i>Phaethornis longirostris</i>	<i>Phaethornis baroni</i>
<i>Phaethornis bourcierii</i>	<i>Phaethornis mexicanus</i>
<i>Phaethornis griseogularis</i>	<i>Phaethornis porcellae</i>
<i>Anthracothorax dominicus</i>	<i>Anthracothorax aurulentus</i>
<i>Stephanoxis lalandi</i>	<i>Stephanoxis loddigesii</i>
<i>Lophornis chalybeus</i>	<i>Lophornis verreauxii</i>
<i>Cyanthus latirostris</i>	<i>Cyanthus doubledayi</i>
<i>Amazilia lactea</i>	<i>Amazilia bartletti</i>
<i>Amazilia viridigaster</i>	<i>Amazilia cupreicauda</i>
<i>Amazilia viridifrons</i>	<i>Amazilia wagneri</i>
<i>Lampornis castaneiventris</i>	<i>Lampornis cinereicauda</i>
<i>Heliodoxa schreibersii</i>	<i>Heliodoxa whitelyana</i>
<i>Urochroa bougueri</i>	<i>Urochroa leucura</i>
<i>Oreotrochilus estella</i>	<i>Oreotrochilus stolzmanni</i>
<i>Coeligena torquata</i>	<i>Coeligena conradii</i>
<i>Coeligena bonapartei</i>	<i>Coeligena eos</i>
<i>Coeligena violifer</i>	<i>Coeligena dichroura</i>
<i>Heliangelus amethysticollis</i>	<i>Heliangelus clarisse</i>
<i>Eriocnemis luciani</i>	<i>Eriocnemis sapphiropygia</i>
<i>Oreonympha nobilis</i>	<i>Oreonympha albolimbata</i>
<i>Oxypogon guerinii</i>	<i>Oxypogon stubelii</i>
<i>Schistes geoffroyi</i>	<i>Schistes albogularis</i>
<i>Buceros hydrocorax</i>	<i>Buceros mindanensis</i>
<i>Ramphastos vitellinus</i>	<i>Ramphastos culminatus</i>

Species C

Relevant CITES listings

	I
	I
<i>Cyanoramphus saisseti</i>	I
	II
<i>Pyrrhura pfrimeri</i>	II
<i>Tyto sororcula</i>	II
	II
	II
	II
	II
	II
	II
	II

Split B

Split C

Split D

Accipiter sylvestris
Accipiter ventralis

Accipiter erythronemius

Trichoglossus rubritorquis

Trichoglossus weberi

Trichoglossus capistratus

Pyrrhura parvifrons

Pyrrhura peruviana

Pionites xanthomerus

Amazona diadema

Cyclopsitta amabilis

Cyclopsitta melanogenia

Psittaculirostris cervicalis

Otus sulaensis
Otus pamela
Otus tempestatis
Otus lempiji
Otus everetti

Otus socotranus

Otus semitorques

Glaucidium cobanense

Glaucidium hoskinsii

Ninox leucopsis

Ninox randi

Ninox leventisi

Ninox hantu

Ninox malaitae

Ninox obscura

Ninox reyi

Ninox forbesi

Ninox roseoaxillaris

Ninox rumseyi

Cynanthus lawrencei

Coeligena eisenmanni

Coeligena consita

Coeligena albicaudata

Heliangelus spencei

Coeligena inca

Coeligena osculans

Oxygogon lindenii

Oxygogon cyanolaemus

Ramphastos citrolaemus

Ramphastos ariel

New lumped species

Lophura edwardsi

Falco peregrinus

Cyanoramphus novaezelandiae

Psittacara holochlorus

Pyrrhura leucotis

Tyto novaehollandiae

Bubo poensis

Campylopterus curvipennis

Chlorostilbon mellisugus

Chlorostilbon poortmani

Thalurania colombica

Penelopides affinis


Split E

Split F



Trichoglossus forsteni

Trichoglossus rosenbergii



Ninox spilonota

Ninox mindorensis