

## CONSIDERATION OF PROPOSALS FOR AMENDMENT OF APPENDICES I AND II

A. Proposal

Transfer of *Sclerocactus spinosior* spp. *blainei* from Appendix II to Appendix I.

B. Proponent

The United States of America.

C. Supporting statement1. Taxonomy

1.1 Class: Dicotyledoneae

1.2 Order: Caryophyllales

1.3 Family: Cactaceae

1.4 Species: *Sclerocactus spinosior* spp. *blainei* (Welsh & Thorne) Hochstetter 1995

1.5 Scientific synonyms: *Sclerocactus blainei* Welsh & Thorne 1985  
*Sclerocactus schlesseri* Heil & Welsh 1986  
*Pediocactus spinosior* spp. *blainei* (Welsh & Thorne) Halda 1998  
*Pediocactus spinosior* spp. *schlesseri* (Heil & Welsh) Halda 1998

1.6 Common names: English: Blaine's pincushion, Blaine's fishhook cactus, Desert valley fishhook cactus, Great Basin eagle-claw cactus, Schlessers' pincushion, Spinier devil's-claw cactus

French:  
 Spanish:

1.7 Code numbers:

2. Biological parameters

## 2.1 Distribution

*S. spinosior* spp. *blainei* is reported as occurring in Nye and Lincoln Counties, Nevada, and Iron County, Utah, United States of America (Anderson, 2001; Kartesz, 1999). The Utah Natural Heritage Program does not consider the taxon to be present in Utah (Franklin, pers. comm., 2002). Taxonomic uncertainty regarding this taxon confounds the issues related to its distribution. According to Dr. Morefield, Nevada Natural Heritage Program, "the descriptions and circumscriptions of *S. blainei*, *S. nyensis*, and *S. schlesseri* are highly inconsistent within and among the recent treatments" (NatureServe, 2002). The treatment given in the CITES Cactaceae Checklist has been followed for the purposes of this proposal. In addition the taxon will be considered as occurring in Utah until the disagreement over distribution has been definitely resolved.

## 2.2 Habitat availability

*S. spinosior* spp. *blainei* is found between 1460 and 1830 meters in elevation (NatureServe, 2002). It is found on alkaline calcareous and volcanic gravelly-clay soils in open valley bottom areas in the shadscale and lower sagebrush zones. Associated species include *Sarcobatus vermiculatus*,

*Pleuraphis jamesii*, *Atriplex confertifolia*, *Artemisia tridentata*, and *Ericameria nauseosa* (Nevada Natural Heritage Program, 2001).

### 2.3 Population status

*S. spinosior* spp. *blainei* is known from about ten locations in Nevada (Morefield, 2002). The total number of individuals is unknown. No information is available on the status of the species in Utah. *S. spinosior* spp. *blainei* is listed as Endangered by the IUCN (Walter and Gillett 1998).

*S. spinosior* spp. *blainei* is ranked as S1 and G1. The ranking of S1 and G1 are applied to taxa that have been identified as critically imperiled in the state (S1) and globally (G1). These taxa are at great risk of extirpation due to extreme rarity, imminent threats, and/or biological factors (NatureServe 2002). Furthermore, the taxon is listed as Special Status Sensitive Species on lands managed by the U.S. Bureau of Land Management (<http://www.blm.gov/nhp/efoia/nv/98im/nvim98-013.htm>).

### 2.4 Population trends

Currently unknown. However, Dr. J. Lüthy of the Management Authority of Switzerland, reports that specimens of *S. spinosior* spp. *blainei* in the wild are rare and hard to find, based on personal observation (Lüthy, 1999).

### 2.5 Geographic trends

Specific information about the geographic trends of *S. spinosior* spp. *blainei* is not available. However, potential habitat for *S. spinosior* spp. *blainei* in Nevada may be affected by agricultural and/or industrial development, off-road vehicle use, and highway maintenance (NatureServe, 2002).

### 2.6 Role of the species in its ecosystem

*S. spinosior* spp. *blainei* usually occurs as solitary plants, 3-15 cm tall. Flowers are reddish purple to violet or lavender and are borne at the tops of the stems in the late spring (Nevada Natural Heritage Program, 2001).

An important aspect of cactus flowers is the presence of different syndromes or sets of characters facilitating pollination by bats, birds, or insects. Red flowers are frequent in regions where bird pollination is common (Anderson, 2001). Cactus fruits are a source of food for numerous animals, including many species of birds. The most common method of cactus fruit or seed dispersal is by birds (Anderson, 2001).

### 2.7 Threats

*S. spinosior* spp. *blainei* is desirable and vulnerable to hobby collectors and poachers (Lüthy, 1999; Morefield, 2002). The seeds of *S. spinosior* spp. *blainei* have become increasingly available on several European web sites (Lüthy, 1999). The Nevada Natural Heritage Program receives requests from individuals in Europe for population location information for *S. spinosior* spp. *blainei* (Morefield, 2002). Most species in the genus have been adversely affected by illegal collection (Anderson, 2001).

## 3. Utilization and trade

### 3.1 National utilization

The seeds of *S. spinosior* spp. *blainei* are commercially available on the Internet.

### 3.2 Legal international trade

The U.S. CITES Annual Report data show no exports of wild specimens of *S. spinosior* spp. *blainei* from 1994 to 2000 (U.S. Fish and Wildlife Service, 2002). Seeds from *S. spinosior* spp. *blainei* are available on the Internet from web sites located in the Austria, the Czech Republic, Germany, Malta, and the Netherlands.

### 3.3 Illegal trade

There is no documentation of illegal international trade in wild-collected specimens from the United States of America.

### 3.4 Actual or potential trade impacts

The potential trade impacts to *S. spinosior* spp. *blainei* include the over-harvest of wild plants and seeds. Because these are slow-growing plants from small populations, unregulated harvest of individual plants may affect the long-term survival of the taxon.

### 3.5 Artificial propagation for commercial purposes (outside country of origin)

Although *S. spinosior* spp. *blainei* is commercially available, the genus *Sclerocactus* is one of the most difficult to propagate (Anderson 2001).

## 4. Conservation and Management

### 4.1 Legal status

#### 4.1.1 National

*S. spinosior* spp. *blainei* is listed on the State of Nevada Rare Species List, and also listed as a Special Status Sensitive Species by the U.S. Bureau of Land Management. The U.S. Lacey Act (described in 4.3.2.) provides protection against the illegal trade of *S. spinosior* spp. *blainei*.

#### 4.1.2 International

*S. spinosior* spp. *blainei* has been listed in CITES Appendix II since 1975.

### 4.2 Species management

#### 4.2.1 Population monitoring

*S. spinosior* spp. *blainei* has not yet been systematically surveyed for in Nevada (Nevada Natural Heritage Program, 2001).

#### 4.2.2 Habitat conservation

*S. spinosior* spp. *blainei* occurs on land managed by the U.S. Bureau of Land Management (BLM) in Nevada, and is listed on the BLM State Director's Sensitive Species List (<http://www.blm.gov/nhp/efoia/nv/98im/nvim98-013.htm>).

#### 4.2.3 Management measures

According to BLM regulations, collection of "sensitive" plants may be permitted only for scientific or educational purposes, or conservation or propagation of the species, and must

be authorized by a BLM permit (BLM Manual 6840.06C.) In addition, the State of Nevada regulates the harvest of *S. spinosior* spp. *blainei* (Nevada Natural Heritage Program, 2001).

#### 4.3 Control measures

##### 4.3.1 International trade

*S. spinosior* spp. *blainei* has been subject to the trade controls of CITES Appendix II since 1975. Thus, all exports of living or dead plants require the issuance of an export permit by the country of origin, and re-export require the issuance of re-export certificates by the country of re-export. However, seeds of *S. spinosior* spp. *blainei* are exempted from the listing in Appendix II by annotation.

##### 4.3.2 Domestic measures

*S. spinosior* spp. *blainei* is protected by the U.S. Lacey Act, which makes it unlawful to possess any wild plant (including roots, seeds, and other parts) that is indigenous to any State and which is either listed in an appendix to the Convention on International Trade in Endangered Species of Wild Fauna and Flora, or listed pursuant to any State law that provides for the conservation of species threatened with extinction. According to the Act, it is unlawful to import, export, transport, sell, receive, acquire, or purchase in interstate or foreign commerce any wild plant (including roots, seeds, and other parts) taken, possessed, transported, or sold in violation of any State law or regulation. It is also unlawful to import, export, transport, sell, receive, acquire, or purchase any wild plant (including roots, seeds, and other parts) taken or possessed in violation of any U.S. law, treaty, or regulation or in violation of Indian tribal law (<http://www.xmission.com/~gastown/herpmed/lacey.htm>).

The Nevada Natural Heritage Program protects location information in Nevada for *S. spinosior* spp. *blainei* because the species is considered especially vulnerable to poaching, vandalism, harassment, and hobby collecting.

#### 5. Information on Similar Species

Currently, taxonomic uncertainty regarding the treatment of *Sclerocactus blainei* exists. Some consider *Sclerocactus spinosior* (Engelmann) Woodruff & L. D. Benson to include *Sclerocactus blainei* S. L. Welsh & K. Thorne, *Sclerocactus spinosior* spp. *blainei* (S. L. Welsh & K. Thorne) Hochstätter, and *Sclerocactus schlesseri* K. D. Heil & S. L. Welsh (Anderson, 2001). Others treat *Sclerocactus blainei* and *Sclerocactus schlesseri* as separate species (e.g., Nevada Natural Heritage Program), or refer to *S. blainei* as a local variant of *S. spinosior* that differs in details of spination only (Hunt, 2002).

*Sclerocactus blainei* is considered an unusual *Sclerocactus*, because some specimens have long ribbon-like spines reminiscent of *S. papyracanthus*. The taxon *S. blainei* differs from other sclerocacti in the region in that the flowers are slightly larger and the upper central spine is longer than those of either *S. pubispinus* or *S. spinosior*. However, a few documented specimens of *S. blainei* are reported to be similar in appearance to *S. pubispinus* and *S. spinosior*. The central spine number varies, and the number of hooked spines may number as many as six, which is characteristic of *S. polyancistrus spinosior* (Porter, 2002).

A few documented specimens of *S. nyensis* have been reported to be similar in appearance to *S. blainei*, *S. pubispinus*, and *S. spinosior* (Porter, 2002).

#### 6. Other Comments

-----

7. Additional Remarks

-----

8. References

- Anderson, E. F. 2001. The Cactus Family. Timber Press, Portland, Oregon. 776 pp.
- Franklin, B. Utah Natural Heritage Program. 2002. *In litt.* to the Division of Scientific Authority, U.S. Fish and Wildlife Service.
- Hunt, D. International Cactaceae Systematics Group. 2002. *In litt.* to the Division of Scientific Authority, U.S. Fish and Wildlife Service.
- Kartesz, J.T. 1999. A Synonymized Checklist and Atlas with Biological Attributes for the Vascular Flora of the United States, Canada, and Greenland. First Edition. In: Kartesz, J.T., and C.A. Meacham. Synthesis of the North American Flora, Version 1.0. North Carolina Botanical Garden, Chapel Hill, NC.
- Lüthy, J. Switzerland Management Authority. 1999. *In litt.* to the Division of Scientific Authority, U.S. Fish and Wildlife Service.
- Morefield, J. Nevada Natural Heritage Program. 2002. *In litt.* to the Division of Scientific Authority, U.S. Fish and Wildlife Service.
- NatureServe: An online encyclopedia of life [web application]. 2001. Version 1.5 . Arlington, Virginia, USA: Association for Biodiversity Information. Available: <http://www.natureserve.org/>. (Accessed:XXX, 2002).
- Nevada Natural Heritage Program. 2001. Rare Plant Fact Sheet: *Sclerocactus Blainei*.
- Porter, J.M. Rancho Santa Ana Botanic Garden. 2002. *In litt.* as attachment to the Division of Scientific Authority, U.S. Fish and Wildlife Service. Unpublished treatment in Flora of North America, Vol. 4.
- Walter, K. S. and Gillett, H. J. [eds]. 1998. *1997 IUCN Red List of Threatened Plants*. Compiled by the World Conservation Monitoring Centre. IUCN – The World Conservation Union, Gland, Switzerland and Cambridge, UK. Ixiv + 862pp.