

CONVENTION ON INTERNATIONAL TRADE IN ENDANGERED SPECIES
OF WILD FAUNA AND FLORA

Seventeenth meeting of the Conference of the Parties
Johannesburg (South Africa), 24 September – 5 October 2016

CONSIDERATION OF PROPOSALS FOR AMENDMENT OF APPENDICES I AND II

A. Proposal

Inclusion of the genus *Dalbergia* in CITES Appendix II with exception to the species included in Appendix I.

The UNEP-WCMC assessed the *Dalbergia* species of Latin America and concluded: "... all populations of *Dalbergia* spp. from South and Central America appear to meet the criteria for listing in CITES Appendix II" (UNEP-WCMC, 2015). Including the whole genus in Appendix II will be essential for the control of international trade by eliminating the arduous task of enforcement and customs officers of differentiating between the hundreds of *Dalbergia* species listed and not listed in CITES. The inclusion will help ensure that legal trade does not become a direct cause of the extinction of these highly threatened species and will help curb illegal trade.

Considering that CITES Appendix II must include all species, which although not necessarily now threatened with extinction may become so unless trade in specimens of such species is subject to strict regulation in order to avoid utilization incompatible with their survival, it is important to include the genus *Dalbergia* in CITES Appendix II.

- a) Resolution Conf. 9.24, Annex 2 a, Criterion A - "It is known, or can be inferred or projected, that the regulation of trade in the species is necessary to avoid it becoming eligible for inclusion in Appendix I in the near future".
- b) Resolution Conf. 9.24, Annex 2 a, Criterion B - "It is known, or can be inferred or projected, that regulation of trade in the species is required to ensure that the harvest of specimens from the wild is not reducing the wild population to a level at which its survival might be threatened by continued harvesting or other influences".
- c) Resolution 9.24 (Rev. CoP15) Annex 2b Criterion A – "The specimens of the species in the form in which they are traded resemble specimens of a species included in Appendix II under the provisions of Article II, paragraph 2 (a), or in Appendix I, so that enforcement officers who encounter specimens of CITES-listed species are unlikely to be able to distinguish between them".

Domestic and international experience has indicated that enforcement and customs officers who encountered specimens of *Dalbergia* products are unlikely to be able to reliably distinguish between the various species of *Dalbergia*.

B. Proponent

Argentina, Brazil, Guatemala and Kenya².

¹ This document has been provided in these languages by the author(s)

² The geographical designations employed in this document do not imply the expression of any opinion whatsoever on the part of the CITES Secretariat (or the United Nations Environment Programme) concerning the legal status of any country, territory, or area, or concerning the delimitation of its frontiers or boundaries. The responsibility for the contents of the document rests exclusively with its author.

C. Supporting statement

1. Taxonomy

- 1.1 Class: Magnoliopsida
- 1.2 Order: Fabales
- 1.3 Family: Leguminosae (Fabaceae) Juss. 1789
- 1.4 Genus, species or subspecies, including author and year: *Dalbergia* L.f. (Grandtner, 2005)³
- 1.5 Scientific synonyms: According to GRIN (2014) the following genera are accepted as synonyms for *Dalbergia*
Amerimnon P. Browne
Coroya Pierre
Ecastaphyllum P. Browne
Miscolobium Vogel
Triptolemea Mart.
- 1.6 Common names⁴:: English: Rosewood, Palisander
French: Palissandre; bois de rose
Spanish: Palo de Rosa, Rosul, Granadillo, Ébano, Corazón Bonito, Cocobolo, Palisandro
- 1.7 Code numbers:

2. Overview

Species of the genus *Dalbergia* belong to the family Leguminosae (Fabaceae) Juss.. The genus includes trees, shrubs and vines, approximately 250 species in the tropics (Mabberley, 2008) and 304 species worldwide (The Plant List, 2013) (see annex 1).

These species are in high demand in the international tropical hardwood trade for the manufacture of furniture. However, the wild populations of several species have experienced a major decline due to legal and illegal overexploitation. CITES regulates 61 species of *Dalbergia*: *D. abrahamii*, *D. andapensis*, *D. aurea*, *D. baronii*, *D. bathiei*, *D. bojeri*, *D. brachystachya*, *D. bracteolata*, *D. calycina*, *D. campenonii*, *D. capuronii*, *D. catipenonii*, *D. chapelieri*, *D. chlorocarpa*, *D. cochinchinensis*, *D. cubilquitzensis*, *D. darienensis*, *D. davidii*, *D. delphinensis*, *D. emimensis*, *D. erubescens*, *D. gautieri*, *D. glaberrima*, *D. glaucocarpa*, *D. glomerata*, *D. granadillo*, *D. greveana*, *D. hildebrandtii*, *D. hirticalyx*, *D. humbertii*, *D. hutibertii*, *D. lemurica*, *D. louvelii*, *D. madagascariensis*, *D. malacophylla*, *D. manongarivensis*, *D. maritima*, *D. masoalensis*, *D. mollis*, *D. monticola*, *D. neoperrieri*, *D. nigra*, *D. normandii*, *D. occulta*, *D. orientalis*, *D. peltieri*, *D. pervillei*, *D. pseudobaronii*, *D. pseudoviguieri*, *D. purpurascens*, *D. retusa*, *D. stevensonii*, *D. suaresensis*, *D. trichocarpa*, *D. tricolor*, *D. tsaratananensis*, *D. tsiandalana*, *D. tucurensis*, *D. urschii*, *D. viguieri*, *D. xerophila*.

While some identification guides for a few *Dalbergia* species exist, distinction between and identification of individual species is very difficult for non-professionals and sometimes even for experts, making it a problem for enforcement and customs officers to comply correctly with inspection and identification of CITES listed *Dalbergia* tree and product shipments.

However, *Dalbergia* species at genus level can be distinguished from other “rosewood” timber genus.

The genus *Dalbergia* is native to the tropical regions of Central and South America, Africa, Madagascar and southern Asia. The distribution of this genus is fragmented and many species populations are in decline mainly due to the loss of forest coverage by human induced disturbances (e.g. non-sustainable agricultural practices, population growth, fires, legal and illegal logging). Several timber tree species of

³ Taxonomic circumscription of the genus is subject to much debate.

⁴ The genus *Dalbergia* has a larger number of trade and vernacular names, depending on the species and the region on which the species occur.

Dalbergia produce fine timbers of high economic value, generally known as “rosewood”, so renowned for its fragrances and colours, used in musical instruments and expensive furniture (e.g. *D. cochinchinensis*, *D. latifolia*, *D. melanoxylon*, *D. nigra*, *D. odorifera* and *D. sissoo*). However, the term rosewood has also been variously ascribed to the genera *Jacaranda*, *Guibourtia*, *Machaerium* and *Pterocarpus*.

Only the heartwood yield quality timber, but the trees have a low regeneration and they are slow in forming heartwood; even large logs often lose much of their volume when the valueless sapwood is removed. The cutting of exploitable individuals in the wild is the main cause of the absence of certain diameter classes and low density of populations.

Rosewood species are currently in high demand by consumers and logs and sawn wood are the main products in trade in most of the producing countries. As a result of an ongoing demand in the Asian market, rosewood has become seriously endangered by their abusive logging and destruction of their habitat which is leading to the decline of wild populations inside and outside protected areas.

This document suggests that the genus *Dalbergia* meets the criteria for listing in CITES Appendix II in compliance with Article II, paragraph 2(a) of the Convention and resolutions Conf. 9.24 (Rev. CoP16) Annex 2 a, Criteria A and B, and Annex 2 b Criterion A.

It is established or it is possible to deduct that regulation of trade in the species of the genus is required to ensure that the harvest of their specimens from the wild is not reducing the population to a level at which their survival might be threatened.

3. Species characteristics

3.1 Distribution

The genus *Dalbergia* has a pantropical distribution with high species diversity in Asia (ca. 119 species), in Africa (ca. 116 species) and in Central and South America (ca. 80 species) (Vatanparast *et al.*, 2013); according to Vaglica (2014), the genus *Dalbergia* is spread worldwide to 102 countries; (see figure 1 and Annex 2).

Global distribution of *Dalbergia* species




Figure 1. Global distribution of *Dalbergia* species (Source: Vaglica, 2014)

3.2 Habitat

The genus *Dalbergia* occurs in different types of ecosystems such as humid evergreen rainforest, semi deciduous rainforest, riparian forest, dry tropical forest, Miombo woodland, dry and subarid forests, savannahs, thickets and pastureland. Species are found in areas with completely different

characteristics, such as totally flat areas with dry rocky sites, areas with possibility of flooding during the rainy season(s), fertile and deep sandy clay or calcareous soils along streams (Standley & Steyermark, 1946; Carvalho, 1997; Stevens *et al.*, 2001; Linares & Sousa, 2007; Lemmens, 2008). Species can be found at altitudes from sea level up to 1.700 m.

3.3 Biological characteristics

The plants of the genus *Dalbergia* are small to medium sized trees, shrubs, or woody climbers and lianas with alternate, imparipinnate and compound leaves (Saha *et al.*, 2013); stipules often small and falling early; leaflets are alternate, rarely sub opposite, and glabrous (Saha *et al.*, 2013).

Flowers small (2 to 20mm): corolla small, rarely fragrant, usually white, light yellow or purple, with light green spot at the center of the vexillum (FAUSAC- FNPV, 2015). Inflorescences are terminal or axillar, racemes or panicles, usually numerous flowered; bracts and bracteoles usually small, rarely persistent, sometimes scorpioid or corymbiform.

The fruit is a pod/legume oblong, ligulate, winged, elliptic, or strap-shaped, rarely half-moon-shaped (Carvalho, 1997; Shu, 2010). Fruits with intact seeds are wind-dispersed (Bawa & Webb, 1984). The fruit has in most of the species between 1 to 3 seeds, with some exceptions as *D. retusa* which has from one to five seeds in the fruit (FAUSAC-FNPV, 2015).

Trees bloom between December and July in Central America after 4 to 5 years, with a second flush in August and September (e.g. *D. retusa*), while in Madagascar trees bloom from September to March. *Dalbergia* trees are slow growing. It has been estimated that trees in Africa reach a size large enough to yield a fair amount of heartwood only after 70–100 years (Lemmens, 2008). In South East Asia, heartwood of trees reaches an average of 13 cm in diameter after 20 years (CITES COP 16 Prop. 60).

Honeybees, beetles and butterflies are the typical pollination agents for the genus that requires cross-pollination (Vasudeva & Sareen, 2009). Bumblebees and wasps are also frequent visitors of the flowers; aphids have been found damaging the flowers structures (FAUSAC-FNPV, 2015).

Information on the breeding system of the genus *Dalbergia* is not available; however, some aspects of the reproductive biology show common features: high levels of seed abortion have been observed in *D. retusa* (Bawa & Webb, 1984); pollen is dispersed by bees in *D. glomerata*, *D. stevensonii* and *D. retusa* (Frankie *et al.*, 2002), and seed dispersed by wind and water in *D. retusa* and *D. stevensonii* (Marín & Flores, 2003).

Natural regeneration of some species (such as *D. retusa*) is scarce, however saplings and juveniles are numerous in areas periodically exposed to fire (Madrigal, 1993; Marín & Flores, 2003). In Guatemala only *D. retusa* seeds germinate in areas affected by fire (FAUSAC-FNPV, 2015).

A very important and common regeneration strategy for *Dalbergia* species in tropical dry forests is sprouting or coppicing. The ability to coppice is used for management systems (including plantation) of some *Dalbergia* species. Examples for well coppicing species are *D. sissoo*, *D. stevensonii* and *D. cochinchinensis*.

3.4 Morphological characteristics of the wood

The wood of the genus *Dalbergia* has a texture and specific colours that makes it highly desirable: red in color (*D. tucurensis* and *D. glabra*), dark brown (*D. calycina*), brown to dark brown with reddish stripes (*D. stevensonii*) dark brown to purplish black (*D. melanoxylon*) (Lemmens, 2008; FAUSAC-FNPV, 2015). According to Condit & Pérez (2002), the colour may change and darkens relatively quickly in the light to become ebony black. The trees that produce rosewood are also much alike: they are medium size, reaching 20-30 m tall, with irregular-shaped trunk, 1-2 m in diameter (NAS/NRC, 1979). Field studies of FAUSAC-FNPV (2015) showed in Guatemala that the biggest trunks reached 0.90 m in DBH (diameter at breast height) for *D. tucurensis*, 1.00 m in *D. calycina*, 0.77 m for *D. retusa* and 0.83 m for *D. stevensonii*.

Only the heartwood yield quality timber: it is very hard, heavy and strong. Thus in manufactured objects the wood is stable, durable and holds its shape. The heartwood is surrounded by clearly differentiated white sapwood; the sapwood, which is as dense as the heartwood, will vary in amount

depending on the species, the age of the tree and the conditions of its habitat (FAUSAC-FNPV, 2015). The trees are slow in forming heartwood and even large logs often lose much of their volume when the valueless sapwood is removed (NAS/NRC, 1979). Wastage may be as high as 70 - 80% as only the finest straight grain logs are used in making bars for marimbas and xylophones (Kline, 1980). However, the amount of figure and contrasting colour varies widely from tree to tree.

3.5 Role of the species in its ecosystem

The *Dalbergia* flowers represent an important food source for honeybees, beetles, wasps, bumble bees, butterflies and other insects (Vasudeva & Sareen, 2009; FAUSAC-FNPV, 2015). *D. retusa* is a highly attractive bee plant in Costa Rica, where up to 60 species of bees visit the flowers (Frankie *et al.*, 2002). *D. stevensonii* in Belize is a dominant component of the southern forest types (Cho & Quiroz, 2005). Mature trees are sites of wasps and bees colonies; beehives have been found in the principal trunks (FAUSAC-FNPV, 2015). Ant colonies are frequently found associated with *D. stevensonii*, especially because the use of leaves as source of food (FAUSAC-FNPV, 2015). Leaves also represent a source food for mammals, such as *Potos flavus*; birds are often seen on the crown of the trees (FAUSAC-FNPV, 2015).

Numerous epiphytes such as orchids, bromeliads, fungi, lichens, ferns, aroids and peperomias live on the branches and in the main trunks of the trees (FAUSAC-FNPV, 2015). Climbing plants are also associated to the genus *Dalbergia*.

The roots of *Dalbergia* species form nodules in symbiosis with nitrogen-fixing soil bacteria (it has been demonstrated to have aescynomenoid type of root nodules): as a result of this nitrogen-fixing symbiosis, *Dalbergia* species play an important role in the natural forest ecosystem, enhancing soil fertility (Rasolomampianina *et al.*, 2005; Sprent, 2009) and this is a requirement for biodiversity preservation (Rasolomampianina *et al.*, 2005).

Logging is likely to disturb the habitat of all the species associates, with the related consequences of road and trail building to transport the logs (Newman, 2004). The protection of the species from further unsustainable logging would allow remaining trees to continue their ecological functions.

4. Status and trends

The IUCN Red List has assessed the genus *Dalbergia* and has classified 83 of the 304 known species of *Dalbergia* (IUCN, 2016): 1 species as **Critically Endangered** (*D. intibucana*); 29 species as **Endangered** (*D. abrahamii*, *D. andapensis*, *D. annamensis*, *D. bariensis*, *D. bathiei*, *D. bojeri*, *D. brachystachya*, *D. cambodiana*, *D. capuronii*, *D. congesta*, *D. davidii*, *D. delphinensis*, *D. erubescens*, *D. glaucocarpa*, *D. gloveri*, *D. hirticalyx*, *D. humberti*, *D. intibucana*, *D. louvelii*, *D. mammosa*, *D. maritima*, *D. normandii*, *D. oliveri*, *D. peishaensis*, *D. setifera*, *D. suarensis*, *D. tsaratanensis*, *D. tsiandalana*, *D. urschii*, *D. xerophila*); 26 species as **Vulnerable** (*D. acariantha*, *D. aurea*, *D. balansae*, *D. baronii*, *D. catipenoni*, *D. chlorocarpa*, *D. cochichinensis*, *D. glaberrima*, *D. hildebrandtii*, *D. latifolia*, *D. lemurica*, *D. madagascarensis*, *D. monticola*, *D. neoperrieri*, *D. nigra*, *D. odorifera*, *D. orientalis*, *D. pseudobaronii*, *D. purpurascens*, *D. retusa*, *D. simposonii*, *D. tonkinensis*, *D. tricolor*, *D. vaccinifolia*, *D. viguieri*); 3 species as **Near Threatened** (*D. chapelieri*, *D. cultrate*, *D. eremicola*); 5 species as **Lower Risk/Near Threatened** (*D. bracteolata*, *D. emimensis*, *D. greveana*, *D. mollis*, *D. pervillei*); 12 species as **Least Concern** (*D. arbutifolia*, *D. assamica*, *D. calycina*, *D. cana*, *D. louisii*, *D. monetaria*, *D. oligophylla*, *D. ovata*, *D. parviflora*, *D. peltieri*, *D. reniformis*, *D. rimosa*); 1 species as **Lower Risk/Least Concern** (*D. thrichocarpa*); 6 species as **Data Deficient** (*D. boniana*, *D. ealaensis*, *D. entadoides*, *D. funera*, *D. menoeides*, *D. sambesiaca*).

4.1 Habitat trends

In Central America the typical habitat has been destroyed and is under continuing pressure particularly from increasing agriculture, cattle ranching, palm plantations, population growth and overexploitation (FAUSAC-FPNV, 2015). The tropical dry forests of Central America, the main habitat for *D. retusa*, have been subject to human influences such as hunting and modification of the vegetation cover for a long as 11,000 years (Murphy & Lugo, 1995). Less than 0.1% of the original dry forest has conservation status in Pacific Mesoamerica (Maass, 1995).

Deforestation has been commonly observed throughout the habitat range of *D. cochinchinensis*, the natural stands of the species are found scattered only in 30 protected areas thus the habitat is

fragmented (CITES COP 16 Prop. 60). In Madagascar in addition to the damage caused by abusive logging of the *Dalbergia* species, the destruction of the habitat is worsened by various anthropological activities (slash-and-burn agriculture, extension of crop fields) (CITES COP 16 Prop. 63). In some areas, in eastern and southern Africa *Dalbergia melanoxylon* is found in the Miombo woodlands which has been highly degraded as a result of human use (Deweese *et al.*, 2011).

4.2 Population size

In Madagascar, the results obtained on the density and dendrometric features of some *Dalbergia* species show a wide variety of individuals ranging between 10 and 320 individuals per hectare. Moreover, the biovolume and the basal area are low. This indicates that most individuals are not yet exploitable (DBEV, 2010). In an inventory of *Dalbergia melanoxylon* in Nachingwea, southern Tanzania, the species accounted for 4% of tree stems of the sampling area (Opulukwa *et al.*, 2002). The density in southern Tanzania corresponds well with the density found in the two forest types in which *D. melanoxylon* occurs in Mozambique with 3% and 5% respectively (Jenkins *et al.*, 2002).

In Thailand, it was estimated that the country had just 80,000-100,000 *D. cochinchinensis* trees (approximately 63,500 m³) left in the forests in 2011. The population size of rosewood in Vietnam has been declining about 50-60% during the past 5-10 years (CITES COP 16 Prop. 60).

According to FAUSAC-FNPV (2015), in Guatemala a few (only four) populations of *D. stevensonii* have been found at the moment at the so called "Franja Transversal del Norte, FTN" (Alta Verapaz and Izabal), with a range from 44 to 800 trees (only 5% of trees have a diameter between 60 and 100 cm DBH). In the Department of Santa Rosa one population of *D. calycina* was identified, around 100 trees (18% of the trees have a diameter between 40 and 80 cm DBH) and scattered trees have been found in all the Department.

Dalbergia retusa was widely distributed at the coastal planes in the South Pacific regions of Guatemala (Standley & Steyermark, 1946). However, only one population of this species has been found in Suchitepéquez (48 trees), and few scattered trees have been found in Santa Rosa and Escuintla. This species shows a high percentage of trees (69%) in the lowest diameter classes range (0-20 cm DBH) indicating recent regeneration, and only 21 % belong to the range 20-40 cm DBH, confirming a high level of over exploitation in the country.

According to FAUSAC-FNPV (2015), in Guatemala, for *Dalbergia tucurensis* only scattered trees have been found in Alta Verapaz and Quiché, of which only 12 % of trees belong to the range of 60-100 cm DBH.

4.3 Population structure

In Madagascar the population structure of the *Dalbergia* species presents a disruption caused by the absence of certain diameter classes inside and outside the protected areas (CITES COP 16 Prop. 63). An inventory of *Dalbergia melanoxylon* in Nachingwea, southern Tanzania, revealed the patchy distribution of the species. The highest amount of trees was found in the two smallest diameter classes up to 20 cm and the lowest in the class bigger than 40 cm, showing the influence of past harvests (Opulukwa *et al.*, 2002).

According to FAUSAC-FNPV (2015), only trees with a diameter class between 20 and 60 cm have been found of *Dalbergia spp.* in the wild, mainly localised in two regions of Guatemala: Alta Verapaz and Peten. The biovolume (m³/ha) and the basal area (m²/ha) are low, and this indicates that most individuals are not exploitable.

4.4 Population trends

Opulukwa *et al.* (2002) noted in a study on a region in southern Tanzania that it was now very hard to find *Dalbergia melanoxylon* trees of a harvestable size (commonly as from a DBH of 24 cm), with reductions in the number and volume of trees due to excessive exploitation. In Madagascar there is a decrease in the number of exploitable individuals in the production areas (CITES COP 16 Prop. 63).

In Thailand, it was estimated that the country had 300,000 natural stands in 2005, but greatly reduced to just 80,000-100,000 trees (approximately 63,500 m³) in 2011. Evidence suggests that the species is threatened with extinction as EIA reported that a major Rosewood trader complained in April 2011

that “the species is finished ... there are only about five years left in the trade.” In Vietnam, the species has been exposed to high rates of exploitation of the prime timber (CITES COP 16 Prop. 60).

According to FAUSAC-FPNV (2015), the combination of forest cover loss and cutting has resulted in a decline of the populations of the species of the genus *Dalbergia* in Guatemala. The wild populations of *Dalbergia* spp. (namely *D. retusa*, *D. tucurensis* and *D. stevensonii*) are likely to be severely diminished as a result of heavy logging and land-change effects during the period 1991-2012.

4.5 Geographic trends

In general, the rate and extent of deforestation in the range States is very high. FAO (2005) reported that the annual rates of forest cover change between -0.4% (Colombia) and -4.6% (El Salvador) for the range States between 1990 and 2000. Between the period 2006-2010, in Guatemala there was a loss in forest covers of 500.210 hectares with a deforestation rate estimated at 1% per year. The main areas affected by deforestation for illegal logging were Verapaz, Petén, Quiché and the central regions Chimaltenango, Guatemala and Santa Rosa which includes the distribution areas of *D. stevensonii*, *D. retusa*, *D. tucurensis* and *D. calycina* (INAB, CONAP, UVG & URL, 2012).

Mexico lost an average of 274,450 ha or 0.39% per year between 1990 and 2010 or a 7.8% total loss of forest cover while in Brazil, the change in forest cover between 1990 and 2010 was a loss of 2,765,850 ha or 0.48% per year, in total a loss of 9.6% of the forest cover (Butler, 2016)

In Asia, deforestation has been commonly observed throughout the habitat range of *D. cochinchinensis*. In Thailand, the habitat area has been continuously reduced due to both deforestation for agriculture and illegal logging. The natural stands of the species are found scattered only in 30 protected areas of 557.76 km². The habitat is thus fragmented (CITES COP 16 Prop. 60). In Malaysia, the change in forest cover between 1990 and 2010 was a loss of 96,000 ha or 0.43% per year, in total a loss of 8.6% of the forest cover (Butler, 2016).

In Africa, Miombo woodlands stretch across southern Africa in a belt from Angola and the Democratic Republic of Congo in the west to Mozambique in the east. In some areas, Miombo has been highly degraded as a result of human use (Deweese *et al.*, 2011). This is also the biome predominant in the countries (Mozambique and Tanzania) from where most of the *Dalbergia melanoxylon* timber is exported (Lemmens, 2008). In Madagascar in addition to the damage caused by abusive logging of the *Dalbergia* species, the destruction of the habitat is worsened by various anthropological activities (slash-and-burn agriculture, extension of crop fields) (CITES COP 16 Prop. 63).

5. Threats

The genus *Dalbergia* is known to be in decline as a result in the extraction of the species for its valuable timber. These woods have a long tradition of use in musical instruments, like marimbas and guitars, wood carvings and currently subject to high demand for luxury furniture trade (Lemmens, 2008, Jenkins *et al.*, 2012). Many species have become endangered or vulnerable due to the overexploitation for its valuable wood like *D. annamensis*, *D. bariensis*, *D. cambodiana*, *D. mammosa*, *D. oliveri* (Nghia, 1998), *D. latifolia* (Asian Regional Workshop, 1998), *D. odorifera* (WCMC, 1998), *D. tonkinensis* (Ban, 1998), as well as the 61 *Dalbergia* species listed in CITES.

The habitat of the genus in Guatemala is known to be affected by a range of threats, and the principal processes that affect forests are: shifting agriculture (palm oil - *Elaeis* spp., coffee - *Coffea* spp. and rubber plantations - *Hevea* spp.), forest fires, use of fuel wood and the illegal extraction and over-exploitation of species with high market values (both nationally and internationally), which represent up to the 95% of deforestation rate, and the extraction of timber by rural communities for local use (Vivero *et al.*, 2005; IARNA-URL, 2009).

In Mexico and Costa Rica, cattle farming and road construction are opening up areas and making accessible for logging sites where *D. calycina* occurs (Groom, 2012).

6. Utilization and trade

6.1 National utilization

Timber species of *Dalbergia* of South America (e.g. *D. brasiliensis*, *D. cearensis*, *D. cubilquitzensis*, *D. cuscatlanica*, *D. decipularis*, *D. foliolosa*, *D. funera*, *D. glomerata*, *D. hortensis*, *D. miscolobium*, *D. spruceana*, *D. villosa*) provide some of the most highly prized woods for the manufacture of fine furniture, marquetry and veneers, also used in cabinetwork, musical instruments and sculptures for local markets and handicrafts (Harris, 2004; Grandtner & Chrevrette, 2013; ILDIS, 2014; INAB-SEINEF, 2015).

In Africa it is sometimes used for rafters and poles in construction, and for implements such as walking sticks, hammers, drumsticks, arrow tips, pestles, cups, plates and combs. The wood is also used for charcoal production and as firewood, although the flame is very hot and may damage cooking pots. In Senegal the stem and root bark is used in traditional medicine to treat diarrhoea in combination with baobab or tamarind fruits. The smoke of burnt roots is inhaled for treatment of headache, bronchitis and colds. In Sudan, patients suffering from rheumatism are exposed to the smoke of burnt stems. In East Africa, a root decoction is used to prevent miscarriage, as an anthelmintic and aphrodisiac, and to treat gonorrhoea, stomach-ache and abdominal pain. A bark decoction or bark powder is used to clean wounds and a leaf decoction to relieve pain in the joints. Leaf sap is taken to treat inflammations in mouth and throat. Bark decoctions and leaf sap are also ingredients of mixtures used to treat various complaints (Lemmens, 2008).

In South East Asia *D. cochinchinensis* wood is used to make furniture, carvings, wood turnery, fine-art articles, musical instruments and sewing machines. The wood from the stumps and roots can also be used for making handicrafts. Root, bark and sap can be part of traditional medicine. The distinctive heartwood makes beautiful patterns after sawing (CITES COP 16 Prop. 60.)

6.2 Legal trade

Rosewood species are exported as raw logs or finished products, furniture and handicrafts. In Madagascar, over 90% of exported products were logs and sawn wood (CITES COP 16 Prop. 63). Heartwood from old trees is valued for having the richest colouration (Zadro, 1975). The burls, highly figured cambium outgrowths, are particularly valued by some timber traders.

Trade in timber products of *Dalbergia* has increased exponentially in the past few years as seen by log imports to Asian markets for the Hongmu (Red Wood) trade that is based on 33 species of tropical hardwood trees of which 16 species are *Dalbergia* (EIA, 2016) (Figure 2). Of these 16 *Dalbergia* species, 7 come from Asia (*D. odorifera*, *D. cultrata*, *D. fusca*, *D. latifolia*, *D. bariensis*, *D. cochinchinensis*, *D. oliveri*), 2 from Africa (*D. melanoxyton*, *D. louvelli*) and 7 from Latin America (*D. nigra*, *D. spruceana*, *D. stevensonii*, *D. cearensis*, *D. frutescens*, *D. granadillo*, *D. retusa*) (EIA, 2016). Only *D. cochinchinensis*, *D. louvelli*, *D. nigra*, *D. stevensonii*, *D. granadillo* and *D. retusa* are regulated by CITES.

Far more *Dalbergia* species are presently traded than those CITES-listed. However, information on trade flows is not easily available.




Figure 2. China hongmu log imports by region of origin (Source: EIA, 2016)

According to the CITES Trade Database , a total of 28,288.45 m³ of *Dalbergia* wood products and 140 t were traded between 2005 and 2014 (UNEP-WCMC, 2016a); the number of *Dalbergia* species in trade increased from 1 to 6 (due to the fact that new species were listed in the CITES Appendices), the number of records of trade also increased from a minimum of 64 in 2006 to a maximum of 220 in 2013 (Figure 3) and the number of importing countries fluctuated from a minimum of 20 to a maximum of 35 importing countries in 2013.




Figure 3. Annual transactions of *Dalbergia* spp. products: year range 2005-2014 (Source: UNEP-WCMC, 2016a).

According to UNEP-WCMC (2015), European importers reported trade in sawn timber of *D. nigra*, *D. retusa*, and *D. stevensonii* from South and Central America. The main exporter countries were Guatemala (*D. stevensonii*), Belize (*D. stevensonii*) and Brazil (*D. nigra*) for commercial purpose, and *D. retusa* from Costa Rica for personal purpose. Between 2004 and 2013 exports from South and Central America to the rest of the world were reported in *D. granadillo*, *D. nigra*, *D. retusa*, *D. stevensonii* and *Dalbergia* spp., except for 2005 and 2006. Commercial trade was predominantly

in timber with some additional commercial trade in veneers and carvings of *D. nigra* (UNEP-WCMC, 2015).

According to The National Forest Institute of Guatemala (INAB, 2015), for the year 2014, *Dalbergia* timber products (logs, tables, sawn wood and furniture) of *D. calycina*, *D. cubilquitzensis*, *D. stevensonii* and *Dalbergia spp.* have been commercialized and exported outside the country.

Consulted the CITES Trade Database (UNEP-WCMC, 2016b) Guatemala reported the export of 463.79 m³ of sawn wood of *Dalbergia spp.* during the period 2011-2014. All exports were done under source code "W" (wild) and with "T" purpose (commercial).

Dalbergia melanoxyton is principally viable for commercial timber extraction only in southeast Tanzania and northern Mozambique (Jenkins *et al.*, 2012). Based on official records of Mozambican timber exports, the total consumption of timber (not only *D. melanoxyton*) domestically and for export was 727,000 m³ of logs equivalent in 2012 (FAEF, 2013). Timber imports to China from Mozambique increased around seven-fold in the last 10 years, the figures given by Chang & Peng (2015) for imported *Dalbergia melanoxyton* timber (round wood equivalents) into China are more than 5,000 m³ in 2004 and more than 33,000 m³ in 2013.

6.3 Parts and derivatives in trade

In Guatemala, logs and sawn wood of *Dalbergia* timber tree species are the main products in the international trade, but wooden furniture, firewood, tables and finished items manufactured (doors, frames, board) and handicrafts are also found in the national and international trade (CONAP, 2015; INAB, 2015; UNEP-WCMC, 2016b).

With an exception for *D. nigra*, the CITES trade database shows that almost all the exports in CITES listed *Dalbergia* species reported by range states for the period 2000-2015 are identified as logs, timber and sawn wood (UNEP-WCMC, 2016c).

Woodcarvings of *D. nigra* an Appendix I species from Brazil are traded internationally (UNEP-WCMC, 2016a).

D. melanoxyton is a favourite wood for musical instruments, especially wind-instruments such as clarinets, oboes, flutes and bagpipes, because of its dark colour, stability and clearness of tone. Jenkins *et al.*, (2012) reported a stable demand for this industry of 255 m³ per year. Usually the wood for this purpose is exported semi-processed as billets. In 1999 it was estimated that 250,000 pieces of carving were exported with a value of US\$970,000. In the past Senegal, Kenya and Malawi produced considerable amounts of African blackwood, especially for the carving industry, but stands have been depleted considerably and carvers often changed to other woods or used African blackwood from Tanzania (in Kenya) or Mozambique (in Malawi) (Lemmens, 2008). Destructive use in Kitui, Kwale and Makueni has included digging up the roots to make tall thin carvings. Most carvers now buy their *Dalbergia melanoxyton* wood from traders who cross the border from Tanzania.

6.4 Illegal trade

The trade in Hongmu *Dalbergia* species has increased manifold in the past few years causing an increase in illegal trade. All Hongmu source countries in the Mekong region have strict log export bans and trade in a number of the more threatened species is completely prohibited. Despite these provisions, illegal cross-border log and sawn timber trade is evident across the Mekong (EIA, 2016). The same occurs in West Africa, where most countries have adopted total bans on harvesting and export and in Latin America, despite several moratoriums on exports and recent CITES listings, the illegal trade in Hongmu species is still rampant (EIA, 2016).

Illegal logging is a prevalent problem in Central America. For instance, it was estimated in 2003 that up to 85% of the total harvest in broadleaf forests in Honduras was illegal. There is very little information on the volume of international trade although cocobolo wood is available from numerous sources online (Jenkins *et al.*, 2012). Extensive illegal trade in rosewoods has been reported, raising concerns that it has accelerated in recent years (Jenkins *et al.*, 2012). Seizures of illegally trafficked timber in Guatemala suggest that there is an organized smuggling ring capable of exporting large quantities. The demand for *D. retusa* from the Darien Region of Panama has been described as "out of control" with "hundreds of settlers looting" the species (Jenkins *et al.*, 2012).

During the period 2011-2014, 38 shipments and vehicles, with a total amount of 906.244 m³ of *Dalbergia* timber (mainly trunks, flitches and tables) of *D. stevensonii*, *D. retusa* and *Dalbergia spp.* (reported as rosul) of illegal origin were confiscated in Guatemala (almost two times the CITES timber reported as legally exported for the same period). With the exception of two shipments destined to Honduras and El Salvador, all the other shipments were destined to Asia.

Dalbergia stevensonii was once locally common in Belize, then faced with rapidly declining stocks, Belize prohibited all raw rosewood exports in 1992 but lifted the ban in 1996. Local estimates in 2010 suggested a loss of 90 per cent of historical rosewood stands and in 2012, Belize enacted a moratorium on the harvest and export of rosewood species. Despite this progressive move, the illegal harvest of the species has continued (EIA, 2016). *D. granadillo* started to be logged in all the sierras of Michoacán, Mexico in 2012 where it is sold for US\$2,500/m³. An environment authority said that a ship container is sold to Asian buyers for US\$50,000 (PROCESO, 2014).

In Mexico the seizures of *D. granadillo* sawn wood destined for the Asian market increased twofold from 318.077 m³ in 2012 to 727.703 m³ in 2014 (PROFEPA, 2014).

In recent years Madagascar has experienced very high levels of illegal logging of ebonies and rosewoods from its rainforest national parks, particularly since a coup d'état in March 2009. This illegal logging has been described as the most severe threat to Madagascar's north-eastern rain forests. The vast quantities being felled have been valued at several hundred million US dollars' worth extracted during 2009. There are estimates of 100 – 200 trees being felled daily with a collective value of up to US\$460,000/day. Most of the rosewood being extracted is destined for Asian markets, for luxury furniture. Foreign traders and local traders dubbed 'timber barons' dominate the illegal logging with the vast majority (Jenkins *et al.*, 2012). Illegal Precious Wood trade appears from this brief look (anecdotally) to have been accelerating in the last few years, particularly for rosewood species. It may be the case that the very recent escalation in illegal rosewood extraction from Central America is due to Madagascar clamping down on the illegal rosewood trade from 2010 and therefore this supply drying up for the Asian market. Areas where rosewood used to occur closer to China have been all but exhausted of rosewood, with estimates from traders that the species has just five years left. This rosewood trade appears to have been accelerating in the last few years, particularly 'species hopping' and 'source area shifting' is a trend that has been known about and commented upon for many years (Jenkins *et al.*, 2012).

Species shifting, as species become commercially extinct, is a common practice. For instance, with the commercial extinction of *D. odorifera* in China and *Pterocarpus santalinus* in India, the trade in *D. cochinchinensis* grew rapidly and it became the most sought-after Hongmu species globally (EIA, 2016). As *D. cochinchinensis* was overexploited the main species now dominating the Hongmu trade in South-East Asia are *D. oliveri*, *D. bariensis*, *Pterocarpus macrocarpus*, and *P. pedatus* (EIA, 2016). In 2014, an estimated 229,796 m³ of *D. oliveri* logs were traded internationally (EIA, 2016). More and more species of *Dalbergia* are entering the trade worldwide as stocks of once abundant species are being depleted.

Hongmu trade is also linked to and drives violence in source and transit countries. In West Africa, Hongmu species are known as "blood timbers" due to connections between illegal Hongmu trade and rebel group uprisings; for example, in the Senegalese Casamance, in Côte d'Ivoire and in northern Nigeria in territories controlled by the Muslim extremist group Boko Haram. In Thailand, more than 150 forest rangers, police, soldiers and illegal loggers have been killed in gunfights during rosewood enforcement operations in recent years (EIA, 2016).

Traffickers exploit any legal loophole to smuggle illegal timber. Traffickers have repeatedly taken advantage of the current gaps in the CITES listings, misdeclaring *Dalbergia retusa* as the unregulated and similar-looking *Dalbergia bariensis* in violation of national moratoriums and CITES listings (EIA, 2016). In Guatemala, the documents accompanying rosewood shipments often recorded the export as recycling material (such as cardboard, junk or scrap metal) or other timber species, such as *Cupressus*, *Dialium* and *Miroxylum*. In Mexico, *D. granadillo* logs are mixed with other species in ship containers to disguise the shipments from authorities since there are no permits to log this protected species (PROCESO, 2014).

6.5 Actual or potential trade impacts

Dalbergia species have a relative low regeneration rate and the absence of certain diameter classes (70-100 cm) and the low density of population of certain diameter classes (20-30, 40-50) further

disturbs the health of regeneration. Wastage may be as high as 70- 80% as only the finest straight grain logs are used in making bars for marimbas and xylophones (Kline, 1980). The trees are slow in forming heartwood, so even large logs lose much of their volume when the sapwood is removed (NAS, 1979). The cutting of exploitable individuals in the wild is the main cause of this disturbance; harvesting for exportation could slow natural regeneration and the international trade has therefore promoted cutting of great many *Dalbergia* spp., leading to the decline of populations inside and outside protected areas.

7. Legal instruments

7.1 National

Most range states have different legal instruments for the protection and conservation of the species, their habitat, declaration of protected areas, measures controlling logging, use, import or export.

7.2 International

CITES regulates 61 species of *Dalbergia*: 1 species in Appendix I, 55 species in Appendix II and 5 species in Appendix III:

Species	Appendix		
	Appendix I	Appendix II	Appendix III
<i>D. nigra</i>	Brazil		
<i>D. cochichinensis</i>		Thailand #5	
<i>D. abrahamii, D. andapensis, D. aurea, D. baronii, D. bathiei, D. bojeri, D. brachystachya, D. bracteolata, D. campenonii, D. capuronii, D. catipenonii, D. chapelieri, D. chlorocarpa, D. davidii, D. delphinensis, D. emirnensis, D. erubescens, D. gautieri, D. glaberrima, D. glaucocarpa, D. greveana, D. hildebrandtii, D. hirticalyx, D. humbertii, D. hutibertii, D. lemurica, D. louvelii, D. madagascariensis, D. malacophylla, D. manongarivensis, D. maritima, D. masoalensis, D. mollis, D. monticola, D. neoperrieri, D. nigra, D. normandii, D. occulta, D. orientalis, D. peltieri, D. pervillei, D. pseudobaronii, D. pseudoviguieri, D. purpurascens, D. suaresensis, D. trichocarpa, D. tricolor, D. tsaratananensis, D. tsiandalana, D. urschii, D. viguieri, D. xerophila</i>		Madagascar #5	
<i>D. granadillo, D. retusa, D. stevensonii</i>		Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama #6	
<i>D. darienensis</i>			Panama #2
<i>D. calycina, D. cubilquitzensis, D. glomerata</i>			Guatemala #6
<i>D. tucurensis</i>			Nicaragua #6

Annotations: #2 All parts and derivatives except seeds and pollen and finished products packaged and ready for retail trade; #5 logs, sawn wood and veneer sheets; #6 logs, sawn wood, veneer sheets and plywood

8. Species management

8.1 Management measures

Most range states have different kinds of management measures for either the species, protected areas, logging, use or export.

8.2 Population monitoring

Very few reports have been published on population monitoring although two exist for Guatemala (Inventario Nacional de Especies Forestales Estratégica por la Convención CITES- Fase I y Fase II; FNPV, 2015). FAUSAC-FNPV (2015) is doing an ongoing population monitoring.

8.3 Control measures

8.3.1 Domestic

Most range states have different legal instruments for the protection and conservation of the species, their habitat, declaration of protected areas, logging, use, import or export.

8.3.2 International

CITES regulates 61 species of *Dalbergia*: 1 species in Appendix I, 55 species in Appendix II and 5 species in Appendix III (see section 7.2).

8.4 Artificial propagation

Due to the value of their timber, NAS (1979) recommended that effort be made to extend *Dalbergia* species cultivations.

Guatemala has registers of plantations of: *D. retusa* and *D. stevensonii* (INAB, 2012; FNPV, 2012). *D. retusa* was included in plantation trials of native precious wood species in Costa Rica, which started in 1992 (Fonseca & Chinchilla, 2002; Fonseca *et al.*, 2002). In managed plantations, trees may reach 13 cm DBH and 8 m in height after 17 years (Marín & Flores, 2003 and references therein). *Dalbergia* trees have been found to grow at an average rate of 1.1 m/year in height (Knowles & Leopold, 1997).

Laos and Thailand have trial plots of *D. cochinchinensis* (CITES COP 16 Prop. 60). In Madagascar propagation trials from cuttings and by layering produced satisfactory results for *D. monticola* (CITES COP 16 Prop 63). There are records of artificial propagation of *D. melanoxyton* in plantations in Kenya and Tanzania (Gregory *et al.*, 1999).

8.5 Habitat conservation

Many species of *Dalbergia* exist within natural protected areas in their respective range states.

9. Information on similar species

Many *Dalbergia* species have the same wood anatomy and the process of identification of different species is very difficult because of the hardness of the wood in the process of preparing thin sections for microscopic analysis (FAUSAC-FNPV, 2015; McLure *et al.*, 2015).

The timber of *D. retusa* is likely to be confused with *D. stevensoni* and *D. tucurensis* (Wiedenhoeft, 2011). *D. retusa* can be most easily separated from them by its distinct reddish-orange color, but also its comparative lack of paratracheal parenchyma and distinct abundance of diffuse-in-aggregate parenchyma; this requires strong technical knowledge (Wiedenhoeft, 2011). *D. tucurensis* and *D. stevensonii* cannot usually be definitively separated (Wiedenhoeft, 2011). The two species can be distinguished by means of density that requires accurate weight and volume measures (Wiemann & Ruffinatto, 2012). *D. retusa* wood is denser and stronger than Brazilian rosewood *D. nigra* (SCMRE, 2002). *D. tiralana* can be confused with *D. stevensonii* (Zamora, 2000).

The timber of *D. granadillo* (range States El Salvador and Mexico) is not distinguishable from that of *D. retusa* (Record & Hess, 1943; Richter, 2006). Although it has the common name "granadillo", it is often traded under the name "cocobolo" (Richter, 2006). The inclusion in CITES Appendix II of the whole genus is therefore proposed for look-alike reasons.

Mexico also made recommendations to the CITES Plants Committee for the evaluation of look-alike species from the genus, due to the difficulty that differentiation of CITES-listed and non-listed *Dalbergia* species presents for implementation of the Convention (PC22 Doc. 22.4), a problem also noted by other Parties in the region (PC22 Doc. 17.2; UNEP-WCMC, 2015).

10. Consultations

Country	Reply to consultation	Range state
Argentina	Supports the proposal and states the will to coproponent ⁵	yes
Australia	Believes that the possibility of full genus listing for <i>Dalbergia</i> could be considered along with other mechanisms to help discourage the trade and sale of illegally harvested timber.	yes
Brazil	Supports the proposal and states the will to coproponent ³	yes
Burkina Faso	Supports the proposal and states the will to coproponent ⁶	yes
China	There is no time to finalize the assessment of the proposal and collect updated information on the genus <i>Dalbergia</i> in China before the submission deadline .	yes
Colombia	Supports the proposal	yes
Comoros	Supports the proposal and states the will to coproponent ⁴	yes
Côte d'Ivoire	It was not possible to gather sufficient information on the genus <i>Dalbergia</i> to allow a decision on the proposal	yes
USA	Agree that there are identification and trade issues that warrant consideration of all non-listed <i>Dalbergia</i> species for inclusion in Appendix II. Propose several changes to the draft proposal, some of which were already considered in the final version	yes
Philippines	Supports the proposal	yes
Guatemala	Supports the proposal- coproponent	yes
Equatorial Guinea	Supports the proposal	yes
Honduras	Supports the proposal and states the will to coproponent ⁴	yes
Iraq	Informs there is no data available for the presence of this genus in Iraq and is glad to cooperate in any concerted effort in this field ⁷	yes
Malaysia	Supports the proposal	yes
Nigeria	Supports the proposal	yes
Perú	Supports the proposal	yes
Singapore	Is going to evaluate the proposal and consider supporting it at COP17	yes
Thailand	There is no time to study the impact of the proposal. It will be studied and maybe it can be supported at the COP.	yes
European Union	Supports the proposal	yes

11. Additional remarks

Trade in *Dalbergia* species has been increasing for many years and more and more species are being logged and traded internationally, most of these illegally. Several factors are causing major problems for

⁵ According to the provisional list of proposals for amendment of Appendices I and II, on the Secretariat webpage, the proponents are Argentina, Brazil, Guatemala and Kenya - <https://cites.org/eng/cop/17/prop/index.php>.

⁶ Despite the reply to the consultation, the country is not on the proponents referred on the provisional list of proposals for amendment of Appendices I and II, on the Secretariat webpage.

⁷ Refused by the Secretariat because there is no record of receiving a request for co-sponsoring the proposal before the deadline of 27th May.

enforcement authorities to comply correctly with the actual CITES listings. For example, the difficulty in differentiating between *Dalbergia* species was discussed in section 9, and many shipments are only identified as *Dalbergia* spp. for which there is no way of knowing if it includes only CITES-listed *Dalbergia* species or illegal non-listed *Dalbergia* species. It has been recognized that enforcement and tracking/reporting of *Dalbergia* species in trade is hampered by the use of common trade names, such as “rosewood” as it can relate to CITES-listed *Dalbergia* species and to many non-listed *Dalbergia* species or species from different CITES or non-listed genera (PC22 Doc 17.6).

It has been recognized as well the problem to identify between *Dalbergia* and other timber look-alike species or improve testing to differentiate between listed and non-listed *Dalbergia* species and look-alike species. For example, the genera *Pterocarpus* spp. and *Machaerium* spp. are closely related to *Dalbergia* while other look-alike genera include *Dicorynia* spp., *Caesalpinia* spp., and *Swartzia* spp. (PC22 Doc 17.6).

These and other problems have been discussed by several experts and working groups, including discussions at the last Plants Committee meeting in Tbilisi, Georgia. The Committee noted document PC22 Doc. 17.6. on the implementation of the Convention for *Dalbergia* spp., presented by the representatives from the European region. The Committee also noted document PC22 Doc. 22.4 and supported its submission for consideration by the Conference of the Parties, including recommendations 9 and 10:

- “9. To take note of the advances made by Mexico with respect to the evaluation of timber species of the genus *Dalbergia*, and also of the principal difficulties that the CITES Authorities of Mexico have encountered in the implementation of the listings of *Dalbergia* spp. in Appendix II (para. 3).
10. Considering that to date there is no reliable method (nor one tried out by Customs officials) to identify at an intra-species level the wood from specimens of *Dalbergia*, and that this may cause problems in implementing the listings of *Dalbergia* in Appendix II, to develop recommendations on the appropriateness of listing the thirteen remaining Mexican timber species in Appendix II, and in that context to consider other timber species of *Dalbergia* that might be suitable for the same treatment, in accordance with the rule on listing of look-alike species (in line with Article II, para. 2b. of the Convention).”

During the discussion of these documents a large number of participants supported the need to list the whole genus *Dalbergia* in Appendix II to safeguard many of the *Dalbergia* species which are now in trade and have yet to be listed as well as to simplify the work of enforcement and customs authorities when facing shipments of *Dalbergia* products.

The full genus listing for *Dalbergia* should be considered along with other mechanisms to help discourage the trade and sale of illegally harvested timber. Multiple mechanisms to help address this issue may be used like those referred in PC22 Doc. 17.6 on Implementation of the Convention to *Dalbergia* spp. and namely the following:

- The development of a Timber Identification Directory for CITES-listed species
- Use of the Global Timber Traffic Network (GTTN) database which will include DNA technology as a mechanism for genus identification;
- The development of robust timber legality frameworks within all Parties that encourage the trade in legally harvested timber; and
- Information sharing initiatives at sub regional, regional and an international scale encouraging enforcement agencies to share intelligence on illegal harvest, processing and trafficking of rosewood timber and timber products.

Non-endangered *Dalbergia* species would be subject to increased controls which may increase the bureaucratic burden to the trade in legally harvested timber and timber products but this is a minor inconvenience regarding the benefits resulting from the whole genus listing.

12. References

Asian Regional Workshop (Conservation & Sustainable Management of Trees, Viet Nam, August 1996).
1998. *Dalbergia latifolia*. The IUCN Red List of Threatened Species 1998: e.T3209 A9675 296 .

- <http://dx.doi.org/10.2305/IUCN.UK.1998.RLTS.T32098A9675296.en>. Downloaded on 03 December 2015.
- Ban, N.T. 1998. *Dalbergia tonkinensis*. The IUCN Red List of Threatened Species 1998: e.T32819A9732061. <http://dx.doi.org/10.2305/IUCN.UK.1998.RLTS.T32819A9732061.en>. Downloaded on 03 December 2015.
- Bawa, K.S. and Webb, C.J. 1984. Flower, fruit and seed abortion in tropical forest trees: Implications for the Evolution of Paternal and Maternal Reproductive Patterns. *American Journal of Botany*. 71(5): 736-751.
- Butler, R. 2016. Tropical rainforests: deforestation rates, tables and charts. Mongabay.com Accessed 20/01/2016 <http://rainforests.mongabay.com/deforestation/>
- Carvalho, A. M. 1997. A synopsis of the genus *Dalbergia* (Fabaceae, Dalbergieae) in Brazil. *Brittonia*. 49 (1): 87-109.
- Chang, Y. and Peng, R. 2015. Timber flow study: export/import discrepancy analysis China vs. Mozambique, Cameroon, Uganda and DRC. IIED Issue Paper. IIED, London.
- Cho, P. and Quiroz, L. 2005. Forest Department, Ministry of Natural Resources, Belmopan, Belize. <http://www.unep-wcmc.org/forest/timber/workshops/reports/MA2005.htm>.
- CITES COP16 Prop. 60. *Dalbergia cochinchinensis* (Palo de rosa de Tailandia) - Inclusión en Apéndice II
- CITES COP16 Prop. 63. *Dalbergia* spp. (Palo de rosa de Madagascar) - Inclusión de las poblaciones de Madagascar en Apéndice II
- CONAP (Consejo Nacional de Areas Protegidas). 2014. Manual de Procedimientos para el Comercio Internacional de Especies de Flora Maderable Contenidas en los Apéndices II y III de la CITES (*Swietenia macrophylla* King, *Swietenia humilis* Zuccarini, *Cedrela odorata* L, *Dalbergia retusa* Hemsl y *Dalbergia stevensonii* Standl). Consejo Nacional de Áreas Protegidas. Guatemala. 73 p.
- Condit, R. and Pérez, R. 2002. *Tree Atlas of the Panama Canal Watershed*. Center for Tropical Forest Science, Panama. Accessed 07/02/2005. <http://ctfs.si.edu/webatlas/maintreeatlas.html>
- EIA (Environmental Investigation Agency). 2016. The Hongmu Challenge: A briefing for the 66th meeting of the CITES Standing Committee, January 2016.
- DBEV, 2010. Evaluation écologique de quelques espèces de bois précieux les plus commercialisées a Madagascar (bois d'ébène, palissandre et bois de rose), Rapport final WWF, 97p.
- Deweese, P., B. Campbell, Y. Katerere, A. Siteo, A.B. Cunningham, A. Angelsen and S. Wunder. 2011. Managing the Miombo Woodlands of Southern Africa: Policies, incentives, and options for the rural poor. Washington DC: Program on Forests (PROFOR).
- FAEF (Faculty of Agronomy and Forestry Engineering). 2013. Assessment of harvested volume and illegal logging in Mozambican natural forest. Report of the Faculty of Agronomy and Forestry Engineering, Eduardo Mondlane University, with support from the Forest Law Enforcement, Governance and Trade Support Programme for African, Caribbean and Pacific Countries (GCP/INT/064/EC). Maputo.
- FAO. 2005. *State of the World's forests*. 6th edition. Food and Agriculture Organisation of the United Nations, Rome.
- FAUSAC-FNPV. 2015. ITTO-CITES Project: "Establishment of a forensic laboratory for identification and description of woods for its application to legal processes and to the systems of traceability of the products included in CITES". Preliminary data and personal consultations.
- FNPV. (Fundacion Naturaleza para la Vida). 2012. Inventario Nacional de Especies Forestales Estrategicas Protegidas por la Convencion CITES: Fase II. Proyecto Inventario Nacional de Caoba, Cedro y Rosul. Informe Final Del Proyecto F12/2011/Fonacon.
- Fonseca G.W. and Chinchilla M.O. 2002. Native species in plantation in the south Pacific region of Costa Rica. Memoria del taller-seminario: especies forestales nativas, Heredia, Costa Rica, 4-5 April 2002, 97-100. Source: CAB Abstracts.
- Fonseca G.W., Chinchilla M.O. and Guerrero R. 2002. Native species in plantation in the dry Pacific region of Costa Rica: the case of the precious woods. Memoria del taller-seminario: especies forestales nativas, Heredia, Costa Rica, 4-5 April 2002, 63-67. Source: CAB Abstracts
- Frankie, G.W., Vinson, S.B., Thorp, R.W., Rizzardi, M.A., Tomkins, M. and Newstrom-Lloyed, L.E. 2002. Monitoring: an essential tool in bee ecology and conservation. *In*: Kevan, P and Imperatriz Fonseca,

- V.L. (Eds). Pollinating bees – the conservation link between agriculture and nature. Ministry of Environment. Brasília. pp. 187-198. <http://www.webbee.org.br>
- Grandtner, M.M. 2005. Elsevier's Dictionary of Trees. Volume 1 North America. Elsevier Science.
- Grandtner, M.M., Chevrette. 2013. Dictionary of Trees, Volume 2: South America: Nomenclature, Taxonomy and Ecology. Academic Press.
- Gregory, A-M., Ball, S.M.J. and Eziefula, U.E. 1999. Tanzanian Mpingo 98 Full Report. Mpingo Conservation Project.
- GRIN. 2014. Germplasm Resource Information Network. U.S. Department of Agriculture /Agricultural Research Services National Plant Germplasm System. <https://npgsweb.ars-grin.gov/gringlobal/taxonomygenus.aspx?id=3367>
- Groom, A. 2012. *Dalbergia glomerata* The IUCN Red List of Threatened Species. Version 2014.3. <www.iucnredlist.org>. Downloaded on 23 April 2015.
- Harris, S. 2004. Woody legumes (excluding Acacias). Tropical Forest. Elsevier Ltd. Pp:1793-1798.
- IARNA-URL. (Instituto de Agricultura, Recursos Naturales y Ambiente de la Universidad Rafael Landívar). 2009. Perfil Ambiental de Guatemala 2008-2009. Guatemala. 319 pp.
- IARNA-URL. Instituto de Agricultura, Recursos Naturales y Ambiente de la Universidad Rafael Landívar. 2006. Perfil Ambiental de Guatemala 2006: Tendencias y reflexiones sobre la gestión ambiental. Guatemala. 249 pp.
- ILDIS (International Legume Database and Information Service). 2014. Accessed on line at <http://www.ildis.org/>.
- INAB-SEINEF (Instituto Nacional de Bosques). 2015. Sistema Electronico de Informacion de Empresas Forestales. Accessed on line: March, 2015: <http://seinef.inab.gob.gt/> .
- INAB (Instituto Nacional de Bosques). 2015. Identificación y Priorización de Recursos Genéticos Forestales para Conservación. Instituto Nacional de Bosques. Guatemala. pp19.
- INAB (Instituto Nacional de Bosques). 2012. Base de datos de plantaciones del PINFOR. Guatemala.
- INAB, CONAP, UVG & URL. 2012. Mapa De Bosques y Uso de la Tierra 2012. Mapa de cambios en Uso de la Tierra 2001-2010 para estimacion de emisiones de gases de efecto invernadero. Documento informativo. Pp. 16.
- INAB, CONAP, UVG & URL. 2012. Mapa de Cobertura Forestal de Guatemala 2010 y Dinámica de la Cobertura Forestal 2006-2010. Guatemala. pp.114.
- INAB – IARNA-URL. (Instituto Nacional de Bosques e Instituto de Agricultura, Recursos Naturales y Ambiente de la Universidad Rafael Landívar). 2012. Primer Informe Nacional sobre el Estado de los Recursos Genéticos Forestales en Guatemala. Guatemala, XX pp. 186.
- IUCN Red List of Threatened Species. Version 2015-4. <www.iucnredlist.org>. Downloaded on 27 January 2016
- Jenkins, A., Bridgland, N., Hembery, R., Malessa, U., Hewitt, J. and Keong, C.H. 2012. "Background Paper1: Precious Woods: Exploitation of the Finest Timber". TRAFFIC. Chatman House Workshop. Tackling the Trade in Illegal Precious Woods. 23-24 April 2012, <http://www.traffic.org/non-traffic-papers/>
- Khare, C.P. 2007. Indian Medicinal Plants: an illustrated Dictionary. Springer-Verlag, New York, USA. Pp. 199-201.
- Kiuru, J. 2003. Asistencia técnica en industrias forestales para asociaciones forestales de Guatemala (Informe de consultoría). Guatemala. Instituto Nacional de Bosques, Programa Regional Forestal de Centroamérica.
- Kline M. 1980. *Dalbergia stevensonii* Honduras Rosewood. In Flynn, J.H. 1994. A guide to useful woods of the world. King Philip Publishing Co: Portland, Maine, US. pp.135-136.
- Klitgaard, B.B., Lavin, M. 2005. Tribe Dalbergieae. In: Lewis, G.P., Schrire, B.D., Lock, M., Mackinder, B. 2003. (Eds.): Legumes of the World. Royal Botanic Gardens, Kew, pp.307-335.
- Knowles, D.B. and Leopold, A.C. 1997. Native tree restoration on abandoned lands in Costa Rica. Poster presentation at the Society for Ecological Restoration Annual Meeting (November 12-15 1997, Ft. Lauderdale, FL).

- Lemmens, R.H.M.J. 2008. *Dalbergia melanoxylo* Guill. & Perr. Published on the Internet in PROTA4U. Louppe, D., Oteng-Amoako, A.A. & Brink, M. (Editors). PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands;
<http://www.prota4u.info/protav8.asp?h=M4&t=Dalbergia,melanoxylo&p=Dalbergia+melanoxylo#Synonyms> (accessed 25 September 2015).
- Lewis, G.P., Schrire, B.D. 2003. Advances in Legume Systematics, Part 10, Higher Level Systematics, 1-3 (Eds: B.B. Klitgaard, A. Bruneae). Royal Botanic Gardens, Kew, England, U.K.
- Linares, J. and Sousa, M.S. 2007. Nuevas especies de *Dalbergia* (Leguminosae: Papilionoideae: Dalbergieae) en México y Centroamérica. *Ceiba*. 48(1-2): 61-82.
- Maass, M. 1995. Conversion of tropical dry forest to pasture and agriculture. In: Bullock, S.H., Mooney, H.A. y Medina, E. 1995. Seasonally Dry Tropical Forests. The University Press, Cambridge. pp 399-422.
- Mabberley, D.J. 2008. Mabberley's plant book. A portable dictionary of plants, their classifications and uses. 3 Edition, University of Washington Botanic Gardens. Seattle.
- Madrigal, Q.J. 1993. Árboles maderables en peligro de extinción en Costa Rica. INCAFO, San José, Costa Rica.
- Marín, W.A. and Flores, E.M. 2003. *Dalbergia retusa* Hemsl. In: Vozzo, J.A. 2003. Tropical Tree seeds Manual. Part II Species descriptions. United States Department of Agriculture Forest Service. pp. 429-431.
- McLure P. J., Chavarria, G.D. and Espinoza E. 2015. Metabolic chemotypes of CITES protected *Dalbergia* timbers from Africa, Madagascar and Asia. *Rapid Commun. Mass Spectrom*: 29: 1–6.
- Murphy, P.G. and Lugo, A.E. 1995. Dry forests of Central America and the Caribbean. In: Bullock, S.H., Mooney, H.A. and Medina, E. 1995. Seasonally Dry Tropical Forests. The University Press, Cambridge. pp 9-34.
- NAS/NRC. 1979. *Tropical Legumes: resources for the future*. Report by an ad hoc advisor panel of the advisory Committee on Technology Innovation, Board on science and technology for International Development, Commission on International Relations, National Academy of Sciences and the National Research Council, Washington, DC.
- Newman, D.H. 2004. (Case Study). *Evaluating the Opportunity Costs in Establishing a Nature Reserve*. In: Groom, M.J., Meffe, G.K and Carroll, R.C. (Eds) Principles of conservation biology. Third edition. Sinauer Press. pp. 529-531.
- Nghia, N.H. 1998. *Dalbergia annamensis*. The IUCN Red List of Threatened Species 1998: e.T35944A9969544. <http://dx.doi.org/10.2305/IUCN.UK.1998.RLTS.T35944A9969544.en>. Downloaded on 03 December 2015.
- Opulukwa, M.J., K. Hamza and Malende, Y. 2002. Inventory of *Dalbergia melanoxylo* (Mpingo) in the southern part of Tanzania: the case of Nachingwea. *African Study Monographs*, 23(1), 1-10.
- PROCESO. 2014. Obedecemos al pueblo, no al gobierno”, advierten comunitarios nahuas de Michoacán. <http://www.proceso.com.mx/?p=371024>.
- PROFEPA. 2014. Data on seizures of tropical wood from 2000-2014. OF. NO. PFPA/5.3/12C.6/06867.
- Rasolomampianina, R., Bailly, X., Fetierson, R., Rabehohitra, R., Béna, G, Ramaroson, L., Rahehimandimby, M., Moulin, L., de Lajudie, P., Dreyfus, B. and Avarre, J-C. 2005. Nitrogen-fixing nodules from rose wood legume trees (*Dalbergia* spp.) endemic to Madagascar host seven different genera belonging to α - and β -Proteobacteria. *Molecular Ecology* (14)13: 4135.
- Record, S.J. and Hess, R.W. 1943. Timbers of the New World. Yale University Press, New Haven; H. Milford, Oxford University Press, London. 640pp.
- Richter, H.G. (2006). Pers. Comm. (email) 27 Nov 2006 from Dr. H.G. Richter, Departamento de Madera, Celulosa y Papel, Universidad de Guadalajara, Jalisco, Mexico, concerning possible inclusion of *Cedrela odorata*, *Dalbergia retusa* and *Dalbergia stevensonii* in CITES Appendix II.
- Saha, S., Shilpi, J.A., Mondal, H., Hossain, F., Anisuzzman, Md., Mahadhi Hasan, Md. and Cordell, G.A., 2013. *Phytopharmacology*. 4(2): 291-346.

- SCMRE. Smithsonian Center for Materials Research and Education. 2002. Microscopy: Technical Information Sheet *Dalbergia retusa*.
http://www.si.edu/scmre/educationoutreach/dalbergia_retusa.htm.
- Shu, H.T. 2010. *Dalbergia* Linnaeus f., Suppl. Pl. 52, 316. 1782, nom. Cons. *Flora of China*. 10: 121-130.
- Sprent, J.I., 2009. *Legume Nodulation*. Wiley.
- Standley, P.C. and Steyermark, J.A. 1946. Flora of Guatemala. Fieldiana: Botany 24 (6). Chicago Natural History Museum.
- Stevens, W.D., Ulloa, C., Pool, A. and Montiel, O.M. 2001. *Flora de Nicaragua*. 85 (2). Missouri Botanic Garden Press.
- The Plant List. 2013. Version 1.1. Published on the Internet; <http://www.theplantlist.org/>. Accessed 01 April 2016
- UNEP-WCMC. 2015. Overview of *Dalbergia* spp. from South and Central America- a basic review. SRG 74/7/3. UNEP-WCMC, Cambridge.
- UNEP-WCMC. 2016a. CITES Trade Database. *Dalbergia* spp. global trade 2005-2014. Downloaded 5 January 2016
- UNEP-WCMC. 2016b. CITES Trade Database. *Dalbergia* spp. exports from Guatemala 2011-2014. Downloaded 01 April 2016
- UNEP-WCMC. 2016c. CITES Trade Database. *Dalbergia* spp. global trade 2000-2015. Downloaded 01 April 2016
- Vaglica, V. 2014 *Dalbergia* spp. A case for CITES listing? Thesis submitted in partial fulfilment of the requirements to obtain the Master's Degree in Management and Conservation of Species in Trade: The International Framework (11th Edition). Universidad Internacional de Andalucía, Sede Antonio Machado de Baeza
- Vasudeva, S.P. and Sareen, T.S. 2009. Pollination biology in *Dalbergia sissoo* Rpxb. (Papilionodeae; Leguminosae). *Indian Forester* 135 (9): 1165-1168.
- Vatanparast, M., Klitgard, B.B., Adema, F.A.C.B., Pennington, R.T., Yahara, T. and Kajita, T. 2013. First molecular phylogeny of the pantropical genus *Dalbergia*: implications for infragenetic circumscription and biogeography. *South African Journal of Botany*. 89: 143-149.
- Vivero, J.L., Szejner, M., Gordon, J. and Magin, G. 2006. *The Red List of Trees of Guatemala*. Global Trees Campaign, IUCN SSC. Fauna & Flora International, Cambridge, UK.
- Wiedenhoeft, A.C. 2011. Identification of Central American Woods. USDA Forest Service and Forest products Society. Pp. 90-95.
- Wiemann, M. C. and Ruffinatto, F. 2012. Separation of *Dalbergia stevensonii* from *Dalbergia tucurensis*. Research Paper FPL-RP-665. Madison, WI: U.S. Department of Agriculture, Forest Service, Forest Products Laboratory. 7 p.
- World Conservation Monitoring Centre. 1998. *Dalbergia odorifera*. The IUCN Red List of Threatened Species 1998: e.T32398A9698077.
<http://dx.doi.org/10.2305/IUCN.UK.1998.RLTS.T32398A9698077.en>. Downloaded on 03 December 2015.
- Zamora, N. 2000. Nuevas especies y combinaciones en Leguminosas de Mesoamérica. *Novon* 10: 175-180.

13. List of Annexes

Annex 1.

Annex 2

Check list of accepted names and synonyms of the genus *Dalbergia* according to “The Plant List”; accepted names are presented in **bold roman** type and synonyms are presented in *italic type*(www.theplantlist.org, accessed on line on 10 April 2016).

All Names	Accepted names
Dalbergia abbreviata Craib	
Dalbergia abrahamii Bosser & R.Rabev.	
<i>Dalbergia acaciifolia Dalzell</i>	Dalbergia pinnata var. acaciifolia (Dalzell) Thoth.
Dalbergia acariantha Harms	
Dalbergia acuta Benth.	
Dalbergia acutifoliolata Mendonca & Sousa	
Dalbergia adami Berhaut	
Dalbergia afzeliana G.Don	
<i>Dalbergia afzelii Baker</i>	Amerimnon afzelii (Baker) Kuntze
Dalbergia agudeloi J. Linares & M. Sousa	
Dalbergia ajudana Harms	
Dalbergia albertisii Prain	
Dalbergia albiflora Hutch. & Dalziel	
<i>Dalbergia albiflora subsp. albiflora</i>	Dalbergia albiflora Hutch. & Dalziel
Dalbergia altissima Baker f.	
Dalbergia amazonica (Radlk.) Ducke	
<i>Dalbergia ambongoensis Baill.</i>	Dalbergia greveana Baill.
Dalbergia amerimmon L. ex B.D. Jacks.	
<i>Dalbergia amerimnum Benth.</i>	Dalbergia brownei (Jacq.) Urb.
Dalbergia andapensis Bosser & R.Rabev.	
<i>Dalbergia angustifolia Hassk.</i>	Millettia sericea Benth.
<i>Dalbergia arborea B.Heyne</i>	Dalbergia lanceolaria L.f.
<i>Dalbergia arborea Willd.</i>	Pongamia pinnata (L.) Pierre
Dalbergia arbutifolia Baker	
Dalbergia arbutifolia subsp. aberrans Polhill	
<i>Dalbergia arbutifolia subsp. arbutifolia</i>	Dalbergia arbutifolia Baker

All Names	Accepted names
Dalbergia armata E.Mey.	
<i>Dalbergia assamica</i> "Benth., p.p."	Dalbergia sericea G.Don
Dalbergia assamica Benth.	
<i>Dalbergia atropurpurea</i> Ducke	Dalbergia foliosa (Benth.) A.M.Carvalho
<i>Dalbergia aturensis</i> Pittier	Dalbergia inundata Benth
Dalbergia aurea Bosser & R.Rabev.	
Dalbergia bakeri Baker	
Dalbergia balansae Prain	
<i>Dalbergia barclayi</i> Hook.	Mundulea barclayi (Hook.) Du Puy.
<i>Dalbergia barclayii</i> Hook. [Spelling variant]	Mundulea barclayi (Hook.) Du Puy.
Dalbergia bariensis Pierre	
Dalbergia baronii Baker	
<i>Dalbergia barretoana</i> Hoehne	Dalbergia villosa var. barretoana
Dalbergia bathiei R.Vig.	
Dalbergia beccarii Prain	
Dalbergia beddomei Thoth.	
<i>Dalbergia benthami</i> Prain [Spelling variant]	Dalbergia benthamii Prain
Dalbergia benthamii Prain	
<i>Dalbergia bernieri</i> Baill.	Dalbergia suaresensis Baill.
Dalbergia berteroi (DC.) Urb.	
<i>Dalbergia bhutanica</i> Thoth.	Dalbergia assamica Benth
Dalbergia bignonae Berhaut	
Dalbergia bintuluensis Sunarno & Ohashi	
<i>Dalbergia blumei</i> Hassk.	Dalbergia pinnata (Lour.) Prain
<i>Dalbergia boehmii</i> subsp. <i>boehmii</i>	Dalbergia boehmii Taub.
Dalbergia boehmii subsp. stuhlmannii (Taub.) Polhill	
Dalbergia boehmii Taub.	
<i>Dalbergia boinensis</i> Jum.	Dalbergia trichocarpa Baker
<i>Dalbergia boivinii</i> Baill.	Dalbergia hildebrandtii Vatke

All Names	Accepted names
Dalbergia bojeri Drake	
Dalbergia boniana Gagnep.	
Dalbergia borneensis Prain	
Dalbergia brachystachya Bosser & R.Rabev.	
Dalbergia bracteolata Baker	
Dalbergia brasiliensis Vogel	
<i>Dalbergia brevicaudata</i> Vatke	Craibia brevicaudata (Vatke) Dunn
<i>Dalbergia brownei</i> (Jacq.) "Schinz, p.p."	Dalbergia ecastaphyllum (Jacq.) Urb..
<i>Dalbergia brownei</i> (Jacq.) Schinz [Spelling variant]	Dalbergia brownie (Jacq.) Urb.
Dalbergia brownei (Jacq.) Urb.	
Dalbergia burmanica Prain	
Dalbergia calderonii Standl.	
<i>Dalbergia calderonii</i> var. <i>calderonii</i>	Dalbergia calderonii Standl.
Dalbergia calycina Benth.	
Dalbergia cambodiana Pierre	
<i>Dalbergia campechiana</i> Benth.	Dalbergia glabra (Mill.) Standl.
Dalbergia campononii Drake	
Dalbergia cana Kurz	
Dalbergia candenatensis (Dennst.) Prain	
Dalbergia canescens (Elmer) Merr.	
Dalbergia capuronii Bosser & R.Rabev.	
Dalbergia carringtoniana Sousa	
<i>Dalbergia cassioides</i> A.Gray	Dalbergia mimosella (Blanco) Prain.
Dalbergia catingicola Harms	
Dalbergia caudata G.Don	
<i>Dalbergia cavaleriei</i> H. Lévl.	Dalbergia stenophylla Prain.
Dalbergia cearensis Ducke	
<i>Dalbergia championii</i> Thwaites	Dalbergia pseudolsisoo Miq.
Dalbergia chapelieri Baill.	

All Names	Accepted names
<i>Dalbergia chermezonii</i> R.Vig.	Dalbergia mollis var. menabeensis (R.Vig.) Bosser & R.Rabev.
Dalbergia chlorocarpa R.Vig.	
Dalbergia chontalensis Standl. & L.O.Williams	
<i>Dalbergia cibix</i> Pittier	Dalbergia glabra (Mill.) Standl.
Dalbergia clarkei Thoth.	
<i>Dalbergia cloiselii</i> Drake	Dalbergia madagascariensis var. poolii (Baker) Bosser & R.Rabev.
Dalbergia cochinchinensis Laness.	
Dalbergia cochinchinensis Pierre	
<i>Dalbergia colettii</i> "Prain, p.p."	Dalbergia yunnanensis var. colettii (Prain) Thoth.
<i>Dalbergia colettii</i> Prain	Dalbergia yunnanensis Franch.
Dalbergia commiphoroides Baker f.	
Dalbergia comorensis Bosser & R. Rabev.	
<i>Dalbergia confertiflora</i> "Benth., p.p."	Dalbergia volubilis Roxb.
Dalbergia confertiflora Benth.	
<i>Dalbergia confertiflora</i> var. <i>confertiflora</i>	Dalbergia cuiabensis Benth
Dalbergia congensis Baker f.	
Dalbergia congesta Wight & Arn.	
Dalbergia congestiflora Pittier	
Dalbergia coromandeliana Prain	
Dalbergia crispa Hepper	
Dalbergia cubilquitzensis (Donn.Sm.) Pittier	
Dalbergia cucullata Pittier	
Dalbergia cuiabensis Benth.	
<i>Dalbergia cujabensis</i> Benth. [Spelling variant]	Dalbergia cuiabensis Benth
Dalbergia cultrata Benth.	
<i>Dalbergia cultrata</i> var. <i>cultrata</i>	Dalbergia cultrata Benth
Dalbergia cumingiana Benth.	
<i>Dalbergia cumingii</i> Benth.	Dalbergia parviflora Roxb.
Dalbergia curtisii Prain	

All Names	Accepted names
Dalbergia cuscatlanica (Standl.) Standl.	
Dalbergia dalzielii Hutch. & Dalziel	
Dalbergia darienensis Rudd	
<i>Dalbergia davaoensis Elmer</i>	Dalbergia mimosella (Blanco) Prain
Dalbergia davidii Bosser & R.Rabev.	
Dalbergia debilis J.F.Macbr.	
Dalbergia decipularis Rizzini & A.Mattos	
<i>Dalbergia delavayi Franch.</i>	Cladrastis delavayi (Franch.) Prain
Dalbergia delphinensis Bosser & R.Rabev.	
Dalbergia densa Benth.	
<i>Dalbergia densa var. densa</i>	Dalbergia densa Benth
<i>Dalbergia densicoma Baill.</i>	Dalbergia pervillei Vatke.
Dalbergia densiflora (Benth.) Benth.	
Dalbergia dialoides (Pierre) Niyomdham	
<i>Dalbergia diphaca Pers.</i>	Ormocarpum cochinchinense (Lour.) Merr.
Dalbergia discolor Blume	
<i>Dalbergia discolor Blume ex Miq.</i>	Amerimnon discolor (Blume ex Miq.) Kuntze.
<i>Dalbergia diversifolia Miq.</i>	Dalbergia pseudolsissoo Miq.
<i>Dalbergia domingensis Pers.</i>	Lonchocarpus domingensis (Pers.) DC.
Dalbergia dongnaiensis Pierre	
Dalbergia duarensis Thoth.	
<i>Dalbergia dubia Elmer</i>	Dalbergia pinnata (Lour.) Prain
Dalbergia duperreana Pierre	
Dalbergia dyeriana Harms	
Dalbergia ealaensis De Wild.	
<i>Dalbergia ecastaphylla (L.) Taub. [Spelling variant]</i>	Dalbergia ecastaphyllum (Jacq.) Urb.
Dalbergia ecastaphyllum (L.) Taub.	
<i>Dalbergia ecastophyllum (L.) Taub. [Spelling variant]</i>	Dalbergia ecastaphyllum (Jacq.) Urb.
<i>Dalbergia elata Harms</i>	Dalbergia boehmii Taub.
Dalbergia elegans A.M.Carvalho	

All Names	Accepted names
<i>Dalbergia emarginata</i> Prain	Dalbergia sissoides Wight & Arn.
<i>Dalbergia emarginata</i> Roxb.	Dalbergia latifolia Roxb.
Dalbergia emirnensis Benth.	
<i>Dalbergia emirnensis</i> var. <i>emirnensis</i>	Dalbergia emirnensis Benth
<i>Dalbergia enneandra</i> Hoehne	Dalbergia riedelii (Benth.) Sandwith.
Dalbergia enneaphylla Pittier	
Dalbergia entadoides Prain	
Dalbergia eremicola Polhill	
Dalbergia ernest-ulei Hoehne	
Dalbergia errans Craib	
Dalbergia erubescens Bosser & R.Rabev.	
<i>Dalbergia eurybothrya</i> Drake	Dalbergia greveana Baill.
Dalbergia falcata Prain	
<i>Dalbergia ferruginea</i> Glaz.	Dalbergia glaziovii Harms.
<i>Dalbergia ferruginea</i> Roxb.	Dalbergia stipulacea Roxb.
<i>Dalbergia ferruginea</i> var. <i>daronensis</i> Elmer	Dalbergia stipulacea Roxb.
<i>Dalbergia ferrugineo-tomentosa</i> Hoehne	Dalbergia cuiabensis Benth
Dalbergia fischeri Taub.	
<i>Dalbergia flexuosa</i> Benth.	Dalbergia reniformis Roxb.
Dalbergia floribunda Craib	
Dalbergia florifera De Wild.	
<i>Dalbergia foliacea</i> "Prain, p.p."	Dalbergia entadoides Prain.
<i>Dalbergia foliacea</i> Benth.	Dalbergia rimosa var. <i>foliacea</i> (Benth.) Thoth.
Dalbergia foliolosa Benth.	
Dalbergia foliosa (Benth.) A.M.Carvalho	
<i>Dalbergia foliosa</i> (Benth.) T.S.Ralph	Dalbergia foliosa (Benth.) A.M.Carvalho
Dalbergia forbesii Prain	
Dalbergia fouilloyana Pellegr.	
<i>Dalbergia frondosa</i> DC.	Dalbergia lanceolaria L.f.
Dalbergia frutescens (Vell.) Britton	

All Names	Accepted names
<i>Dalbergia frutescens</i> var. <i>frutescens</i>	Dalbergia frutescens (Vell.) Britton
Dalbergia frutescens var. tomentosa (Vogel) Benth.	
Dalbergia funera Standl.	
Dalbergia fusca Pierre	
<i>Dalbergia fusca</i> Prain	Dalbergia cultrata Benth
<i>Dalbergia fusca</i> var. <i>enneandra</i> S.Q.Zou & J.H.Liu	Dalbergia fusca Pierre
Dalbergia gardneriana Benth.	
Dalbergia gentilii De Wild.	
<i>Dalbergia gerardiana</i> Benth. [Spelling variant]	Dalbergia gardneriana Benth
Dalbergia gilbertii Cronquist	
<i>Dalbergia gilletii</i> De Wild.	Dalbergia hostilis Benth
Dalbergia glaberrima Bosser & R.Rabev.	
<i>Dalbergia glaberrima</i> subsp. <i>glaberrima</i>	Dalbergia glaberrima Bosser & R.Rabev.
Dalbergia glabra (Mill.) Standl.	
<i>Dalbergia glabra</i> var. <i>chontalensis</i> (Standl. & L.O.Williams) Rudd	Dalbergia chontalensis Standl. & L.O.Williams.
<i>Dalbergia glabra</i> var. <i>glabra</i>	Dalbergia glabra (Mill.) Standl.
Dalbergia glandulosa Benth.	
<i>Dalbergia glauca</i> (Desv.) Amshoff	Dalbergia foliosa (Benth.) A.M.Carvalho
<i>Dalbergia glauca</i> Kurz	Dalbergia obtusifolia (Baker) Prain
<i>Dalbergia glauca</i> Wall.	Dalbergia ovata Benth
Dalbergia glaucescens (Benth.) Benth.	
Dalbergia glaucocarpa Bosser & R.Rabev.	
Dalbergia glaziovii Harms	
Dalbergia glomerata Hemsl.	
<i>Dalbergia glomeriflora</i> Kurz	Dalbergia ovata var. glomeriflora (Kurz) Thoth.
Dalbergia godefroyi Prain	
<i>Dalbergia goetzei</i> Harms	Dalbergia bracteolata Baker
Dalbergia gossweileri Baker f.	
Dalbergia gracilis Benth.	

All Names	Accepted names
Dalbergia granadillo Pittier	
Dalbergia grandibracteata De Wild.	
<i>Dalbergia grandidieri Baill.</i>	Dalbergia bracteolata Baker
Dalbergia grandistipula A.M.Carvalho	
Dalbergia greveana Baill.	
<i>Dalbergia gummifera DC.</i>	Ateleia gummifera (DC.) D.Dietr.
Dalbergia guttembergii A.M.Carvalho	
Dalbergia hainanensis Merr. & Chun	
Dalbergia hancei Benth.	
<i>Dalbergia harmsiana De Wild.</i>	Dalbergia boehmii Taub.
<i>Dalbergia havilandi Prain [Spelling variant]</i>	Dalbergia havilandii Prain
Dalbergia havilandii Prain	
<i>Dalbergia hemsleyi Prain</i>	Dalbergia lanceolaria subsp. paniculata (Roxb.) Thoth.
Dalbergia henryana Prain	
<i>Dalbergia heptaphylla Poir.</i>	Lonchocarpus heptaphyllus (Poir.) DC.
<i>Dalbergia heterophylla Willd.</i>	Derris trifoliata Lour.
Dalbergia heudelotii Stapf	
Dalbergia hiemalis Malme	
<i>Dalbergia hildebrandtii var. hildebrandtii</i>	Dalbergia hildebrandtii Vatke
Dalbergia hildebrandtii var. scorpioides (Baker) Bosser & R.Rabev.	
Dalbergia hildebrandtii Vatke	
<i>Dalbergia hircina Benth.</i>	Dalbergia sericea G.Don.
<i>Dalbergia hircina Wall.</i>	Dalbergia sericea G.Don.
Dalbergia hirticalyx Bosser & R.Rabev.	
<i>Dalbergia hookeri Bojer</i>	Mundulea sericea subsp. madagascariensis Du Puy & Labat
Dalbergia horrida (Dennst.) Mabb.	
<i>Dalbergia horrida Graham</i>	Dalbergia spinosa Roxb.
Dalbergia horrida var. glabrescens (Prain) Thoth. & Nair	

All Names	Accepted names
<i>Dalbergia horrida</i> var. <i>horrida</i>	Dalbergia horrida (Dennst.) Mabb.
Dalbergia hortensis Heringer & al.	
Dalbergia hoseana Prain	
Dalbergia hostilis Benth.	
Dalbergia hullettii Prain	
Dalbergia humbertii R.Vig.	
Dalbergia hupeana Hance	
<i>Dalbergia hupeana</i> var. <i>hupeana</i>	Dalbergia hupeana Hance
Dalbergia hygrophila (Benth.) Hoehne	
<i>Dalbergia hygrophila</i> var. <i>nephrocarpa</i> (Ducke) Hoehne	Dalbergia amazonica (Radlk.) Ducke
<i>Dalbergia hypargyrea</i> Harms	Machaerium floribundum var. hypargyream (Harms) Rudd.
<i>Dalbergia hypoleuca</i> Pittier	Dalbergia retusa Hemsl.
<i>Dalbergia ikopensis</i> Jum.	Dalbergia greveana Baill.
<i>Dalbergia inaequilatera</i> (Poepp.) Wiehler	Columnnea inaequilatera Poepp. & Endl.
<i>Dalbergia insularis</i> Pulle	Dalbergia beccarii Prain.
Dalbergia intermedia A.M.Carvalho	
Dalbergia intibucana Standl. & L.O.Williams	
Dalbergia inundata Benth.	
Dalbergia iquitosensis Harms	
<i>Dalbergia isaloensis</i> R.Vig.	Dalbergia greveana Baill.
<i>Dalbergia isangiensis</i> De Wild.	Dalbergia saxatilis Hook.f.
Dalbergia jaherii Burck	
<i>Dalbergia javanica</i> Miq.	Dalbergia sissoides Wight & Arn.
Dalbergia jingxiensis S.Y. Liu	
<i>Dalbergia junghuhnii</i> "Baker, p.p."	Dalbergia curtisii Prain
Dalbergia junghuhnii Benth.	
Dalbergia kerrii Craib	
Dalbergia kingiana Prain	
Dalbergia kisantuensis De Wild. & T.Durand	

All Names	Accepted names
Dalbergia kostermansii Sunarno & Ohashi	
<i>Dalbergia krowee Roxb.</i>	Derris robusta (DC.) Benth
Dalbergia kunstleri Prain	
Dalbergia kurzii Prain	
<i>Dalbergia kurzii var. kurzii</i>	Dalbergia kurzii Prain
<i>Dalbergia laccifera Laness.</i>	Dalbergia oliveri Prain
Dalbergia lacei Thoth.	
Dalbergia lactea Vatke	
<i>Dalbergia laevigata Standl.</i>	Acosmium panamense (Benth.) Yakovlev
Dalbergia lakhonensis Gagnep.	
<i>Dalbergia lakhonensis var. lakhonensis</i>	Dalbergia lakhonensis Gagnep.
<i>Dalbergia lanata (Seem.) Wiehler</i>	Columnea lanata (Seem.) Kuntze
<i>Dalbergia lanceolaria Hemsl.</i>	Dalbergia balansae Prain
Dalbergia lanceolaria L.f.	
<i>Dalbergia lanceolaria Litv.</i>	Dalbergia mimosella (Blanco) Prain
<i>Dalbergia lanceolaria Moon</i>	Pericopsis mooniana Thwaites
<i>Dalbergia lanceolaria Nav. & Ferdinand</i>	Dalbergia mimosella (Blanco) Prain
<i>Dalbergia lanceolaria subsp. lanceolaria</i>	Dalbergia lanceolaria L.f.
Dalbergia lanceolaria subsp. paniculata (Roxb.) Thoth.	
<i>Dalbergia lanceolaria var. assamica (Benth.) Thoth.</i>	Dalbergia assamica Benth
Dalbergia lanceolaria var. maymyensis (W. G. Craib) Thoth.	
Dalbergia lastoursvillensis Pellegr.	
Dalbergia lateriflora Benth.	
Dalbergia latifolia Roxb.	
<i>Dalbergia latifolia var. sissoides (Wight & Arn.) Baker</i>	Dalbergia sissoides Wight & Arn.
<i>Dalbergia laurentii De Wild.</i>	Aganope lucida (Baker) Polhill
<i>Dalbergia lavigata Standl. [Spelling variant]</i>	Acosmium panamense (Benth.) Yakovlev
Dalbergia laxiflora Micheli	
Dalbergia lemurica Bosser & R.Rabev.	

All Names	Accepted names
Dalbergia librevillensis Pellegr.	
<i>Dalbergia limonensis Benth.</i>	Dalbergia stipulacea Roxb.
<i>Dalbergia lindeniana (Benth.) Hoehne</i>	Machaerium lindenianum Benth
<i>Dalbergia lineata Pittier</i>	Dalbergia retusa Hemsl.
<i>Dalbergia livida "Wall., p.p."</i>	Dalbergia pinnata (Lour.) Prain
Dalbergia longepedunculata J. Linares & M. Sousa	
Dalbergia louisii Cronquist	
Dalbergia louvelii R.Vig.	
Dalbergia luteola J. Linares & M. Sousa	
<i>Dalbergia luzonensis Vogel</i>	Dalbergia stipulacea Roxb.
<i>Dalbergia macrophylla (Kunth) Spreng.</i>	Lonchocarpus macrophyllus Kunth
Dalbergia macrosperma Baker	
<i>Dalbergia macrothyrsa Harms</i>	Dalbergia saxatilis Hook.f.
<i>Dalbergia </i> <i>madagascariensis</i> <i>subsp.</i> <i>madagascariensis</i>	Dalbergia madagascariensis Vatke
Dalbergia madagascariensis var. poolii (Baker) Bosser & R.Rabev.	
Dalbergia madagascariensis Vatke	
Dalbergia malabarica Prain	
Dalbergia malangensis Sousa	
<i>Dalbergia malifolia Welw. ex Baker</i>	Amerimnon malifolium (Welw. ex Baker) Kuntze
Dalbergia mammosa Pierre	
Dalbergia marcaniana Craib	
<i>Dalbergia marginata Roxb.</i>	Derris marginata (Roxb.) Benth
Dalbergia maritima R.Vig.	
<i>Dalbergia maritima var. maritima</i>	Dalbergia maritima R.Vig.
Dalbergia martinii F.White	
<i>Dalbergia maymensis Craib</i>	Dalbergia lanceolaria subsp. paniculata (Roxb.) Thoth.
<i>Dalbergia maymyensis W. G. Craib</i>	Dalbergia lanceolaria var. maymyensis (W. G. Craib) Thoth.
Dalbergia mayumbensis Baker f.	

All Names	Accepted names
<i>Dalbergia megalocarpa</i> Harms	Dalbergia arbutifolia subsp. aberrans Polhill.
Dalbergia melanocardium Pittier	
Dalbergia melanoxydon Guill. & Perr.	
<i>Dalbergia mengsuoensis</i> Y.Y. Qian	Dalbergia rimosa Roxb.
Dalbergia menoeides Prain	
Dalbergia mexicana Pittier	
<i>Dalbergia mexicana</i> var. <i>hygrophila</i> (Mart. ex Benth.) J.F. Macbr.	Dalbergia hygrophila (Benth.) Hoehne.
<i>Dalbergia microcarpa</i> Baker f.	Dalbergia microphylla Chiov.
<i>Dalbergia microcarpa</i> R. Vig.	Dalbergia peltieri Bosser & R.Rabev.
Dalbergia microphylla Chiov.	
Dalbergia millettii Benth.	
<i>Dalbergia millettii</i> Prain	Dalbergia millettii var. <i>mimosoides</i> (Franch.) Thoth.
Dalbergia millettii var. <i>mimosoides</i> (Franch.) Thoth.	
Dalbergia mimosella (Blanco) Prain	
Dalbergia mimosoides Franch.	
<i>Dalbergia minahassae</i> Koord.	Dalbergia mimosella (Blanco) Prain
Dalbergia miscolobium Benth.	
Dalbergia modesta J. Linares & M. Sousa	
Dalbergia mollis Bosser & R.Rabev.	
Dalbergia mollis var. <i>menabeensis</i> (R.Vig.) Bosser & R.Rabev.	
<i>Dalbergia mollis</i> var. <i>mollis</i>	Dalbergia mollis Bosser & R.Rabev.
Dalbergia monetaria L.f.	
<i>Dalbergia monetaria</i> var. <i>brachyphylla</i> Radlk.	Dalbergia monetaria L.f.
<i>Dalbergia monetaria</i> var. <i>hygrophila</i> (Benth.) J.F. Macbr.	Dalbergia hygrophila (Benth.) Hoehne.
<i>Dalbergia monetaria</i> var. <i>monetaria</i>	Dalbergia monetaria L.f.
Dalbergia monophylla G.A. Black	
<i>Dalbergia monosperma</i> Dalzell	Dalbergia candenatensis (Dennst.) Prain
Dalbergia monticola Bosser & R.Rabev.	

All Names	Accepted names
<i>Dalbergia mooniana</i> Thwaites	Pericopsis mooniana Thwaites.
<i>Dalbergia multiflora</i> B.Heyne	Dalbergia horrida (Dennst.) Mabb.
<i>Dalbergia multiflora</i> Prain	Dalbergia horrida (Dennst.) Mabb.
<i>Dalbergia multiflora</i> var. <i>glabrescens</i> Prain	Dalbergia horrida var. <i>glabrescens</i> (Prain) Thoth. & Nair
Dalbergia multijuga E.Mey.	
<i>Dalbergia myriabotrys</i> Baker	Dalbergia greveana Baill.
Dalbergia negrensis (Radlk.) Ducke	
<i>Dalbergia nelsii</i> Schinz	Philenoptera nelsii (Schinz) Schrire
Dalbergia neoperrieri Bosser & R.Rabev.	
<i>Dalbergia nephrocarpa</i> Ducke	Dalbergia amazonica (Radlk.) Ducke.
Dalbergia ngounyensis Pellegr.	
Dalbergia nigra (Vell.) Benth.	
<i>Dalbergia nigrescens</i> Kurz	Dalbergia lanceolaria subsp. <i>paniculata</i> (Roxb.) Thoth.
Dalbergia nitida (Benth.) Hoehne	
Dalbergia nitidula Baker	
Dalbergia noldeae Harms	
Dalbergia normandii Bosser & R.Rabev.	
<i>Dalbergia novo-guineense</i> Merr. & L.M.Perry	Dalbergia beccarii Prain
Dalbergia oblongifolia G.Don	
Dalbergia obovata E.Mey.	
<i>Dalbergia obtusa</i> Lecomte	Dalbergia pervillei Vatke
Dalbergia obtusifolia (Baker) Prain	
<i>Dalbergia obtusifolia</i> var. <i>obtusifolia</i>	Dalbergia obtusifolia (Baker) Prain
<i>Dalbergia ochracea</i> Harms	Dalbergia arbutifolia Baker
Dalbergia odorifera T.C.Chen	
Dalbergia oligophylla Hutch. & Dalziel	
Dalbergia oliveri Prain	
<i>Dalbergia oojeinensis</i> Roxb.	Desmodium oojeinense (Roxb.) H.Ohashi
Dalbergia orientalis Bosser & R.Rabev.	

All Names	Accepted names
Dalbergia ovata Benth.	
<i>Dalbergia ovata</i> Graham	Dalbergia ovata Benth
Dalbergia ovata var. glomeriflora (Kurz) Thoth.	
<i>Dalbergia ovata</i> var. <i>obtusifolia</i> Baker	Dalbergia obtusifolia (Baker) Prain
<i>Dalbergia ovata</i> var. <i>ovata</i>	Dalbergia ovata Benth
<i>Dalbergia oxyphylla</i> Harms	Dalbergia cuiabensis Benth
Dalbergia pachycarpa (De Wild. & T.Durand) De Wild.	
<i>Dalbergia pachycarpa</i> Ducke	Dalbergia riedelii (Benth.) Sandwith
<i>Dalbergia pacifica</i> Standl. & Steyerm.	Dalbergia cuscatlanica (Standl.) Standl.
Dalbergia palo-escrito Rzed.	
<i>Dalbergia paniculata</i> Roxb.	Dalbergia lanceolaria subsp. paniculata (Roxb.) Thoth.
<i>Dalbergia papuana</i> Pulle	Dalbergia albertisii Prain
Dalbergia parviflora Roxb.	
Dalbergia paucifoliolata Lundell	
Dalbergia peguensis Thoth.	
Dalbergia peishaensis Chun & T.Chen	
Dalbergia peltieri Bosser & R.Rabev.	
<i>Dalbergia pentaphylla</i> Poir.	Lonchocarpus heptaphyllus (Poir.) DC.
<i>Dalbergia perrieri</i> Drake	Dalbergia trichocarpa Baker
<i>Dalbergia perrieri</i> Jum.	Dalbergia greveana Baill.
Dalbergia pervillei Vatke	
<i>Dalbergia phyllanthoides</i> Miq.	Dalbergia junghuhnii Benth
Dalbergia pierreana Prain	
<i>Dalbergia pinatubensis</i> Elmer	Dalbergia pinnata (Lour.) Prain
Dalbergia pinnata (Lour.) Prain	
Dalbergia pinnata var. acaciifolia (Dalzell) Thoth.	
<i>Dalbergia pinnata</i> var. <i>badia</i> Merr.	Dalbergia pinnata (Lour.) Prain
<i>Dalbergia pinnata</i> var. <i>pinnata</i>	Dalbergia pinnata (Lour.) Prain
Dalbergia pluriflora Baker f.	

All Names	Accepted names
Dalbergia polyadelpha Prain	
Dalbergia polyphylla Benth.	
<i>Dalbergia polyphylla Poir.</i>	Glottidium vesicarium (Jacq.) R.M.Harper
<i>Dalbergia poolii Baker</i>	Dalbergia madagascariensis var. poolii (Baker) Bosser & R.Rabev.
<i>Dalbergia praecox Hochst. ex A. Rich.</i>	Pterocarpus lucens Guill. & Perr.
Dalbergia prainii Thoth.	
<i>Dalbergia prazeri Prain</i>	Dalbergia oliveri Prain
<i>Dalbergia preussii Harms</i>	Dalbergia saxatilis var. preussii (Harms) Cronquist
Dalbergia pseudo-ovata Thoth.	
Dalbergia pseudo-sissoo Miq.	
Dalbergia pseudobaronii R.Vig.	
<i>Dalbergia pterocarpiflora Baill.</i>	Dalbergia chapelieri Baill.
<i>Dalbergia pterocarpiflora Baker</i>	Dalbergia chapelieri Baill.
<i>Dalbergia pubescens Hook.</i>	Dalbergia subcymosa Ducke
Dalbergia purpurascens Baill.	
<i>Dalbergia purpurea "Benth., p.p."</i>	Dalbergia volubilis Roxb.
<i>Dalbergia purpurea Kurz</i>	Dalbergia kurzii Prain
<i>Dalbergia purpusii Brandegees</i>	Dalbergia glabra (Mill.) Standl.
<i>Dalbergia radiata Prain</i>	Dalbergia pseudo-sissoo Miq.
Dalbergia reniformis Roxb.	
Dalbergia reticulata Merr.	
<i>Dalbergia retusa Baill.</i>	Dalbergia pervillei Vatke
Dalbergia retusa Hemsl.	
<i>Dalbergia retusa var. lineata (Pittier) Rudd</i>	Dalbergia retusa Hemsl.
<i>Dalbergia retusa var. retusa</i>	Dalbergia retusa Hemsl.
Dalbergia revoluta Ducke	
Dalbergia rhachiflexa J. Linares & M. Sousa	
<i>Dalbergia richardii Baill.</i>	Dalbergia bracteolata Baker
Dalbergia richardsii Sunarno & Ohashi	

All Names	Accepted names
Dalbergia riedelii (Benth.) Sandwith	
Dalbergia rimosa Roxb.	
Dalbergia rimosa var. foliacea (Benth.) Thoth.	
<i>Dalbergia rimosa var. rimosa</i>	Dalbergia rimosa Roxb.
Dalbergia riparia (Mart.) Benth.	
<i>Dalbergia rivularis Merr. & L.M.Perry</i>	Dalbergia stipulacea Roxb.
<i>Dalbergia robusta DC.</i>	Derris robusta (DC.) Benth
<i>Dalbergia rosei Hoehne</i>	Dalbergia lateriflora Benth
Dalbergia rostrata Hassk.	
<i>Dalbergia rostrata Prain</i>	Dalbergia pseudo-sissoo Miq.
<i>Dalbergia rotundifolia Sond.</i>	Pterocarpus rotundifolius (Sond.) Druce
Dalbergia rubiginosa Roxb.	
Dalbergia ruddiae J. Linares & M. Sousa	
Dalbergia rufa G.Don	
<i>Dalbergia rufa Graham</i>	Dalbergia pinnata (Lour.) Prain
Dalbergia rugosa Hepper	
Dalbergia sacerdotum Prain	
<i>Dalbergia saigonensis Pierre</i>	Dalbergia nigrescens var. saigonensis (Pierre) Gagnep.
Dalbergia salvanaturae J. Linares & M. Sousa	
Dalbergia sambesiaca Schinz	
Dalbergia sampaioana Kuhl. & Hoehne	
Dalbergia sandakanensis Sunarno & Ohashi	
Dalbergia saxatilis Hook.f.	
Dalbergia saxatilis var. preussii (Harms) Cronquist	
<i>Dalbergia scandens Roxb.</i>	Derris scandens (Roxb.) Benth
<i>Dalbergia scorpioides Baker</i>	Dalbergia hildebrandtii var. scorpioides (Baker) Bosser & R.Rabev.
Dalbergia scortechinii (Prain) Prain	
<i>Dalbergia sericea Bojer</i>	Mundulea sericea (Willd.) A.Chev.

All Names	Accepted names
Dalbergia sericea G.Don	
<i>Dalbergia sessiliflora Harms</i>	Dalbergia obovata E.Mey.
Dalbergia setifera Hutch. & Dalziel	
<i>Dalbergia simplicifolia Merr.</i>	Dalbergia rostrata Hassk.
Dalbergia simpsonii Rudd	
<i>Dalbergia sissoides Graham</i>	Dalbergia sissoides Wight & Arn.
Dalbergia sissoides Wight & Arn.	
Dalbergia sissoo DC.	
Dalbergia spinosa Roxb.	
<i>Dalbergia spinosa Wight & Arn.</i>	Dalbergia coromandeliana Prain
Dalbergia spruceana (Benth.) Benth.	
<i>Dalbergia stenocarpa Kurz</i>	Dalbergia sericea G.Don
<i>Dalbergia stenocarpa var. menabeensis R. Vig.</i>	Dalbergia mollis var. menabeensis (R.Vig.) Bosser & R.Rabev.
<i>Dalbergia stenocarpa var. typica R. Vig.</i>	Dalbergia mollis Bosser & R.Rabev.
Dalbergia stenophylla Prain	
Dalbergia stercoracea Prain	
Dalbergia stevensonii Standl.	
Dalbergia stipulacea Roxb.	
<i>Dalbergia stipulacea var. stipulacea</i>	Dalbergia stipulacea Roxb.
<i>Dalbergia stipulaceae Fern.-Vill.</i>	Dalbergia stipulacea Roxb.
<i>Dalbergia stipulata Wall.</i>	Amerimnon stipulatum (Wall.) Kuntze
<i>Dalbergia stocksii Benth.</i>	Dalbergia melanoxydon Guill. & Perr.
<i>Dalbergia striata Bojer</i>	Mundulea sericea subsp. madagascariensis
<i>Dalbergia stuhlmannii Taub.</i>	Dalbergia boehmii subsp. stuhlmannii
Dalbergia suaresensis Baill.	
<i>Dalbergia subalternifolia (Elmer) Merr.</i>	Dalbergia rostrata Hassk.
Dalbergia subcymosa Ducke	
Dalbergia succirubra Gagnep. & Craib	
<i>Dalbergia succirubra Gagnepain & Craib</i>	Dalbergia velutina var. succirubra (Gagnepain & Craib) Niyomdham

All Names	Accepted names
<i>Dalbergia sympathetica</i> Nimmo	Dalbergia horrida (Dennst.) Mabb.
<i>Dalbergia szemaoensis</i> Prain	Dalbergia assamica Benth
<i>Dalbergia tabascanana</i> Pittier	Dalbergia glabra (Mill.) Standl.
<i>Dalbergia tamarindifolia</i> Roxb.	Dalbergia pinnata (Lour.) Prain
<i>Dalbergia tamarindifolia</i> var. <i>acaciifolia</i> (Dalzell) Baker	Dalbergia pinnata var. acaciifolia (Dalzell) Thoth.
<i>Dalbergia tamarindifolia</i> var. <i>pubescens</i> Baker	Dalbergia malabarica Prain
Dalbergia teijsmannii Sunarno & Ohashi	
Dalbergia teixeirae Sousa	
<i>Dalbergia telfairii</i> Bojer	Mundulea sericea subsp. madagascariensis Du Puy & Labat
Dalbergia thomsonii Benth.	
Dalbergia thorelii Gagnep.	
Dalbergia tilarana N.Zamora	
<i>Dalbergia timoriensis</i> DC.	Derris scandens (Roxb.) Benth
<i>Dalbergia tingens</i> Baill.	Rourea orientalis Baill.
Dalbergia tinneveliense Thoth.	
<i>Dalbergia tomentosa</i> (Benth.) Ducke	Dalbergia intermedia A.M.Carvalho
<i>Dalbergia tomentosa</i> Vogel	Dalbergia frutescens var. tomentosa (Vogel) Benth.
Dalbergia tonkinensis Prain	
<i>Dalbergia torta</i> A.Gray	Dalbergia candenatensis (Dennst.) Prain
<i>Dalbergia torta</i> Graham	Dalbergia candenatensis (Dennst.) Prain
<i>Dalbergia toxicaria</i> Baill.	Rourea orientalis Baill.
Dalbergia travancorica Thoth.	
Dalbergia trichocarpa Baker	
Dalbergia tricolor Drake	
<i>Dalbergia tricolor</i> var. <i>tricolor</i>	Dalbergia tricolor Drake
Dalbergia tsaratananensis Bosser & R.Rabev.	
Dalbergia tsiandalana R.Vig.	
Dalbergia tsoi Merr. & Chun	
Dalbergia tucurensis Donn.Sm.	

All Names	Accepted names
Dalbergia uarandensis (Chiov.) Thulin	
Dalbergia urschii Bosser & R.Rabev.	
Dalbergia vacciniifolia Vatke	
<i>Dalbergia variabilis</i> var. <i>bahiensis</i> Hoehne	Dalbergia cearensis Ducke
<i>Dalbergia variabilis</i> var. <i>cubilquitzensis</i> Donn.Sm.	Dalbergia cubilquitzensis (Donn.Sm.) Pittier
<i>Dalbergia variabilis</i> var. <i>tomentosa</i> (Vogel) Benth.	Dalbergia frutescens var. <i>tomentosa</i> (Vogel) Benth
<i>Dalbergia variabilis</i> Vogel	Dalbergia frutescens (Vell.) Britton
Dalbergia velutina Benth.	
Dalbergia velutina var. <i>succirubra</i> (Gagnepain & Craib) Niyomdham	
<i>Dalbergia velutina</i> var. <i>velutina</i>	Dalbergia velutina Benth
Dalbergia verrucosa Craib	
Dalbergia viguieri Bosser & R.Rabev.	
Dalbergia villosa (Benth.) Benth.	
Dalbergia villosa var. <i>barretoana</i> (Hoehne) A.M.Carvalho	
<i>Dalbergia villosa</i> var. <i>divaricata</i> Hoehne	Dalbergia villosa (Benth.) Benth
<i>Dalbergia villosa</i> var. <i>villosa</i>	Dalbergia villosa (Benth.) Benth
<i>Dalbergia violacea</i> (Jacq.) Hoffmanns.	Dalbergia miscolobium Benth
<i>Dalbergia violacea</i> (Vogel) Malme	Dalbergia miscolobium Benth
<i>Dalbergia volubilis</i> (L.) Urb.	Dalbergia monetaria L.f.
Dalbergia volubilis Roxb.	
<i>Dalbergia volubilis</i> var. <i>cuspidigera</i> Hoehne	Dalbergia monetaria L.f.
<i>Dalbergia volubilis</i> var. <i>volubilis</i>	Dalbergia volubilis Roxb.
Dalbergia wattii C.B.Clarke	
Dalbergia xerophila Bosser & R.Rabev.	
Dalbergia ximengensis Y.Y. Qian	
Dalbergia yunnanensis Franch.	
Dalbergia yunnanensis var. <i>collettii</i> (Prain) Thoth.	
<i>Dalbergia yunnanensis</i> var. <i>yunnanensis</i>	Dalbergia yunnanensis Franch.

Distribution of accepted species names of the genus *Dalbergia* according to “The International Legume Database and Information Service” (ILDIS, 2013).

Note. Different reference sources were also utilised and those are indicated by specific numbers after the ISO codes of countries mentioned in Annex 2.

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergia abbreviata</i> Craib	TH
<i>Dalbergia abrahamii</i> Bosser & R.Rabev.	MG
<i>Dalbergia acariiantha</i> Harms	TZ
<i>Dalbergia acuta</i> Benth.	BR
<i>Dalbergia acutifoliolata</i> Mendonca & Sousa	ZM
<i>Dalbergia adamii</i> Berhaut	SN
<i>Dalbergia afzeliana</i> G. Don	AO, CM, CF, CD, GA, GN, GW, NG, SL, TG
<i>Dalbergia agudeloii</i> J. Linares & M. Sousa	SV ¹ , GT ¹ , HN ¹ , NI ¹
<i>Dalbergia ajudana</i> Harms	BJ
<i>Dalbergia albertsii</i> Prain	ID, PG, SB
<i>Dalbergia albiflora</i> Hutch. & Dalziel	CI, NG, SL
<i>Dalbergia altissima</i> Baker f.	VE ¹
<i>Dalbergia amazonica</i> (Radlk.) Ducke	BR, CO ⁴ , FR ¹ , GY, VE
<i>Dalbergia amerimmon</i> L. ex B.D. Jacks.	US ¹
<i>Dalbergia andapensis</i> Bosser & R.Rabev.	MG
<i>Dalbergia arbutifolia</i> Baker	MW, MZ, TZ, ZM, ZW
<i>Dalbergia arbutifolia</i> subsp. <i>aberrans</i> Polhill	CD, TZA
<i>Dalbergia armata</i> E. Mey.	MZ, ZA, LK, SZ, TZ
<i>Dalbergia assamica</i> Benth.	BD ² , BT ² , KH ² , CN, IN, KE, LA ¹ , MM, TH ¹ , VN ¹
<i>Dalbergia aurea</i> Bosser & R.Rabev.	MG
<i>Dalbergia bakeri</i> Baker	AO, CD, GA
<i>Dalbergia balansae</i> Prain	CN, VN
<i>Dalbergia bariensis</i> Pierre	KH ² , LA ² , TH, VN

¹ www.tropicos.org

² www.iucnredlist.org

³ Saha *et al.*, 2013

⁴ Grandter & Chevrette, 2013

⁵ Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiabaronii</i> Baker	MG, TZ
<i>Dalbergiabathie</i> R. Vig.	MG
<i>Dalbergiabeccarii</i> Prain	BN, ID, MY, PG, SB
<i>Dalbergiabeddomei</i> Thoth.	IN
<i>Dalbergiabentharii</i> Prain	CN, VN ¹
<i>Dalbergiaberteri</i> (DC.) Urb.	CU ¹ , HT ¹ , JM ¹
<i>Dalbergiabignoniae</i> Berhaut	GW, CI, SN, TG
<i>Dalbergiabintuluensis</i> Sunarno & Ohashi	MY
<i>Dalbergiaboehmii</i> subsp. <i>stuhlmannii</i> (Taub.) Polhill	
<i>Dalbergiaboehmii</i> Taub.	AO, CM, CF, CD, GW, KE, MW ³ , MY, MZ, SL, SD, TZ, ZM, ZW
<i>Dalbergiabojeri</i> Drake	MG
<i>Dalbergiaboniana</i> Gagnep.	VN
<i>Dalbergiaborneensis</i> Prain	ID, MY
<i>Dalbergia brachystachya</i> Bosser & R. Rabev.	MG
<i>Dalbergiabraceolata</i> Baker	KE, MG, MZ, TZ
<i>Dalbergiabrasiliensis</i> Vogel	BR
<i>Dalbergiabrownei</i> (Jacq.) Urb.	BZ, CO, CR, CU ¹ , DO ¹ , EC ¹ , SV ¹ , GT, HT ¹ , HN ¹ , JM ¹ , MX, NI, PA, TT ¹ , GB ¹ , US, VE
<i>Dalbergiabormanica</i> Prain	CN, MM
<i>Dalbergiacalderonii</i> Standl.	BZ ¹ , SV, GT ¹ , HN ¹ , MX, NI
<i>Dalbergiacalycina</i> Benth.	BZ ² , CR ¹ , SV ¹ , GT, HN ¹ , MX, NI
<i>Dalbergiacambodiana</i> Pierre	KH, VN
<i>Dalbergiacampenonii</i> Drake	MG
<i>Dalbergiacana</i> Kurz	LA ² , MM, TH
<i>Dalbergiacandenatensis</i> (Dennst.) Prain	AU, BD, BN, KH ¹ , CN, FJ, IN, ID, MY, FM, PG, PH, MM, SG, SB, LK, TH, VN
<i>Dalbergiacanescens</i> (Elmer) Merr.	MY, PH
<i>Dalbergia capuronii</i> Bosser & R. Rabev.	MG

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiacarringtoniana</i> Sousa	AO
<i>Dalbergiacatingicola</i> Harms	BR
<i>Dalbergiacaudata</i> G.Don	
<i>Dalbergiacearensis</i> Ducke	BR, MX ³
<i>Dalbergiachapelier</i> Baill.	MG
<i>Dalbergiachlorocarpa</i> R.Vig.	MG
<i>Dalbergiachontalensis</i> Standl. &L.O.Williams	SV ¹ , GT ¹ , NI
<i>Dalbergiaclarkei</i> Thoth.	IN
<i>Dalbergiacochinchinensis</i> Laness.	KH, LA ² , PG, SB, TH, VN
<i>Dalbergiacochinchinensis</i> Pierre	KH, LA, PG, SB, TH, VN
<i>Dalbergiacommiphorooides</i> Baker f.	ET, KE, SO
<i>Dalbergia comorensis</i> Bosser & R. Rabev.	KM ¹
<i>Dalbergiaconfertiflora</i> Benth.	BD, IN, MX
<i>Dalbergiacongensis</i> Baker f.	AO, CF
<i>Dalbergiacongesta</i> Wight & Arn.	IN
<i>Dalbergiacongestiflora</i> Pittier	SV ¹
<i>Dalbergiacoromandeliana</i> Prain	IN
<i>Dalbergiacrispa</i> Hepper	NG, SL
<i>Dalbergiacubilquitzensis</i> (Donn.Sm.) Pittier	BZ ¹ , CR, MX, NI
<i>Dalbergiacucullata</i> Pittier	VE
<i>Dalbergiacuiabensis</i> Benth.	BO, BR
<i>Dalbergiacultrata</i> Benth.	KH ² , CN ⁵ , IN, LA, MM, TH, VN
<i>Dalbergiacumingiana</i> Benth.	PH
<i>Dalbergiacurtisii</i> Prain	ID, LA, MY, TH, VN
<i>Dalbergiacuscatlanica</i> (Standl.) Standl.	CR, SV ¹ , GT, MX, PA
<i>Dalbergiadalzielii</i> Hutch. & Dalziel	CM, GA, NG, TG
<i>Dalbergiadariensis</i> Rudd	PA
<i>Dalbergia davidii</i> Bosser & R.Rabev.	MG

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergia debilis</i> J.F.Macbr.	PE
<i>Dalbergiadelphipularis</i> Rizzini&A.Mattos	BR
<i>Dalbergiadelphinensis</i> Bosser&R.Rabev.	MG
<i>Dalbergiadensa</i> Benth.	AU, ID, PG, PH
<i>Dalbergiadensiflora</i> (Benth.) Benth.	BR
<i>Dalbergiadialoides</i> (Pierre) Niyomdham	
<i>Dalbergiadicolor</i> Blume	ID, MY, PH, TH
<i>Dalbergiadongnaiensis</i> Pierre	KH, TH, VN
<i>Dalbergiaduarensis</i> Thoth.	IN
<i>Dalbergiaduperreana</i> Pierre	KH, TH
<i>Dalbergiadyeriana</i> Harms	CN
<i>Dalbergiaealaensis</i> De Wild.	AO, CM, CD, GA
<i>Dalbergiaecastaphyllum</i> (L.) Taub.	AO, AG, BS, BB, BZ, BR, CM, CO, CR, CU, DM, DO, GQ, FR, GM, GH, GD, GT, GN, GW, GY, HT, HN ¹ , IN, CI, JM ¹ , LR, MU ³ , MX, NI, NG, NL, PA, PY ¹ , PE, VC, ST, SN, SL, SR, TG, TT, GB, US
<i>Dalbergia elegans</i> A.M.Carvalho	BR
<i>Dalbergiaemirnisensis</i> Benth.	MG
<i>Dalbergiaenneaphylla</i> Pittier	VE
<i>Dalbergiaentadoides</i> Prain	KH, LA, TH, VN
<i>Dalbergiaeremicola</i> Polhill	KE, SO
<i>Dalbergiaernest-ulei</i> Hoehne	BR
<i>Dalbergiaerrans</i> Craib	TH
<i>Dalbergia erubescens</i> Bosser & R.Rabev.	MG
<i>Dalbergiafalcata</i> Prain	BN, ID, MY
<i>Dalbergiafischeri</i> Taub.	MW, MZ, TZ, ZM, ZW
<i>Dalbergia floribunda</i> Craib	TH
<i>Dalbergiaflorifera</i> De Wild.	AO, CF, CD
<i>Dalbergiafoliolosa</i> Benth.	BO ¹ , BR

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergia foliosa</i> (Benth.) A.M.Carvalho	BR, CO, FR, GY ¹ , SR, VE
<i>Dalbergiaforbesii</i> Prain	ID, LA, TH
<i>Dalbergiafouilloiyana</i> Pellegr.	GA
<i>Dalbergiafrutescens</i> (Vell.) Britton	AR, BO, BR, CO ¹ , CR ¹ , EC ¹ , GY, PY, PE, VE
<i>Dalbergia frutescens</i> var. <i>tomentosa</i> (Vogel) Benth.	BO ¹ , BR, SR, VE ⁴
<i>Dalbergiafunera</i> Standl.	SV, GT
<i>Dalbergiafusca</i> Pierre	CN, MM ¹ , VN ¹
<i>Dalbergiagardneriana</i> Benth.	IN
<i>Dalbergiagentilii</i> De Wild.	CD
<i>Dalbergiagilbertii</i> Cronquist	CD
<i>Dalbergia glaberrima</i> Bosser & R.Rabev.	MG
<i>Dalbergiaglabra</i> (Mill.) Standl.	BZ, CR, SV ¹ , GT, HN, MX, NI
<i>Dalbergiaglandulosa</i> Benth.	BR
<i>Dalbergiaglaucescens</i> (Benth.) Benth.	BR
<i>Dalbergia glaucocarpa</i> Bosser & R.Rabev.	MG
<i>Dalbergiaglaziovii</i> Harms	BR
<i>Dalbergiaglomerata</i> Hemsl.	CR ¹ , GT ² , HN ² , MX
<i>Dalbergiagodefroyi</i> Prain	KH, LA, TH
<i>Dalbergiagossweleri</i> Baker f.	AO
<i>Dalbergiagracilis</i> Benth.	BO, BR
<i>Dalbergiagranadillo</i> Pittier	SV ¹ , MX
<i>Dalbergiagrandibracteata</i> De Wild.	CD, GN
<i>Dalbergia grandistipula</i> A.M.Carvalho	BR
<i>Dalbergiagreveana</i> Baill.	MG ²
<i>Dalbergia guttembergii</i> A.M.Carvalho	BR
<i>Dalbergiahainanensis</i> Merr. & Chun	CN
<i>Dalbergiahance</i> Benth.	CN, TH, VN
<i>Dalbergiahavilandii</i> Prain	BN, MY

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiahenryana</i> Prain	CN
<i>Dalbergiaheudelotii</i> Stapf	CD, GA, GH, GN, CI, LR, NG, SN, SL
<i>Dalbergiahiemalis</i> Malme	BR
<i>Dalbergiahildebrandtii</i> var. <i>scorpioides</i> (Baker) Bosser&R.Rabev.	MG
<i>Dalbergiahildebrandtii</i> Vatke	MG
<i>Dalbergiahirticalyx</i> Bosser&R.Rabev.	MG
<i>Dalbergiahorrida</i> (Dennst.) Mabb.	IN, LA, TH, VN
<i>Dalbergia horrida</i> var. <i>glabrescens</i> (Prain) Thoth. & Nair	
<i>Dalbergiahortensis</i> Heringer& al.	BR
<i>Dalbergiahoseana</i> Prain	MY
<i>Dalbergiahostilis</i> Benth.	AO, BJ, CM, CF, CD, GQ, GA, GH, GN, GW, CI, LR, NG, SL, TG
<i>Dalbergiahullettii</i> Prain	ID, MY, SG
<i>Dalbergiahumbertii</i> R.Vig.	MG
<i>Dalbergiahupeana</i> Hance	CN, LA, VN
<i>Dalbergiahygrophila</i> (Benth.) Hoehne	BR, CO ⁴ , PE, VE
<i>Dalbergia intermedia</i> A.M.Carvalho	BR, GY ¹ , VE ¹
<i>Dalbergiaintibucana</i> Standl. &L.O.Williams	HN
<i>Dalbergiainundata</i> Benth.	BR, CO, GY, PE, VE
<i>Dalbergiaiquitosensis</i> Harms	BR, PE
<i>Dalbergiajahaerii</i> Burck	ID
<i>Dalbergia jingxiensis</i> S.Y. Liu	CN ¹
<i>Dalbergiajunghunii</i> Benth.	ID, MY, SG
<i>Dalbergiakerrii</i> Craib	LA, TH
<i>Dalbergiakingiana</i> Prain	CN, MM
<i>Dalbergia kisantuensis</i> De Wild. & T.Durand	AO, CD, GA
<i>Dalbergiakostermansii</i> Sunamo&Ohashi	ID, MY
<i>Dalbergiakunstleri</i> Prain	MY

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiakurzii</i> Prain	TH
<i>Dalbergialacei</i> Thoth.	MM, TH
<i>Dalbergialactea</i> Vatke	BI, CM, CD, ET, GA, KE, MW, MZ, NG, RW, SD, TZ, UG, ZM, ZW
<i>Dalbergialakhonensis</i> Gagnep.	LA, TH
<i>Dalbergialanceolaria</i> L.f.	BD, KH, HN ¹ , IN, PK, MM, LK, TH, VN
<i>Dalbergia lanceolaria</i> subsp. <i>paniculata</i> (Roxb.) Thoth.	KH, IN, LA, NP, MM, TH, VN
<i>Dalbergialanceolaria</i> var. <i>maymyensis</i> (W. G. Craib) Thoth.	
<i>Dalbergialastoursvillensis</i> Pellegr.	GA
<i>Dalbergialateriflora</i> Benth.	BR
<i>Dalbergialatifolia</i> Roxb.	BD ² , FR, IN, ID, KE, MY, MU, NP, NG, PK, SG, LK, TZ, UG
<i>Dalbergialaxiflora</i> Micheli	CD
<i>Dalbergia lemurica</i> Bosser & R.Rabev.	MG
<i>Dalbergialibrevillensis</i> Pellegr.	GA
<i>Dalbergia longepedunculata</i> J. Linares & M. Sousa	HN ¹ , MX ¹
<i>Dalbergialouisii</i> Cronquist	BJ ² , CM ² , CG ² , CD, GA ² , NG
<i>Dalbergialouvelii</i> R.Vig.	MG
<i>Dalbergia luteola</i> J. Linares & M. Sousa	GT ¹ , MX ¹
<i>Dalbergiamacrosperma</i> Baker	AO
<i>Dalbergia madagascariensis</i> var. <i>poolii</i> (Baker) Bosser & R.Rabev.	MG
<i>Dalbergiamadagascariensis</i> Vatke	MG
<i>Dalbergiamalabarica</i> Prain	IN
<i>Dalbergiamalangensis</i> Sousa	AO, TZ, ZM
<i>Dalbergiamammosa</i> Pierre	VN
<i>Dalbergiamarcaniana</i> Craib	TH
<i>Dalbergiamaritima</i> R.Vig.	MG
<i>Dalbergiamartini</i> F.White	NA, ZM, ZW
<i>Dalbergiamayumbensis</i> Baker f.	AO

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiamelanocardium</i> Pittier	BZ ¹ , CR, SV, GT, HN ¹ , MX, NI ¹
<i>Dalbergiamelanoxylon</i> Guill. &Perr.	AO, BW ¹ , BF, BI ³ , CF, TD, CD, ET, IN, CI, KE, MW, ML, MZ, NA ² , NG, SN, ZA, SS ² , LK, SD, TZ, UG, ZM, ZW
<i>Dalbergiamenooides</i> Prain	ID, MY
<i>Dalbergiamexicana</i> Pittier	MX ¹
<i>Dalbergiamicrophylla</i> Chiov.	ET, KE, SO, TZ
<i>Dalbergiamillettii</i> Benth.	BT, CN, IN, MM
<i>Dalbergiamillettii</i> var. <i>mimosoides</i> (Franch.) Thoth.	BT, IN, MM
<i>Dalbergiamimosella</i> (Blanco) Prain	ID, MY, PH
<i>Dalbergiamimosoides</i> Franch.	CN, IN ¹
<i>Dalbergiamiscobium</i> Benth.	BR
<i>Dalbergia modesta</i> J. Linares & M. Sousa	MX ¹
<i>Dalbergia mollis</i> Bosser & R.Rabev.	MG
<i>Dalbergiamollis</i> var. <i>menabeensis</i> (R.Vig.) Bosser&R.Rabev.	MG
<i>Dalbergiamonetaria</i> L.f.	BZ, BO ¹ , BR, CO, CR, CU, DO, EC ¹ , FR, GT, GY, HT, HN, JM, MX, NI, PA, PE, LC ² , VC, SR, TT, US ¹ , VE
<i>Dalbergiamonophylla</i> G.A.Black	BR
<i>Dalbergia monticola</i> Bosser & R.Rabev.	MG
<i>Dalbergiamultijuga</i> E.Mey.	ZA
<i>Dalbergianegrensis</i> (Radlk.) Ducke	BR
<i>Dalbergia neoperrieri</i> Bosser & R.Rabev.	MG
<i>Dalbergiangounyensis</i> Pellegr.	CD, GA, SL
<i>Dalbergianigra</i> (Vell.) Benth.	BR
<i>Dalbergianitida</i> (Benth.) Hoehne	BO, TZ
<i>Dalbergianitidula</i> Baker	AO, BI, CD, MW, MZ, RW, ZA, UG, ZM, ZW
<i>Dalbergianoideae</i> Harms	AO, GW
<i>Dalbergia normandii</i> Bosser & R.Rabev.	MG ²
<i>Dalbergiaoblongifolia</i> G.Don	GH, GW, CI, LR, SN, SL
<i>Dalbergiaobovata</i> E.Mey.	MZ, ZA, TZ

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiaobtusifolia</i> (Baker) Prain	CN, MM, TH
<i>Dalbergia odorifera</i> T.C.Chen	CN
<i>Dalbergiaoligophylla</i> Hutch. & Dalziel	CM, NG, SL
<i>Dalbergiaoliveri</i> Prain	IN, MY, MM, TH, VN
<i>Dalbergia orientalis</i> Bosser & R.Rabev.	MG
<i>Dalbergia ovata</i> Benth.	BD, LA ² , MM, TH, VN
<i>Dalbergia ovata</i> var. <i>glomeriflora</i> (Kurz) Thoth.	MM, TH
<i>Dalbergiapachycarpa</i> (De Wild. & T.Durand) De Wild.	AO, CM, CD
<i>Dalbergiapalo-escrito</i> Rzed.	MX
<i>Dalbergiaparviflora</i> Roxb.	BN, ID, LA, MY, MM, SG, TH
<i>Dalbergiapaucifoliolata</i> Lundell	MX ¹
<i>Dalbergiapeguensis</i> Thoth.	MM
<i>Dalbergiapeishaensis</i> Chun & T.Chen	CN
<i>Dalbergia peltieri</i> Bosser & R.Rabev.	MG
<i>Dalbergiapervillei</i> Vatke	MG
<i>Dalbergiapierreana</i> Prain	KH
<i>Dalbergiapinnata</i> (Lour.) Prain	BD, BT, CN, IN, ID, LA, MY, NP, PG, PH, MM, TH, VN
<i>Dalbergia pinnata</i> var. <i>acaciifolia</i> (Dalzell) Thoth.	IN
<i>Dalbergiapluriflora</i> Baker f.	AO
<i>Dalbergiapolyadelpa</i> Prain	CN
<i>Dalbergiapolyphylla</i> Benth.	PH
<i>Dalbergiaprainii</i> Thoth.	MM
<i>Dalbergia pseudo-ovata</i> Thoth.	MM
<i>Dalbergia pseudo-sissoo</i> Miq.	IN, MY, LK, TH
<i>Dalbergiapseudobaronii</i> R.Vig.	MG
<i>Dalbergiapurpurascens</i> Baill.	
<i>Dalbergiareniformis</i> Roxb.	BD, IN, MM
<i>Dalbergiareticulata</i> Merr.	PH

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergia retusa</i> Hemsl.	BZ ² , CO ¹ , CR, SV, GT ¹ , HN ¹ , MX, NI, PA
<i>Dalbergia revoluta</i> Ducke	BR, VE
<i>Dalbergia rhachiflexa</i> J. Linares & M. Sousa	MX ¹
<i>Dalbergia richardsii</i> Sunarno & Ohashi	MY
<i>Dalbergia riedelii</i> (Benth.) Sandwith	BO ¹ , BR, CO, EC ¹ , FR ¹ , GY, PE, SR, VE
<i>Dalbergia rimosa</i> Roxb.	BD, BT, KH ¹ , CN, IN, ID ¹ , LA, MY ⁵ , MM, TH, VN
<i>Dalbergia rimosa</i> var. <i>foliacea</i> (Benth.) Thoth.	MM, TH
<i>Dalbergia riparia</i> (Mart.) Benth.	BO ¹ , BR, EC ¹ , GY, PE
<i>Dalbergia rostrata</i> Hassk.	BN, IN, ID, MY, PH, SG, LK
<i>Dalbergia rubiginosa</i> Roxb.	CN ³ , IN
<i>Dalbergia ruddiae</i> J. Linares & M. Sousa	CR ¹ , MX ¹
<i>Dalbergia rufa</i> G. Don	CM, CD, GN, GW, NG, SN, SL
<i>Dalbergia rugosa</i> Hepper	GR, LR, SL
<i>Dalbergia sacerdotum</i> Prain	CN
<i>Dalbergia salvanaturae</i> J. Linares & M. Sousa	SV ¹
<i>Dalbergia sambesiaca</i> Schinz	MZ
<i>Dalbergia sampaoana</i> Kuhlm. & Hoehne	BR
<i>Dalbergia sandakanensis</i> Sunarno & Ohashi	MY
<i>Dalbergia saxatilis</i> Hook.f.	AO, CM, CD, GA, GH, GW, CI, LR, NG, SN, SL, TG
<i>Dalbergia saxatilis</i> var. <i>preussii</i> (Harms) Cronquist	
<i>Dalbergia scortechinii</i> (Prain) Prain	ID, MY
<i>Dalbergia sericea</i> G. Don	BT, CN, KE ³ , NP, MM ⁵ , VN ¹
<i>Dalbergia setifera</i> Hutch. & Dalziel	GH, CI
<i>Dalbergia simpsonii</i> Rudd	PE
<i>Dalbergia sissooides</i> Wight & Arn.	IN, ID
<i>Dalbergia sissoo</i> DC.	AF, AU, BD, BT, TD, CN, DO, GH, GW, HN ¹ , IN, ID, IR, IQ, KE, MY, MZ, NP, NE, NG, OM, PK, MM, SN, SL, SD, TZ, UG, US, ZM, ZW
<i>Dalbergia spinosa</i> Roxb.	BD, IN, KY, MM

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiaspruceana</i> (Benth.) Benth.	BO, BR, VE
<i>Dalbergiastenophylla</i> Prain	CN, TH ⁵ , VN ¹
<i>Dalbergiastercoracea</i> Prain	ID, MY, SG
<i>Dalbergiastevensonii</i> Standl.	BZ, GT, HN ¹ , MX
<i>Dalbergiastipulacea</i> Roxb.	BD, BT, KH ¹ , CN, IN, ID, LA, MY, NP, PG, PH, MM, SB, TH, VN
<i>Dalbergiasuaresensis</i> Baill.	MG
<i>Dalbergiasubcymosa</i> Ducke	BO ¹ , BR, FR, GY, PE, SR, VE
<i>Dalbergiasuccirubra</i> Gagnep. & Craib	TH
<i>Dalbergiateijsmannii</i> Sunarno&Ohashi	ID
<i>Dalbergiateixeirae</i> Sousa	AO
<i>Dalbergiathomsonii</i> Benth.	IN
<i>Dalbergiathorelii</i> Gagnep.	KH, LA, TH
<i>Dalbergiatilarana</i> N.Zamora	CR, MX ¹ , NI, PA
<i>Dalbergiatinnevelliensis</i> Thoth.	IN
<i>Dalbergiatonkinensis</i> Prain	CN, VN
<i>Dalbergiatravancorica</i> Thoth.	IN
<i>Dalbergiatrichocarpa</i> Baker	MG
<i>Dalbergiatricolor</i> Drake	MG
<i>Dalbergia tsaratananensis</i> Bosser & R.Rabev.	MG
<i>Dalbergiatsiandalana</i> R.Vig.	MG
<i>Dalbergiatso</i> Merr. & Chun	CN
<i>Dalbergiatucurensis</i> Donn.Sm.	BZ, CR, SV ¹ , GT, HN, MX, NI ¹
<i>Dalbergiauarandensis</i> (Chiov.) Thulin	SO
<i>Dalbergia urschii</i> Bosser & R.Rabev.	MG
<i>Dalbergiavacciniifolia</i> Vatke	KE, TZ
<i>Dalbergiavelutina</i> Benth.	BD, IN, ID, LA, MY, MM, SG, TH
<i>Dalbergia velutina</i> var. <i>succirubra</i> (Gagnepain & Craib) Niyomdham	

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu, T.H., 2010

ACCEPTED SPECIES NAMES	RANGE STATES
<i>Dalbergiaverrucosa</i> Craib	TH
<i>Dalbergia viguieri</i> Bosser & R.Rabev.	MG
<i>Dalbergiavillosa</i> (Benth.) Benth.	BO ¹ , BR
<i>Dalbergia villosa</i> var. <i>barretoana</i> (Hoehne) A.M.Carvalho	BR
<i>Dalbergiavolubilis</i> Roxb.	BD, CN ¹ , IN, LA, NP, LK, TH, VN ³
<i>Dalbergiawattii</i> C.B.Clarke	IN
<i>Dalbergia xerophila</i> Bosser & R.Rabev.	MG
<i>Dalbergia ximengensis</i> Y.Y. Qian	CN ¹
<i>Dalbergiayunnanensis</i> Franch.	CN, IN ¹
<i>Dalbergiayunnanensis</i> var. <i>collettii</i> (Prain) Thoth.	MM

¹www.tropicos.org

²www.iucnredlist.org

³Saha *et al.*, 2013

⁴Grandter & Chevrette, 2013

⁵Shu,T.H., 2010