

CONVENTION SUR LE COMMERCE INTERNATIONAL DES ESPECES  
DE FAUNE ET DE FLORE SAUVAGES MENACEES D'EXTINCTION



Soixante-dixième session du Comité permanent  
Rosa Khutor, Sochi (Fédération de Russie), 1 – 5 octobre 2018

Questions spécifiques aux espèces

GRANDS SINGES (*HOMINIDAE SPP.*)

1. Le présent document a été préparé par le Secrétariat.

Résolution Conf. 13.4 (Rev CoP16)

2. Dans la résolution Conf. 13.4 (Rev. CoP16), *Conservation et commerce des grands singes*, la Conférence des Parties CHARGE le Secrétariat de faire rapport au Comité permanent sur l'application de la présente résolution à chacune de ses sessions ordinaires [paragraphe 2 d)], et CHARGE le Comité permanent d'examiner l'application de la présente résolution à chaque session ordinaire en s'appuyant sur les rapports du Secrétariat [paragraphe 3 a)].
3. Au paragraphe 4 de cette résolution, la Conférence des Parties prie instamment le Secrétariat, le Comité permanent et le Comité pour les animaux de travailler en étroite collaboration avec le GRASP<sup>1</sup> (GReat Apes Survival Partnership) une alliance regroupant plus de 100 gouvernements nationaux, organisations œuvrant en faveur de la conservation, instituts de recherche, agences des Nations Unies et entreprises privées, qui se sont engagés à assurer la survie à long terme des grands singes et de leurs habitats en Afrique et en Asie. Un document d'information distribué lors de la présente session fait le point sur les activités du GRASP.
4. Les membres du GRASP ne se sont pas réunis depuis la dernière session du Comité permanent en novembre 2017. Toutefois, le Secrétariat est toujours un membre actif du Comité exécutif du GRASP et il a maintenu des contacts réguliers avec le Secrétariat du GRASP, en particulier pour s'assurer que les décisions 17.232 et 17.233 sont pleinement mises en œuvre.

Mise en œuvre des décisions 17.232 et 17.233

5. À sa 17<sup>e</sup> session (CoP17, Johannesburg, 2016), la Conférence des Parties a adopté les décisions 17.232 et 17.233 sur les *Grand singes* comme suit:

**À l'adresse du Secrétariat**

17.232 *Le Secrétariat collabore avec le groupe CSE/UICN de spécialistes des primates, le GRASP et d'autres spécialistes, et, sous réserve que des fonds suffisants soient disponibles, produit un rapport relatif à l'état des grands singes et à l'impact relatif du commerce illégal et des autres pressions sur cet état, en vue d'un examen par le Comité permanent.*

---

<sup>1</sup> GRASP = Great Apes Survival Partnership, une initiative des Nations Unies engagée en faveur de la survie à long terme des chimpanzés, gorilles, bonobos, and orang-outans, ainsi que de leurs habitats en Afrique et en Asie ([www.un-grasp.org](http://www.un-grasp.org))

## **À l'adresse du Comité permanent**

17.233 *Le Comité permanent examine le rapport préparé conformément à la décision 17.232 et, s'il y a lieu, élabore des recommandations pour des actions futures qui seront examinées à la 18<sup>e</sup> session de la Conférence des Parties.*

6. Lors de la 69<sup>e</sup> session du Comité permanent (SC69, Genève, novembre 2017), le Secrétariat a fait une présentation orale sur les activités menées en rapport avec la résolution Conf. 13.4 (Rev. CoP16). Il a expliqué que, pour ce qui est de la décision 17.232, le rapport aurait dû être soumis à la présente session, mais que, pour diverses raisons, cela n'a pas été possible. Pour faire avancer l'étude, le GRASP, le Secrétariat de la CITES et l'Union internationale pour la conservation de la nature (UICN) ont proposé un calendrier révisé, lequel a été accepté par le Comité permanent, de sorte que le projet final du rapport de situation puisse être présenté pour examen à la présente réunion du Comité permanent.
7. Comme l'étude demandée dans la décision 17.232 est essentiellement de nature scientifique, il semble qu'elle bénéficierait largement de la contribution du Comité pour les animaux. Le président du Comité pour les animaux a déclaré que le Comité pour les animaux serait heureux d'examiner et de commenter l'étude, une fois reçue. Le Comité permanent a demandé au Secrétariat de présenter au Comité pour les animaux pour examen le rapport qu'il a commencé à préparer conformément à la décision 17.232.
8. Le Secrétariat du GRASP a préparé un projet de rapport en collaboration avec un certain nombre d'experts, notamment ceux du groupe de spécialistes des primates de la Commission de la sauvegarde des espèces (CSE) de l'UICN, et l'a soumis pour examen et commentaires au Comité pour les animaux lors de sa 30<sup>e</sup> session (voir le document AC30 Doc. 26).
9. Par l'intermédiaire d'un groupe de travail en session, le Comité pour les animaux a recommandé que le Secrétariat fournisse une série d'observations aux auteurs sur la manière de mettre à jour le rapport et d'en améliorer la présentation [voir le document AC30 Com 2 (Rev. par Sec.)].
10. Un rapport final révisé a été préparé pour la soixante-dixième session du Comité permanent, compte tenu des commentaires du Comité pour les animaux. Ce rapport figure en annexe du présent document.
11. Pour faciliter la mise en œuvre de la décision 17.233, le Comité permanent a créé un groupe de travail intersessions sur les grands singes lors de la soixante-neuvième session du Comité permanent doté du mandat suivant : examiner le rapport finalisé par le Secrétariat conformément à la décision 17.232 et toute recommandation émanant du Comité pour les animaux; et faire rapport au Comité permanent lors de sa soixante-dixième session avec tout projet de recommandation. Le groupe est composé des membres suivants: Canada (présidence), Chine, États-Unis d'Amérique, France, Indonésie, Malaisie, Nigéria, Royaume-Uni de Grande-Bretagne et d'Irlande du Nord, Thaïlande et Union européenne; et *Animal Advocacy and Protection (AAP)*, *Born Free Foundation*, Convention sur les espèces migratrices, Organisation des Nations Unies pour l'alimentation et l'agriculture, TRAFFIC, Union internationale pour la conservation de la nature, *Wildlife Conservation Society*, *Wildlife Impact* et *World Association of Zoos and Aquariums*.
12. Malheureusement, en raison du court intervalle de temps entre la fin de la session du Comité pour les animaux et la date de soumission des documents pour la présente session, le groupe de travail intersessions n'a pas été en mesure d'achever son mandat en temps voulu pour soumettre un rapport écrit à la présente session.
13. Maintenant que le document est disponible, le groupe de travail intersessions a confirmé qu'il poursuivra ses travaux jusqu'à la présente session afin de s'acquitter des tâches décrites dans son mandat. Le président du groupe de travail présentera un rapport oral lors de cette session.

### Recommandations

14. Le Comité permanent est invité à:
  - a) examiner le rapport figurant en annexe, compte tenu de toute information émanant du groupe de travail intersessions du Comité permanent sur les grands singes, et à examiner les recommandations concernant les mesures à venir qui pourraient s'avérer utiles;

- b) envisager de demander au Secrétariat, en consultation avec le GRASP et le président du Comité permanent, de revoir les dispositions actuelles figurant dans la résolution Conf. 13.4 (Rev. CoP16) *Conservation et commerce des grands singes* et, compte tenu des recommandations issues du rapport, proposer des amendements appropriés à apporter à la résolution, pour examen à la 18<sup>e</sup> session de la Conférence des Parties ; et
- c) confirmer que les décisions 17.232 et 17.233 ont été totalement mises en œuvre.

## **Great Apes Status Report (August 2018)**

**Great Apes Survival Partnership & International Union for Conservation of Nature**

**Editors: J. Refisch<sup>1</sup>, S.A. Wich<sup>2,3,4</sup> & E.A. Williamson<sup>3,5</sup>**

**Population data analysis, compilation and maps: T. Sop, IUCN SSC A.P.E.S. database manager<sup>6</sup>**

**With contributions and reviews by (in alphabetical order): M. Ancrenaz<sup>3,4,7</sup>, R.A. Bergl<sup>3,4,8</sup>,  
D. Greer<sup>3,9</sup>, T. Humle<sup>3,10</sup>, R. Kormos<sup>11</sup>, H.S. Kühl<sup>3,4,6</sup>, E.J. Macfie<sup>3</sup>, F. Maisels<sup>3,5,12</sup>,  
J.F. Oates<sup>3,13</sup>, M.M. Robbins<sup>4,6</sup>, J. Sherman<sup>14</sup> & T. Sop<sup>6</sup>**

<sup>1</sup>Great Apes Survival Partnership, United Nations Environment Programme (UNEP), P.O. Box 30552, 00100 Nairobi, Kenya

<sup>2</sup>John Moores University Liverpool, School of Natural Sciences and Psychology, Liverpool L3 5UA, UK

<sup>3</sup>Executive Committee of IUCN SSC Primate Specialist Group (PSG) Section on Great Apes (SGA)

<sup>4</sup>Scientific Commission of Great Apes Survival Partnership (GRASP)

<sup>5</sup>Faculty of Natural Sciences, University of Stirling, Stirling, FK9 4LA, UK

<sup>6</sup>Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany

<sup>7</sup>HUTAN – Kinabatangan Orangutan Conservation Programme, P.O. Box 17793-88874, Kota Kinabalu, Malaysia

<sup>8</sup>North Carolina Zoo, 4401 Zoo Parkway, Asheboro, North Carolina, USA

<sup>9</sup>WWF Uganda, Kampala, Uganda

<sup>10</sup>Durrell Institute of Conservation and Ecology (DICE), School of Anthropology and Conservation, University of Kent, Canterbury CT2 7NR, UK

<sup>11</sup>Department of Integrative Biology, University of California, Berkeley, California, USA

<sup>12</sup>Wildlife Conservation Society, Global Conservation Program, 2300 Southern Boulevard, Bronx, New York, NY 10460, USA

<sup>13</sup>Department of Anthropology, Hunter College, 695 Park Avenue, New York, NY 10065, USA

<sup>14</sup>Wildlife Impact, P.O. Box 31062, Portland, OR 97231, USA

Citation: GRASP & IUCN (2018). Report to the CITES Standing Committee on the Status of Great Apes. United Nations Environment Programme Great Apes Survival Partnership, Nairobi, and International Union for Conservation of Nature, Gland.

## Table of contents

<b>1. Introduction</b>	p. 3
<b>2. African great apes</b>	p. 3
2.1. <i>Current distribution of African great apes</i>	p. 3
2.2. <i>African great ape population sizes</i>	p. 4
2.2.1. Population size estimates per country	p. 4
2.2.2. Population size estimates per taxon	p. 6
2.3. <i>Temporal trends in African great ape populations</i>	p. 7
2.4. <i>Threats to African great apes</i>	p. 9
<b>3. Asian great apes</b>	p. 12
3.1. <i>Current distribution of Asian great apes</i>	p. 12
3.2. <i>Asian great ape population sizes</i>	p. 13
3.2.1. Population size estimates per country	p. 13
3.2.2. Population size estimates per taxon	p. 14
3.3. <i>Temporal trends in Asian great ape populations</i>	p. 14
3.4. <i>Threats to Asian great apes</i>	p. 15
<b>4. Description of threats</b>	p. 16
<b>5. International agreements specific to great ape range States</b>	p. 19
<b>6. Challenges and solutions</b>	p. 20
<b>7. Recommendations</b>	p. 22
<b>8. Main sources of information</b>	p. 23
<b>9. Acknowledgements</b>	p. 23
<b>10. References</b>	p. 24
<b>Annex I. Acronyms</b>	p. 34

## Report to the CITES Standing Committee on the Status of Great Apes

### 1. Introduction

The 65<sup>th</sup> meeting of the Standing Committee of the Convention of International Trade in Endangered Species of Wild Fauna and Flora (CITES) mandated the CITES Secretariat (SC65, Doc37) to collaborate with the IUCN SSC Primate Specialist Group, the Great Apes Survival Partnership (GRASP) and other experts, to prepare a report on the status of great apes and the relative impact of illegal trade and other pressures on their status, for consideration by the Standing Committee. This report constitutes fulfilment of the mandate received from the CITES Secretariat and will address distribution and abundance trends related to all great ape species and sub-species, as well as threats to their conservation.

The great apes (bonobos, chimpanzees, eastern and western gorillas, Bornean, Sumatran and Tapanuli orangutans) face significant conservation threats and are listed as either Endangered or Critically Endangered by the IUCN Red List of Threatened Species (IUCN, 2018) and on Appendix I of CITES (CITES, 2017). All great apes have slow rates of reproduction due to their prolonged maturation and high investment in single (occasionally twin) offspring: age at first reproduction is late, their young take a long time to develop, and interbirth intervals are 4–9 years, depending on the species. Their populations are, therefore, highly vulnerable to even low levels of offtake, and unable to cope with significant and continued losses of individuals.

Great ape populations in Africa and Asia are threatened by the combined impacts of habitat loss, degradation and fragmentation, poaching, disease and illegal trade. It is illegal to kill or capture great apes and to trade live animals or their body parts in all great ape range States (IUCN, 2018). However, despite legal protection, law enforcement remains a major challenge in many countries, and poaching, especially for the illegal domestic (and some international, albeit mostly regional, between neighbouring countries) trade in bushmeat, is the most significant threat to the survival of most great apes (see Section 4 and Table 10).

This report presents the current distribution of African and Asian great apes, as well as temporal population trends (changes over time) and the main threats to their survival. It also highlights current conservation challenges and provides a list of recommendations to the CITES Parties, the CITES Secretariat and other relevant stakeholders.


### 2. African great apes

#### 2.1. Current distribution of African great apes

African great apes occur in 21 countries across Equatorial Africa (Figure 1). There are four species and nine taxa overall (Table 1).

**Table 1.** The African great apes

Genus	Species	Subspecies
<b>Gorilla</b> 2 species	<b>eastern gorilla</b> <i>Gorilla beringei</i> 2 subspecies	<b>Grauer's gorilla</b> <i>Gorilla beringei graueri</i>
		<b>mountain gorilla</b> <i>Gorilla beringei beringei</i>
	<b>western gorilla</b> <i>Gorilla gorilla</i> 2 subspecies	<b>Cross River gorilla</b> <i>Gorilla gorilla diehli</i>
		<b>western lowland gorilla</b> <i>Gorilla gorilla gorilla</i>
<b>Pan</b> 2 species	<b>chimpanzee</b> <i>Pan troglodytes</i> 4 subspecies	<b>central chimpanzee</b> <i>Pan troglodytes troglodytes</i>
		<b>eastern chimpanzee</b> <i>Pan troglodytes schweinfurthii</i>
		<b>Nigeria-Cameroon chimpanzee</b> <i>Pan troglodytes ellioti</i>
		<b>western chimpanzee</b> <i>Pan troglodytes verus</i>
	<b>bonobo</b> <i>Pan paniscus</i>	<b>No subspecies currently recognized</b>



**Figure 1.** Geographic distribution of African great ape taxa and sites (IUCN SSC A.P.E.S. database, 2017<sup>2</sup>), Max Planck Institute for Evolutionary Anthropology.

## 2.2. African great ape population sizes

The African great ape population estimates in this report are based on surveys conducted in the past 10 years. “Site” in this text refers to any area in which surveys were conducted in the last 10 years, including amongst others protected areas and their buffer zones, a logging concession or a group of concessions. Population estimates presented in this report are drawn from peer-reviewed publications, published or unpublished reports, data from research and conservation organizations, or are expert estimates. Estimating population size is complex, because great apes are difficult to observe. All great apes build a new nest to sleep in every night, and these can be used as proxies for independent nest-building animals. A software package (Distance) incorporates animal sign production (in this case night nests) and decay rate to calculate the density of animal sign and of population (Thomas, et al., 2010). Because sample size can be small where ape density is low, the resulting estimates of abundance often show a great deal of variation in precision (see Kühl, et al., 2008). At some sites, genetic censusing is used (e.g. Arandjelovic, et al., 2011; Gray, et al., 2013; Roy, et al., 2014). An overview of survey methods can be found in Kühl, et al. (2008).

### 2.2.1. Population size estimates per country

There is great disparity in population estimates from taxon to taxon depending on the survey method used and sampling effort. Chimpanzee, gorilla and bonobo population estimates per range State are presented in Table 2. The Democratic Republic of Congo (DRC) and Congo host the highest combined population sizes of great apes in Africa, followed by Cameroon, Guinea and Gabon. Countries with the lowest great ape population numbers include Burundi, Ghana, Rwanda, Mali and Senegal, which each hosts a few hundred great apes.

**Table 2.** African great ape population estimates by country. *Except for mountain gorillas, estimates are based on the number of “nest builders”, thus excluding infants. Estimates are derived from both surveys and modelling approaches.*

<sup>2</sup> Ape Populations, Environments and Surveys database (<http://apesportal.eva.mpg.de/>)

*This is the best information available. However, survey effort is often highest in protected areas, while other areas remain unsampled, thus can lead to low country estimates.*

Country	Taxon	Abundance	Survey period	Source
Angola	western lowland gorilla	1,652 (1,174–13,311)	2005–2013	Strindberg, et al. (2018)*
	central chimpanzee	1,705 (1,027–4,801)	2005–2013	
Burundi	eastern chimpanzee	204 (122–339)	2011–2013	Hakizimana & Huynen (2013)
Cameroon	Cross River gorilla	132–194	2014	Dunn, et al. (2014)
	western lowland gorilla	38,654 (34,331–112,881)	2005–2013	Strindberg, et al. (2018)*
	central chimpanzee	21,489 (18,575–40,408)	2005–2013	IUCN SSC A.P.E.S. database (2017); Strindberg, et al. (2018)
	Nigeria-Cameroon chimpanzee	3,000–7,060	2004–2006	Morgan, et al. (2011); Mitchell, et al. (2015); Oates, et al. (2016)
Central African Republic	western lowland gorilla	5529 (3,635–8,581)	2015	N’Goran, et al. (2016)
	central chimpanzee	2,843 (1,194–4,855)	2015	Strindberg, et al. (2018)
	eastern chimpanzee	910 (538–1,534)	2012–2016	Aebischer, et al. (2017)
Congo	western lowland gorilla	215,799 (180,814–263,913)	2005–2013	Strindberg, et al. (2018)*
	central chimpanzee	55,397 (42,433–64,824)	2005–2013	
Côte d’Ivoire	western chimpanzee	580 (332–940)	2007–2015	Tiédoúé, et al. (2016); IUCN SSC A.P.E.S. database (2017)
Democratic Republic of Congo	Grauer’s gorilla	3,800	2011–2015	Plumptre, et al. (2016a)
	mountain gorilla	N/A <sup>†</sup>		
	central chimpanzee	present (population size unknown)	N/A	Inogwabini, et al. (2007)
	eastern chimpanzee	173,000–248,000	2010	Plumptre, et al. (2010)
	bonobo	15,000–20,000 minimum	2012	IUCN & ICCN (2012)
Equatorial Guinea	western lowland gorilla	1,872 (1,082–3,165)	2005–2013	Strindberg, et al. (2018)*
	central chimpanzee	4,290 (2,894–7,985)	2005–2013	
Gabon	western lowland gorilla	99,245 (67,117–178,390)	2005–2013	Strindberg, et al. (2018)*
	central chimpanzee	43,037 (36,869–60,476)		
Ghana	western chimpanzee	264	2009	Danquah, et al. (2012)
Guinea	western chimpanzee	21,210 (10,007–43,534)	2009–2014	WCF (2012; 2014) Kühl, et al. (2017)
Guinea Bissau	western chimpanzee	1,000–1,500	2016	Chimbo Foundation (2017) unpubl. data
Liberia	western chimpanzee	7,008	2011–2012	Tweh, et al. (2015)

		(4,260–11,590)		
<b>Mali</b>	western chimpanzee	present (population size unknown)	2014	PanAf (2014) unpubl. data
<b>Nigeria</b>	Cross River gorilla	85–115	2014	Dunn, et al. (2014)
	Nigeria-Cameroon chimpanzee	730–2,095	2005	Morgan, et al. (2011); Oates, et al. (2016); S. Nixon, pers. comm.
<b>Rwanda</b>	mountain gorilla	N/A <sup>†</sup>		
	eastern chimpanzee	430	2009–2014	IUCN SSC A.P.E.S. database (2017)
<b>Sierra Leone</b>	western chimpanzee	5,580 (3,052–10,446)	2009	Brncic, et al. (2010)
<b>Senegal</b>	western chimpanzee	500–600	2016–2017	Pruetz & Wessling unpubl. data
<b>South Sudan</b>	eastern chimpanzee	present	2011	Plumptre, et al. (2016b)
<b>Tanzania</b>	eastern chimpanzee	2,500	2010–2012	Plumptre, et al. (2016b)
<b>Uganda</b>	mountain gorilla	400 <sup>†</sup>	2011	Roy, et al. (2014)
	eastern chimpanzee	5,000	2006–2017	Plumptre, et al. (2016b)

\* Strindberg, et al. (2018) estimate population sizes in 2013; western lowland gorilla numbers will have declined by a further 13% overall by the end of 2018.

<sup>†</sup> Bwindi mountain gorilla population only.

### 2.2.2. Population size estimates per taxon

Western lowland gorillas and eastern chimpanzees are the most numerous great ape taxa, while the Cross River gorilla has the smallest population size, with approximately 300 or fewer mature individuals remaining.

**Table 3.** African great ape population estimates by taxon in descending order of abundance. *Except for mountain gorillas, estimates are based on the number of “nest builders”, thus excluding infants. Estimates are derived from both surveys and modelling approaches. The IUCN Red List classification of most great ape taxa is based on the rate of decline over three generations (one generation time equating to 20–25 years, depending on the taxon).*

<b>Taxon</b>	<b>Abundance</b>	<b>IUCN Status</b>	<b>Source</b>
<b>western lowland gorilla</b> <i>Gorilla g. gorilla</i>	361,919 (302,973–460,093)*	Critically Endangered	Strindberg, et al. (2018)
<b>eastern chimpanzee</b> <i>Pan t. schweinfurthii</i>	181,000–256,000	Endangered	Plumptre, et al. (2010; 2016b)
<b>central chimpanzee</b> <i>Pan t. troglodytes</i>	128,760 (114,208–317,039)	Endangered	Strindberg, et al. (2018)
<b>western chimpanzee</b> <i>Pan t. verus</i>	18,000–65,000	Critically Endangered	Humle, et al. (2016); Kühl, et al. (2017)
<b>bonobo</b> <i>Pan paniscus</i>	15,000–20,000 minimum	Endangered	IUCN & ICCN (2012)
<b>Nigeria-Cameroon chimpanzee</b> <i>Pan t. ellioti</i>	4,400–9,345	Endangered	Morgan, et al. (2011); Oates, et al. (2016)
<b>Grauer’s gorilla</b> <i>Gorilla b. graueri</i>	3,800 (1,280–9,050)	Critically Endangered	Plumptre, et al. (2015; 2016b)
<b>mountain gorilla</b> <i>Gorilla b. beringei</i>	>1,000	Critically Endangered	Roy, et al. (2014); Hickey, et al. (in prep.) <sup>§</sup>
<b>Cross River gorilla</b> <i>Gorilla g. diehli</i>	<300	Critically Endangered	Dunn, et al. (2014); Bergl, et al. (2016); R. Bergl & J. Oates pers. comm.

\* Estimated in 2013. At an annual rate of decline of 2.7%, the population will be ~316,000 by the end of 2018.

<sup>§</sup> Hickey, et al. (in prep.) will report the results of the 2015–2016 survey in the Virungas; a survey of Bwindi Impenetrable National Park is still in progress.

### 2.3. Temporal trends in African great ape populations

With the exception of the mountain gorillas, all African great ape taxa are facing an overall decline, some of them drastic. The extent of the decline, however, differs between taxa, with Grauer's gorillas suffering the highest estimated losses over a short period of time (Table 4)<sup>3</sup>.

**Grauer's gorilla, *Gorilla beringei graueri***

In 1995, the Grauer's gorilla population was estimated at 16,900 (Hall, et al., 1998). By 2015, however, estimates placed their population at only 3,800 individuals, with an annual rate of decline of 7.2%. This represents a decline estimated at between 84–93% across the subspecies' range. Comparing the 1994 data to that of 2011–2015, researchers found a 77% reduction in abundance. The subspecies is classified as Critically Endangered. Source: Plumptre, et al. (2015; 2016a).

**Mountain gorilla, *Gorilla beringei beringei***

Whilst the mountain gorilla is the only great ape taxon increasing in numbers, it has been listed as Critically Endangered, with an estimated population of 880 individuals in the early 2010s, increasing to over 1,000 individuals in 2016. There are two small isolated populations, the Virunga and the Bwindi populations, both of which have been the subject of intense conservation, research and survey efforts over the last 40 years (e.g. Robbins, et al., 2011). Numbers in the Virungas dwindled to an estimated 250 individuals in 1981, before the population was allowed to recover and almost doubled between 1989 and 2010. Between 2003 and 2010, the population increased by 26% at an annual rate of 3.7% (Gray, et al., 2013).

**Cross River gorilla, *Gorilla gorilla diehli***

Cross River gorilla population changes are unknown. The subspecies is classified as Critically Endangered because of a small and highly fragmented population: each subpopulation is likely to number fewer than 50 mature individuals. Source: Bergl, et al. (2016).

**Western lowland gorilla, *Gorilla gorilla gorilla***

The Critically Endangered western lowland gorilla population declined by 19.4% between 2005 and 2013, an annual loss of approximately 2.7% (Strindberg, et al., 2018). Although their geographic range is large, they are threatened by bushmeat poaching, disease (including Ebola virus disease), and habitat loss and degradation.

**Central chimpanzee, *Pan troglodytes troglodytes***

This subspecies is classified as Endangered, having experienced a significant population reduction since the 1970s. The principal threats to this taxon, as for western lowland gorillas, are bushmeat poaching, disease, and habitat loss and degradation. An analysis of nest survey data collected between 2003 and 2013 across the entire geographic range did not detect a statistically significant decline (Strindberg, et al., 2018). Unlike gorillas, where an adult male will confront danger, chimpanzees slip away stealthily, and more often escape being killed. However, the factors known to reduce chimpanzee populations (lack of forest guards, remoteness from roads, human population density, degree of forest intactness, etc.) were found to be significantly correlated with population density in the direction predicted (Strindberg, et al., 2018). All of these factors are increasing in either extent or intensity, or both, so we predict that a decline will be statistically easier to detect in the future.

**Eastern chimpanzee, *Pan troglodytes schweinfurthii***

The Albertine Rift escarpment in DRC is a stronghold for eastern chimpanzees, but recent surveys indicate 80–98% declines at some key sites in just 20 years, contributing to the major population declines seen in recent decades. It is estimated that populations in eastern DRC declined significantly (range of estimated decline 22–45%) between 1994 and 2015. Eastern chimpanzees are listed as Endangered. Source: Plumptre, et al. (2016b).

**Nigeria-Cameroon chimpanzee, *Pan troglodytes ellioti***

This taxon has the lowest estimated population size of any chimpanzee subspecies. Its range is limited to clearly defined areas in southern Nigeria and central-southwest Cameroon. Its Endangered listing is based on an inferred population size reduction of between 50–80% over a three-generation period from the mid-1980s to 2060. Source: Oates, et al. (2016). Past presence of chimpanzees in Benin and Togo has been very poorly documented, but if chimpanzees were in those countries, it is probable that they belonged to this subspecies.

**Western chimpanzee, *Pan troglodytes verus***

This subspecies, *Pan troglodytes verus*, has recently been upgraded to Critically Endangered as it is expected to experience a decline exceeding 80% over the next 69 years, i.e. three generations (Humble et al., 2016). It is found in West Africa from Senegal to Ghana but has almost certainly become extinct in Burkina Faso in the 20<sup>th</sup> century (Ginn, et al., 2013; Campbell & Hounbedji, 2015). With a likelihood of probably more than 18,850 individuals, Guinea now hosts the largest remaining western chimpanzee population. Liberia and Sierra Leone are also strongholds for this subspecies

---

<sup>3</sup> Some text in Sections 2.3 and 3.3 is taken from the IUCN Red List assessments, as indicated by "Source".

(Kühl, et al., 2017). Approximately 17,000 individuals (half of the total western chimpanzee population) live in the Fouta Djallon region of Guinea (Regnaut & Boesch, 2012). Between 1990 and 2015, the Côte d’Ivoire population declined by 80% (Kühl, et al., 2017); only a few hundred individuals remain in Tai and Comoé National Parks (Campbell, et al., 2008). Between 2008 and 2011, the population in Lagoas de Cufadas National Park in Guinea-Bissau declined by more than 60% (Carvalho, et al., 2013).

#### Bonobo, *Pan paniscus*

Endemic to the DRC, the *Pan paniscus* population is estimated to have declined sharply in the last 15–20 years. This decline is projected to exceed 50% over a 75-year period from 2003 to 2078. The species is listed as Endangered. Source: Fruth, et al. (2016).

**Table 4.** African great ape population trends by taxon in descending order of abundance. *Except for mountain gorillas, estimates are based on the number of “nest builders”, thus excluding infants. Estimates are based on both surveys and modelled results.*

Taxon	Abundance	Trend	Annual rate of change	Total estimated change	Period assessed	Source of trend
<b>western lowland gorilla</b> <i>Gorilla g. gorilla</i>	361,919 (302,973–460,093)*	Declining ↓	–2.7%	–19.4%	2005–2013	Strindberg, et al. (2018)
<b>eastern chimpanzee</b> <i>Pan t. schweinfurthii</i>	181,000–256,000	Declining ↓	–5.1%	–22–45% †	1994–2014	Plumptre, et al. (2015; 2016b)
<b>central chimpanzee</b> <i>Pan t. troglodytes</i>	128,760 (114,208–317,039)*	Declining ↓	>–4.95%	>–50%	2005–2013	Maisels, et al. (2016b)
<b>western chimpanzee</b> <i>Pan t. verus</i>	18,000–65,000*	Declining ↓	–6.43%	–80%	1990–2015	Kühl, et al. (2017)
<b>bonobo</b> <i>Pan paniscus</i>	15,000–20,000 minimum	Declining ↓	–5.95% §	–54.9%	2003–2015	Fruth, et al. (2016)
			–1% ¶	>–50%	2003–2078	
<b>Nigeria-Cameroon chimpanzee</b> <i>Pan t. ellioti</i>	4,400–9,345	Declining ↓	–0.92–2.14%	–50–80%	1985–2060	Oates, et al. (2016)
<b>Grauer’s gorilla</b> <i>Gorilla b. graueri</i>	3,800 (1,280–9,050)	Declining ↓	–7.2%	–94.2%	1994–2015	Plumptre, et al. (2016a, 2016c)
<b>mountain gorilla</b> <i>Gorilla b. beringei</i>	>1,000	Increasing ↑	+3.7%	+26% ‡	2003–2010	Gray, et al. (2013); Roy, et al. (2014); Hickey, et al. (in prep.)
<b>Cross River gorilla</b> <i>Gorilla g. diehli</i>	<300	Declining ↓	N/A	N/A	N/A	Dunn, et al. (2014); R. Bergl & J. Oates pers. comm.

\* Surveys conducted in 2003–2013 in western equatorial Africa were used to estimate total population size.

† 22–45% decline estimated for eastern DRC only, not for the entire geographic range.

§ The confidence interval for this analysis is very large, suggesting an uncertainty in the data.

¶ There is uncertainty surrounding the 5.95% annual decline; however, Fruth, et al. (2016) state that an annual loss of 1% would still lead to >50% decline of the bonobo population by 2078.

‡ Virunga population only. No estimate available for the Bwindi population due to changes in sampling method.

#### 2.4. Threats to African great apes

Threats are discussed in detail in Section 4. The most important threats to great apes in Africa are poaching for bushmeat, habitat loss and degradation, and infectious diseases. The term “poaching” is used as a synonym for illegal killing, but can have different motives, such as obtaining bushmeat or retaliation for crop foraging. Great apes may also become accidental victims of snares set for other species (see poaching categories in Table 10 of Section 4). Illegal trafficking of

live infants is also an issue in some areas. The levels of these threats vary greatly between taxa (Table 5). Western lowland gorillas and central chimpanzees are treated together because their ranges overlap by 97% (Strindberg, et al., 2018).

#### **Grauer's gorilla, *Gorilla beringei graueri***

Even though all killing, capture and consumption of great apes is illegal in the DRC, bushmeat poaching presents the most serious and immediate threat to Grauer's gorillas. This concerns the entire geographic range with a high demand for bushmeat created by the growing human population and widespread artisanal mining in remote areas (Kirkby, et al., 2015; Plumtre, et al., 2015, 2016a). Miners working in Grauer's gorilla habitat admit to poaching gorillas, considering them to be relatively easy to hunt with guns and providing large quantities of meat (Kirkby, et al., 2015, Spira, et al., 2017). Indiscriminate and conflict poaching also occur. Habitat loss and degradation, mainly driven by artisanal mining, farming (slash-and-burn agriculture) and livestock ranching to supply regional markets, also constitutes a major threat. There is currently no commercial logging within the Grauer's range, although artisanal logging is widespread. As parts of DRC emerge from civil war, new concessions for timber, minerals and possibly petroleum will pose conservation challenges in the future. Another major threat to Grauer's gorillas is civil unrest in eastern DRC, which has massively exacerbated the decline of this subspecies since the mid-1990s. Long-term conflict and insecurity resulted in rebel and civilian occupation of the forests, including protected areas, and severely restricted the ability of conservation organizations to monitor and protect gorillas, enforce hunting and protected area regulations, and address intensifying threats. Only 26% of the predicted range of Grauer's gorillas overlaps with national parks and nature reserves. Source: Plumtre, et al. (2016c).

#### **Mountain gorilla, *Gorilla beringei beringei***

Mountain gorillas live in the Bwindi Impenetrable National Park in Uganda and the Virunga Massif, shared by DRC, Rwanda and Uganda. Mountain gorilla habitat is well protected and conservation measures, including long-term activities such as intensive law enforcement, research, tourism and veterinary care have been proven to lower the impact of the main threats, which are habitat loss, indiscriminate poaching, and mortality due to disease (Robbins, et al., 2011). As a result of these conservation activities, population numbers have increased. Despite these positive results, challenges remain. Most poaching is indiscriminate (snares). There are occasional conflict killings and, in the past, mountain gorillas have been shot dead in both politically-motivated incidents and in crossfire (collateral). The mountain gorilla population is growing, but the habitat is surrounded by farms, limiting options for expansion. Another issue is climate change, which is likely to result in changes in food availability and habitat quality for the mountain gorillas, as well as the surrounding local communities, which could increase the challenges to conservation efforts. Source: Plumtre, et al. (2016d).

#### **Cross River gorilla, *Gorilla gorilla diehli***

Cross River gorillas occur in 13 small forest fragments totalling approximately 600 km<sup>2</sup> in a landscape of 13,000 km<sup>2</sup>. These fragments are surrounded by densely populated human settlements. Many of the subspecies' subpopulations are outside of protected areas and are at most risk from hunting and habitat loss. Although poachers do not specifically target gorillas, it is estimated that opportunistic bushmeat poaching removes 1–3 individuals from the population annually (though this may be an underestimate; Dunn, et al., 2014). Cross River gorillas are also occasionally killed or injured in snares (indiscriminate poaching). Their habitat is also under threat. Much of the suitable habitat in Cameroon has no protected status and there is rapid, ongoing conversion of forest to agriculture and grazing. There is also some habitat loss even inside the protected areas, and corridors between subpopulations are particularly vulnerable. For example, the Okwangwo Division of Nigeria's Cross River National Park and the adjacent Takamanda National Park in Cameroon contain enclaves of human settlements whose farmlands have spread beyond their legal boundaries. The small size of the Cross River population in general and its high level of fragmentation also make it more vulnerable to disease. The Ebola virus, which has caused significant mortality in *Gorilla g. gorilla* populations, has not been reported in *Gorilla g. diehli* populations. Nevertheless, their proximity to dense human populations and livestock heightens the risk of disease transmission (Dunn, et al., 2014). Source: Bergl, et al. (2016).

#### **Western lowland gorilla, *Gorilla gorilla gorilla* and central chimpanzee, *Pan troglodytes troglodytes***

Poaching for bushmeat is the primary driver of decline in western lowland gorilla and central chimpanzee populations. Most of the terra firma forests outside the protected areas of their geographic range are now logging concessions (Global Forest Watch, 2017). A network of new logging roads provides rapid access to poachers and traffickers into previously inaccessible forests. Consignments of bushmeat can be rapidly sent, according to estimations, hundreds of kilometres out of the forests (Maisels, et al., 2016a). Infectious disease, especially the Ebola virus, is the second major driver of their decline. Surveys carried out since the 1980s show that a series of large great ape die-offs have occurred in a large forest region that straddles the border between northeastern Gabon and northwestern Congo. Approximately 14% of the total range of these taxa is thought to have been affected by Ebola virus disease. At present, habitat loss (as opposed to habitat degradation) in the region is low, but this will change in the near future: 42% of the western lowland gorilla and central chimpanzee geographic range is considered suitable for oil palm, the progressive development of which could become a major threat to these taxa (Wich, et al., 2014). Approximately 80% of these two great ape subspecies live outside formally protected areas (Strindberg, et al., 2018), making them and their habitat vulnerable to habitat loss and bushmeat poaching. Sources: Maisels, et al. (2016a, 2016b).

### **Eastern chimpanzee, *Pan troglodytes schweinfurthii***

Bushmeat poaching is the greatest threat to great apes in eastern DRC, where large populations of eastern chimpanzees occur. Eastern chimpanzees are poached for bushmeat, especially around artisanal mining and logging camps, where bushmeat is often the main source of protein available. When adult chimpanzees are killed for bushmeat, their infants may end up in the ape trade as pets (Hicks, et al., 2010). Bushmeat poaching is evidenced by ongoing confiscations, as well as the hundreds of eastern chimpanzees housed in sanctuaries in the DRC (Plumptre, et al., 2015). In eastern DRC, armed groups involved in artisanal mining are responsible for much of the bushmeat poaching. In both eastern and northern DRC, poaching and illegal wildlife trafficking have been driven by insecurity over the past 20 years (Ondoua Ondoua, et al., 2017). Another major threat is habitat loss and degradation due to smallholder and shifting agriculture. Industrial agriculture (e.g. oil palm plantations) poses a potential future threat as eastern DRC stabilizes (Plumptre, et al., 2016b). Infectious diseases were identified as the major cause of death in chimpanzees at Gombe and Mahale in Tanzania (e.g. Goodall, 1986; Nishida, et al., 2003). The frequency of encounters between chimpanzees, humans and human waste is increasing as human populations expand, leading to higher risks of disease transmission. Source: Plumptre, et al. (2016b).

### **Nigeria-Cameroon chimpanzee, *Pan troglodytes ellioti***

The two main threats to the survival of Nigeria-Cameroon chimpanzees are human hunting (bushmeat poaching) and habitat loss (Morgan, et al., 2011). These threats are exacerbated by the expansion of human populations in the subspecies' geographic range, as well as economic growth in Cameroon and Nigeria. Poaching represents the greatest threat to the survival of this taxon, both supplying bushmeat trade and, to a lesser extent, providing body parts for traditional medicine (superstition poaching) (Oates, et al., 2016). In 2002–2003, a six-month study in rural markets in southeastern Nigeria and southwestern Cameroon recorded 240 chimpanzee carcasses (Fa, et al., 2006). Furthermore, a five-week survey of eight markets in the transboundary region of Cross River State in Nigeria recorded six chimpanzee carcasses in 2009 (Oates, et al., 2016). Suitable habitat in Nigeria and Cameroon continues to be lost, degraded and fragmented by agriculture, logging, grazing and fire. In eastern Nigeria in particular, several forest reserves have been converted to farmland and to commercial oil palm and rubber plantations, while large areas of forest surrounding key protected areas, such as Okomu National Park, have already been converted to oil palm plantations. In Cameroon, extensive new oil palm developments are underway in both Littoral and Southwest Regions, and new logging concessions continue to be established. The combined impacts of habitat loss and poaching have gradually fragmented *Pan t. ellioti* populations, so that many of those remaining are small and isolated. They are therefore at increased risk of extinction from disease and other unpredictable events. Source: Oates, et al. (2016).

### **Western chimpanzee, *Pan troglodytes verus***

Half of western chimpanzees reside in the Fouta Djallon region of Guinea (Regnaut & Boesch, 2012). This area is characterized by traditional small-scale farming practices, predominantly by Fulani people, who neither eat nor kill chimpanzees for cultural reasons (Ham, 1998). The Fouta Djallon also contains the world's largest bauxite deposits and it is likely that industrial mining will occur in much of the chimpanzee habitat within the next decade (Kormos, et al., 2014). If mining proceeds at the scale planned, it will most certainly cause further population declines and thus threaten this stronghold of the subspecies (Kühl, et al., 2017). Mines also need electricity for their operations, and there are plans for hydrodams throughout the Fouta Djallon region, which would accelerate habitat loss (R. Kormos, pers. comm.). Other causes of western chimpanzee decline include several types of poaching (bushmeat, indiscriminate, conflict, superstition), capture (and associated poaching and illegal trafficking), habitat loss, and infectious diseases (Humble, et al., 2016). These threats and their underlying drivers have caused local extirpation of chimpanzee populations, especially in Ghana and Côte d'Ivoire. Population declines in Côte d'Ivoire have also been caused by large-scale deforestation inside and outside of protected areas and classified forests. Such deforestation is a result of rapid growth of the human population, massive immigration from the Sahel Belt and industrial-scale agricultural production of coffee, cacao, rubber and palm oil (Campbell, et al., 2008; Kühl, et al., 2017). The remaining strongholds of western chimpanzees are Guinea, Liberia and Sierra Leone. Most chimpanzees in Liberia and Sierra Leone, however, occur outside protected areas, where subsistence agriculture is a major driver of forest loss, and where coexistence with people is strained because chimpanzees consume crops and compete over wild resources, such as the oil palm. Chimpanzees in Liberia are threatened by bushmeat poaching both inside and outside protected areas (Greengrass, 2016), as well as by the rapidly developing mining, forestry and industrial-agricultural sectors (Junker, et al., 2015; Tweh, et al., 2015). Over 80% of the western chimpanzee's geographic range in Liberia and Sierra Leone is outside protected areas and threatened by industrialised agriculture and oil palm developments (Wich, et al., 2014) and associated infrastructure, such as roads and hydrodams.

### **Bonobo, *Pan paniscus***

The most significant threat to bonobos is bushmeat poaching, followed by habitat loss through deforestation and fragmentation. Disease is likely to pose a threat in the future, with increased exposure to human populations. A number of indirect threats exist, including the proliferation of weapons in the region, weak law enforcement, weak stakeholder commitment to conservation, expansion of slash-and-burn agriculture and industrial-scale commercial activities. Not only is there a massive demand for bushmeat stemming from the cities, but rebel factions and poorly-paid government soldiers add to that demand. Source: Fruth, et al. (2016).

**Table 5.** Major threats affecting African great apes at taxon level. *Some direct threats have a larger impact on great ape populations than others, but no quantitative comparisons are possible.*

<b>Taxon</b>	<b>Main threats (direct)</b>	<b>Source</b>
<b>Grauer's gorilla</b> <i>Gorilla b. graueri</i>	<b>Poaching</b> (types: bushmeat, indiscriminate, conflict, superstition, collateral). N.B. Illegal trafficking of live orphans is a by-product of bushmeat trade	Plumptre, et al. (2015; 2016c)
	<b>Habitat loss, fragmentation and degradation</b> due to artisanal mining, shifting and commercial agriculture	
	<b>Disease</b>	
	<b>Climate change</b>	
<b>mountain gorilla</b> <i>Gorilla b. beringei</i>	<b>Poaching</b> (types: indiscriminate, conflict, politically-motivated)	Gray, et al. (2010); Robbins, et al. (2011); Roy, et al. (2014)
	<b>Disease</b>	
	<b>Climate change</b>	
<b>Cross River gorilla</b> <i>Gorilla g. diehli</i>	<b>Habitat loss, fragmentation and degradation</b> due to shifting and commercial agriculture	Bergl, et al. (2016)
	<b>Poaching</b> (types: bushmeat, indiscriminate, conflict)	
<b>western lowland gorilla</b> <i>Gorilla g. gorilla</i>	<b>Poaching</b> (types: bushmeat, indiscriminate, conflict). N.B. Illegal trafficking of live orphans is a by-product of bushmeat trade	Maisels, et al. (2016a)
	<b>Disease</b>	
	<b>Habitat loss and degradation and degradation</b> due to extractive industries, commercial agriculture and infrastructure development	
<b>western chimpanzee</b> <i>Pan t. verus</i>	<b>Habitat loss, fragmentation and degradation</b> due to shifting and commercial agriculture, extractive industries and infrastructure development	Humle, et al. (2016); Kühl, et al. (2017)
	<b>Poaching</b> (types: bushmeat, indiscriminate, conflict, superstition, live capture)	
	<b>Illegal trafficking of live animals</b>	
	<b>Disease</b>	
<b>Nigeria-Cameroon chimpanzee</b> <i>Pan t. ellioti</i>	<b>Poaching</b> (types: bushmeat, indiscriminate, conflict)	Oates, et al. (2016)
	<b>Habitat loss fragmentation and degradation</b> due to shifting and commercial agriculture	
	<b>Disease</b>	
<b>central chimpanzee</b> <i>Pan t. troglodytes</i>	<b>Poaching</b> (types: bushmeat, indiscriminate). N.B. Illegal trafficking of live orphans is a by-product of bushmeat trade	Maisels, et al. (2016b)
	<b>Disease</b>	

	<b>Habitat loss, fragmentation and degradation</b> due to extractive industries, commercial agriculture and infrastructure development	
<b>eastern chimpanzee</b> <i>Pan t. schweinfurthii</i>	<b>Poaching</b> (types: bushmeat, indiscriminate, conflict, superstition). N.B. Illegal trafficking of live orphans is a by-product of bushmeat trade	Plumptre, et al. (2015; 2016b)
	<b>Habitat loss, fragmentation and degradation</b> due to shifting and commercial agriculture, artisanal and industrial mining, and infrastructure development	
	<b>Disease</b>	
	<b>Climate change</b>	
<b>bonobo</b> <i>Pan paniscus</i>	<b>Poaching</b> (types: bushmeat, indiscriminate, superstition). N.B. Illegal trafficking of live orphans is a by-product of bushmeat trade	Fruth, et al. (2016)
	<b>Habitat loss and degradation</b> from shifting agriculture, mining and infrastructure development	
	<b>Disease</b>	Sakamaki, et al. (2009); IUCN & ICCN (2012)

### 3. Asian great apes


#### 3.1. Current distribution of Asian great apes

Orangutans are the only great apes found in Asia and they exist solely on the islands of Sumatra and Borneo (Figure 2), in Indonesia and Malaysia (Wich, et al., 2008). There are three species: the Bornean orangutan (*Pongo pygmaeus*), the Sumatran orangutan (*Pongo abelii*), and the recently described Tapanuli orangutan (*Pongo tapanuliensis*). Bornean orangutans are further divided into three subspecies: *Pongo pygmaeus pygmaeus*, *Pongo pygmaeus wurmbii* and *Pongo pygmaeus morio* (Table 6).

**Table 6.** The Asian great apes

Genus	Species	Subspecies
<b>Pongo</b> 3 species	<b>Bornean orangutan</b> <i>Pongo pygmaeus</i> 3 subspecies	<b>northwest Bornean orangutan</b> <i>Pongo pygmaeus pygmaeus</i>
		<b>southwest Bornean orangutan</b> <i>Pongo pygmaeus wurmbii</i>
		<b>northeast Bornean orangutan</b> <i>Pongo pygmaeus morio</i>
	<b>Sumatran orangutan</b> <i>Pongo abelii</i>	<b>No subspecies currently recognized</b>
	<b>Tapanuli orangutan</b> <i>Pongo tapanuliensis</i> *	<b>No subspecies currently recognized</b>

\* New species described by Nater, et al. (2017)



**Figure 2.** Geographic distribution of Asian great ape taxa (orangutans; IUCN SSC A.P.E.S. database, 2017), Max Planck Institute for Evolutionary Anthropology.

### 3.2. Asian great ape population sizes

#### 3.2.1. Population size estimates per country

**Table 7.** All Asian great ape population estimates per country†

Country	Taxon	Abundance*	IUCN status	Year of estimate	Source
Indonesia	<b>northwest Bornean orangutan</b> <i>Pongo p. pygmaeus</i>	5,200 (3,800–7,200)	Critically Endangered	2018	Calculated from Voigt, et al. (2018)
	<b>southwest Bornean orangutan</b> <i>Pongo p. wurmbii</i>	97,000 (73,800–135,000)	Critically Endangered	2018	Calculated from Voigt, et al. (2018)
	<b>northeast Bornean orangutan</b> <i>Pongo p. morio</i>	24,800 (18,100–35,600)	Critically Endangered	2018	Calculated from Voigt, et al. (2018)
	<b>Sumatran orangutan</b> <i>Pongo abelii</i>	13,900 (5,400–26,100)	Critically Endangered	2016	Wich, et al. (2016)
	<b>Tapanuli orangutan</b> <i>Pongo tapanuliensis</i>	800 (300–1,400)	Critically Endangered	2016	Wich, et al. (2016)
Malaysia	<b>southwest Bornean orangutan</b> <i>Pongo p. wurmbii</i>	1,100 (800–1,600)	Critically Endangered	2018	Calculated from Voigt, et al. (2018)
	<b>northeast Bornean orangutan</b>	11,000 (8,000–18,000)	Critically Endangered	2005	Ancrenaz, et al. (2005)

	<i>Pongo p. morio</i>				
--	-----------------------	--	--	--	--

† A PHVA workshop took place in Indonesia in 2016. The report awaits final approval.

\* Numbers rounded to the nearest 100. 95% confidence intervals in parentheses.

### 3.2.2. Population size estimates per taxon

#### Bornean orangutan, *Pongo pygmaeus*

Abundance estimates for *P. pygmaeus pygmaeus*, *P. pygmaeus wurmbii* and *P. pygmaeus morio*, all listed as Critically Endangered, are presented in Table 8.

#### Sumatran orangutan, *Pongo abelii*

Previously estimated to number 6,600 (Wich, et al., 2008), the most recent abundance estimate for the Sumatran orangutan is 13,900 individuals, in a total area of 16,775 km<sup>2</sup> of forest (Wich, et al., 2016). This latest estimate does not reflect an increase in Sumatran orangutan numbers, but it is explained by much improved survey techniques and coverage, and hence more accurate data. Considering only populations that are potentially viable over the long term (i.e. > 250 individuals in each), there are effectively just 13,587 individuals remaining. The vast majority (95%) occurs in the Leuser ecosystem, while other populations are found in the Sidiangkat, Pakpak and Batang Toru forests (Singleton, et al., 2017). The overall number continues to decline dramatically (Wich, et al., 2016). The species is classified as Critically Endangered.

#### Tapanuli orangutan, *Pongo tapanuliensis*

This new species was first described in 2017 (Nater, et al., 2017). The Nater, et al. (2017) study showed that an isolated orangutan population found in the Batang Toru ecosystem of Sumatra, in the southernmost range of extant Sumatran orangutans, is distinct from other Sumatran and Bornean populations. With a total estimated population of fewer than 800 individuals (Wich, et al., 2016), *Pongo tapanuliensis* is the great ape species with the lowest total number of individuals, and is Critically Endangered.

### 3.3. Temporal trends in Asian great ape populations

**Table 8.** Asian great ape population decline by taxon

Species	Subspecies	Abundance	Trend	Annual rate of change	Total estimated change	Period	Source of trend
Bornean orangutan	northwest Bornean orangutan <i>Pongo p. pygmaeus</i>	6,300 (4,700–8,600)	Declining ↓	–4.71%	–53%	1999–2015	Calculated from Voigt, et al. (2018)
	southwest Bornean orangutan <i>Pongo p. wurmbii</i>	97,000 (73,800–135,000)	Declining ↓	–4.71%	–53%	1999–2015	Calculated from Voigt, et al. (2018)
	northeast Bornean orangutan <i>Pongo p. morio</i>	30,900 (22,800–44,200)	Declining ↓	–4.45%	–52%	1999–2015	Calculated from Voigt, et al. (2018)
Sumatran orangutan	Sumatran orangutan <i>Pongo p. abelii</i> *	13,900 (5,400–26,100)	Declining ↓	–2.37%	–30%*	2015–2030	Wich, et al. (2016)
Tapanuli orangutan	Tapanuli orangutan <i>Pongo tapanuliensis</i>	800 (300–1,400)	Declining ↓	–2.36% (for 1985–2060)	–83%	1985–2060	Nowak, et al. (2017)

\*See temporal trends text for the Sumatran orangutan based on forest loss. Confidence intervals rounded to the nearest 100, 95% confidence intervals are in parentheses. Under the current land-use scenario, as many as 4,500 individuals could disappear by 2030. Other scenarios are mentioned in Wich, et al. (2016).

#### Bornean orangutan, *Pongo pygmaeus*; subspecies: *Pongo p. pygmaeus*, *Pongo p. wurmbii*, *Pongo p. morio*

Recent studies on the temporal trends of Bornean orangutans found declines of 25% over a 10-year period (2000–2010; Santika, et al., 2017) and approximately 50% for the period 1999–2015 (Voigt, et al., 2018). Based on predicted land-

cover change alone, a further 43,000 Bornean orangutans could be lost by 2050, which is equivalent to 50–60% of the current estimated population (Voigt, et al., 2018). Climate change and land-cover prediction models further show that, under a business as usual scenario, 68–81% of the Bornean orangutan habitat will be lost by 2080 (Wich, et al., 2015).

#### **Sumatran orangutan, *Pongo abelii***

The decline of the Sumatran orangutan to its current estimated population of 13,900 in 2016 (Wich, et al., 2016) has not been systematically quantified. However, forest loss data indicate that the decline must have been large. Between 1985 and 2007, the Sumatran orangutan lost 60% of its key forest habitat (<500 m). Future predictions of forest loss indicate that a further 4,500 Sumatran orangutans, about one-third of the current population, could be lost by 2030 (Wich, et al., 2016).

#### **Tapanuli orangutan, *Pongo tapanuliensis***

A quantitative population viability analysis estimated that, in 1985, there would have been ~1,489 individuals of the Tapanuli orangutan, and that the population would decline to only 257 individuals over a 75-year period by 2060 (Nowak, et al., 2017). If the key threats are not effectively reduced, an 83% decline over the course of three generations is predicted. Source: Nowak, et al. (2017).

### *3.4. Threats to Asian great apes*

#### **Bornean orangutan, *Pongo pygmaeus*; subspecies: *Pongo p. pygmaeus*, *Pongo p. wurmbii*, *Pongo p. morio***

Widespread forest clearance for industrial plantations, cultivation for food, mining infrastructure and rural development, combined with illegal logging, fire and several types of poaching (bushmeat, conflict and live capture), are the main threats to this taxon and have dramatically reduced the number of Bornean orangutans (Wich, et al., 2008; Wich, et al., 2012b; Santika, et al., 2017; Voigt, et al., 2018). Data from the Global Forest Resources Assessment shows a 2.4% forest loss between 2000 and 2015 and a 0.7% loss between 2010 and 2015 for the whole of Borneo (FAO, 2015). This translates to an annual rate of deforestation estimated at more than 3,000 km<sup>2</sup> per year between 2000 and 2010 (Gaveau, et al., 2014). If this deforestation rate continues, it is predicted that 32,000 km<sup>2</sup> of forest could be lost by 2020, 129,000 km<sup>2</sup> by 2050 and 226,000 km<sup>2</sup> by 2080 (Wich, et al., 2015). Most of this deforestation occurs in peatlands that generally harbour large orangutan populations. In 2010, 80% of the range in Kalimantan was located outside protected areas, consisting of commercial forest reserves exploited for timber and forest areas earmarked for conversion to agriculture. The situation is different in Malaysia, where currently more than 80% of orangutans are found in fully-protected forests. Besides forest loss, poaching is another major cause of Bornean orangutan decline (see poaching categories in Table 10 of Section 4). On average, an estimated 1,950–3,100 orangutans were killed per year within the lifetime of survey respondents (Meijaard, et al., 2011), principally for meat consumption (bushmeat poaching) or during human-orangutan conflict (conflict poaching) (Davis, et al., 2013; Voigt, et al., 2018). This means that habitat protection alone will not ensure the survival of orangutans and that effective reduction of orangutan killings is urgently needed (Ancrenaz, et al., 2016). Other threats are fires that contribute to habitat loss and fragmentation, lack of environmental awareness and climate change (Ancrenaz, et al., 2016; Santika, et al., 2017; Voigt, et al., 2018).

#### **Sumatran orangutan, *Pongo abelii***

Habitat loss and fragmentation seriously threaten the survival of the Sumatran orangutan (Wich, et al., 2011; Wich, et al., 2016). Forests continue to be cleared on a large scale (hundreds of square kilometres) for oil palm plantations. On a smaller scale, logging for timber (both legal and illegal) remains a threat. In addition, the creation of new roads fragment populations and provide access for illegal settlement and further encroachment for agriculture and plantations (also frequently illegal) and to wildlife poachers (Singleton, et al., 2017). Sumatran orangutans are often killed deliberately during human-wildlife conflict (conflict poaching), and surviving infants end up in the illegal pet trade (Nijman, 2009; Singleton, et al., 2017). A significant threat to the Sumatran orangutan comes from the 2013 Aceh province spatial land-use plan allowing large areas of the Leuser ecosystem, which hosts 90% of Sumatran orangutans, to be designated for oil palm plantations as well as timber and mining concessions (Wich, et al., 2016). With the same Leuser ecosystem designated in 1997/98 as a National Strategic Area for its environmental function under Aceh's special autonomy law, and it being the main stronghold of the Sumatran orangutan, revising the spatial land-use plan as a priority would have meaningful benefits for future orangutan populations.

#### **Tapanuli orangutan, *Pongo tapanuliensis***

The Tapanuli orangutan is under considerable threat from high levels of habitat loss and fragmentation, as well as from bushmeat poaching, killing during conflicts over crops (conflict poaching), and illegal trade in young orangutans, fuelled by a human population influx from the west of Sumatra. A substantial section of the range is threatened by habitat conversion for small-scale agriculture, mining exploration and exploitation, a large-scale hydroelectric scheme and geothermal development (Nowak, et al., 2017). Due to their slow reproduction rate, with a generation time of about 25 years (Wich, et al., 2004; Wich, et al., 2009), orangutans on Sumatra are unable to cope with significant and continued individual losses. The small population size and geographic isolation of *Pongo tapanuliensis* may lead to inbreeding depression (Hedrick & Kalinowski, 2000) and threaten population persistence (Allendorf, et al., 2013).

**Table 9.** Summary of threats facing Asian great apes per taxon

Species	Main threats	Source
<b>Bornean orangutan</b> <i>Pongo pygmaeus</i>	<b>Habitat loss, fragmentation and degradation</b> due to agriculture, extractive industries and fire	Ancrenaz, et al. (2016); Voigt, et al. (2018)
	<b>Poaching</b> (types: bushmeat, conflict)	
	<b>Climate change</b>	Struebig, et al. (2015)
<b>Sumatran orangutan</b> <i>Pongo abelii</i>	<b>Habitat loss, fragmentation and degradation</b> due to agriculture, extractive industries, and infrastructure (roads)*	Wich, et al. (2012a, 2016); Singleton, et al. (2017)
	<b>Poaching</b> (type: conflict)	Wich, et al. (2012a); Singleton, et al. (2017)
<b>Tapanuli orangutan</b> <i>Pongo tapanuliensis</i>	<b>Habitat loss, fragmentation and degradation</b> due to agriculture, extractive industries and construction of large-scale infrastructure (e.g. hydroelectric dam)	Nowak, et al. (2017)
	<b>Poaching</b> (types: bushmeat, conflict)	Wich, et al. (2012a)

\*Cases of illegal trade as a by-product of habitat loss have been reported (Singleton, et al., 2017), but in these cases trade is not a direct threat; the direct threat is habitat loss.

#### 4. Description of threats

The most important direct threats to great apes are habitat loss, degradation and fragmentation, infectious disease, bushmeat poaching, indiscriminate poaching, and deliberate killing due to conflicts over resources, usually cultivated foods. The relative importance of these threats varies by taxon and location. Areas with high human population densities tend to have the most degraded habitats and the lowest ape densities. Many range States are among the poorest countries in the world, with growing populations and development aspirations, including DRC, Guinea, Liberia, Rwanda and Uganda. Sub-Saharan African human population growth rates are among the highest in the world, at about 2.7% annually (World Bank, 2018). This places ever-increasing pressure on land, natural resources and wildlife. Consequently, a thorough understanding of local circumstances is required to address the main threats to great apes, as they cannot be considered in isolation from each other.

##### *Habitat loss, degradation and fragmentation*

Great apes are dependent on forest ecosystems, but these are increasingly threatened by industrial and small-scale agriculture, resource extraction such as logging and mining, as well as infrastructure development (e.g. Ancrenaz, et al., 2015). Mirroring the major role that oil palm has played in the loss of great ape habitats in Asia, the progressive development of plantations is a future threat to the African great apes (Wich, et al., 2014). Once a forest habitat is lost, it is very difficult to restore and repopulate it. Laurance, et al. (2015) describe the environmental costs of 33 planned or under-construction development corridors in Sub-Saharan Africa. It is important to note that some great ape taxa can survive in reduced-impact logging areas (gorillas and chimpanzees: Morgan, et al., 2017; orangutans: Ancrenaz, et al., 2010). However, survival in low impact logging areas is only possible when the logging is carefully controlled, does not lead to subsequent uncontrolled logging and further habitat deterioration, and when bushmeat poaching is not a threat (Morgan & Sanz, 2007; Ancrenaz, et al., 2010). Any progressive loss of forest resources (as opposed to rapid loss or habitat conversion) also necessitates high levels of tolerance from people in surrounding areas if great apes are to persist, especially if great apes turn to foraging on crops to supplement their diet, which is the case for orangutans in Asia and many chimpanzee populations in West and East Africa (Hockings & Humle, 2009; Campbell-Smith, et al., 2012; Bryson-Morrison, et al., 2016; Garriga, et al., 2017).

##### *Disease*

Great apes and humans are very closely related and are susceptible to the same deadly diseases (Gilardi, et al., 2015). Such diseases include Ebola, anthrax, Marburg viruses and respiratory diseases, with transmission risks growing as human populations expand further into great ape habitat (Leendertz, et al., 2006). Pathogen transfer goes both ways and bushmeat consumption can lead to infections in humans (Leendertz, et al., 2017). The spread of disease is facilitated by habitat fragmentation, close proximity between great ape populations and human settlements, and high frequencies of human-great ape interactions (Gilardi, et al., 2015).

## Poaching

The term “poaching” is used as a synonym for illegal killing, but can have different purposes (see Table 10). We categorize (but do not attempt to quantify) the key types of poaching according to motivation and/or the context in which killing takes place.

**Table 10.** Key categories of poaching (illegal killing)

<b>Type 1: Great apes are targeted and killing is intentional</b>		
a	Bushmeat	Most killing in this context is opportunistic, i.e. great apes are killed while hunters are tracking other species, such as antelopes or monkeys. However, in some range States, great apes are occasionally killed deliberately to serve their meat as a delicacy or high-status food at traditional or political functions. N.B. If infants are captured alive when their mothers are killed for bushmeat, these orphans may be illegally trafficked. This trade is a by-product of poaching for bushmeat, so is classed as bushmeat poaching (see also 1d below).
b	Conflict	Great apes are killed over competition for natural resources (see Hockings & Humle, 2009). Retribution killings motivated by crop-foraging or injuries to people are in this category. Illegal killing and capture may occur on a large-scale when land is cleared or converted to other land uses.
c	Superstition	Great apes are sometimes killed for traditional ceremonies and “medicine” – superstitious beliefs, including black magic. N.B. Body parts from a dead ape that are used for superstitious purposes when the primary motivation for killing was to obtain meat (1a above) are secondary uses and therefore classed as bushmeat poaching; e.g. bones or digits used to confer “strength” upon hunters or human infants.
d	Live	Live capture and associated killing covers taking live infant great apes as the primary motivation (as opposed to the primary motivation being to obtain bushmeat). Live capture is in itself illegal, but obtaining an infant necessitates killing the mother and usually other group or community members. N.B. The majority of orphaned infants thought to be destined for the illegal wildlife trade were not targeted by poachers with the intention of trading them – this trafficking is a “by-product” of bushmeat poaching (1a above).
e	Politically-motivated	Killing of great apes can result from conflict with government, usually the protected area authorities, or specifically with law enforcement agents.

<b>Type 2: Great apes are not targeted or killing is not intentional</b>		
a	Indiscriminate	Great apes are accidentally caught in snares intended for other species. Often the apes are maimed and lose a hand or foot. These incidents are fatal when the animals are unable to free or feed themselves, or when they develop gangrene or septicemia.
b	Collateral	Great apes have been shot in civil conflicts – killed in crossfire. Even when the killing is accidental, the ape is usually eaten. N.B. Civil war and insecurity exacerbate illegal activities, including bushmeat poaching (1a above).

CITES, INTERPOL and World Customs Organization (WCO) databases (SC65, Doc. 37) only record international trade, thereby under-representing domestic trade. This highlights the need for domestic monitoring.

### A) Domestic trade

Poaching is a key threat to all great ape taxa (Tables 5 and 9). Almost all wildlife species (mammals, birds, reptiles, amphibians and even insects) are eaten in much of West and Central Africa, and a huge bushmeat trade network has developed, where very large numbers of animals are hunted in remote forest areas and brought to industrial camps (logging, mining), towns and cities for profit, often over very long distances (e.g. Nasi, et al., 2008; Fa & Brown, 2009; Wilkie, et al., 2011; Ziegler, 2016). For example, animals are hunted in Salonga National Park in central DRC and their meat is smoked and then transported on foot and by bicycle, hundreds of kilometres south to the mining towns in Katanga, where the purchasing power of consumers is far greater than in the source villages (e.g. Colom, 2006; Steel, et al., 2008; Abernethy, et al., 2010).

Great ape populations in West and Central Africa are highly threatened by the commercial bushmeat trade (e.g. Refisch & Koné, 2005; Plumptre, et al., 2010, 2015; Fruth, et al., 2016; Maisels, et al., 2016c). New roads into remote forest areas facilitate expansion of the bushmeat trade. For example, in Congo, the road from Brazzaville to Ouessou opens access to remote landscapes and enables transport of bushmeat. Illegal and unsustainable killing has brought many species to the brink of extinction, causing the “empty forest” syndrome (Redford, 1992; Wilkie, et al., 2011). Although great apes are

killed for their meat, they are not generally specifically targeted. There is evidence of deliberate hunting of Cross River gorillas for bushmeat in Cameroon (Dunn, et al., 2014). The killing of a mother may enable opportunistic capture of infants, which may then be trafficked illegally (e.g. Plumptre, et al., 2016b, 2016c; Singleton, et al., 2017). Some communities, particularly in Africa, consider great apes to be their close relatives and hence the killing of them is a taboo (e.g. in some parts of West Africa; Kormos, et al., 2003); or along the Congolese coast (Dowsett & Dowsett-Lemaire, 1991). However, in Central Africa, taboos against eating western lowland gorillas and central chimpanzees only exist in approximately 1% and 5% of their geographic ranges, respectively (Strindberg, et al., 2018). African apes are also killed for traditional medicine or witchcraft. For example, Sá, et al. (2012) suggest that poaching for superstitious reasons threatens chimpanzee populations in Guinea-Bissau.

Poaching of orangutans for food occurs extensively in Borneo, especially in the Indonesian parts of the island (Davis, et al., 2013), and to a lesser extent in the range of the Tapanuli orangutan on Sumatra (Wich, et al., 2012b). It is not known how much trade in orangutan meat occurs, but orangutans have been killed to sell their meat to local communities (Davis, et al., 2013). Sumatran and Tapanuli orangutans are similarly at risk from bushmeat poaching and killing as a result of habitat destruction or human-wildlife conflict (Nowak, et al., 2017; Singleton, et al., 2017).

Although funding has gone into providing alternative sources of protein, the impact of such investments has been marginal and, indeed, rarely quantified. Wicander & Coad (2015) investigated 64 projects in Central Africa that aimed to encourage the production of alternative protein sources and, in some cases, to increase revenue to the target communities. Unfortunately, very few of these projects monitored the impact of their efforts, so no conclusions could be reached as to whether they succeeded in their aims. Possibly because there were, in most cases, either no mechanisms for compliance, or no sanctions for noncompliance, or both, many participants were likely to adopt new protein sources as an additional activity, rather than an alternative activity. In other words, new activities were added to poaching and bushmeat trade rather than replacing them. In some cases, commercial hunters from outside the communities posed a far higher threat to wildlife or ecosystems than did the local communities themselves, rendering project activities ineffective in the face of external pressure. It is essential to build in and budget for the monitoring of project impacts, or it will be impossible to gauge success and failure in order to improve the effectiveness of future approaches.


#### *B) International trade*

Poaching for bushmeat poses particular challenges to enforcement, with species-specific identification of bushmeat being a key to addressing this threat. Numerous publications have investigated bushmeat issues in Africa and Asia (for examples see: Nasi, et al., 2008; Fa & Brown, 2009; Meijaard, et al., 2011; Foerster, et al., 2012; Coad, et al., 2013; Wilkie, et al., 2016; Ziegler, 2016). However, the volume of bushmeat that crosses international borders – which is specifically relevant to CITES – is particularly difficult to estimate. The vast majority of transborder trade in great ape meat is not transcontinental, but rather across neighbouring country boundaries, where detection is weak. Smuggled goods, including great ape meat, can easily be transported in small boats across rivers, which form highly porous international boundaries. Across terrestrial borders, trafficked goods often travel undiscovered in bush taxis, minibuses and on motorcycles, or even on foot through unpatrolled forests. The vast majority of gorillas, for example, live in Gabon and Congo and their meat is trafficked mostly within those two countries and across borders to the cities of neighbouring countries. Kinshasa in DRC, Yaoundé and Douala in Cameroon, and Bata in Equatorial Guinea are major destinations for all types of bushmeat, including that of great apes, which is trafficked from the more wildlife-rich parts of the region. Intercontinental illegal trade does occur, but is completely dwarfed by the subregional illegal trade. Only two studies at airports in Europe have been carried out, but these indicate that there is an active international bushmeat trade from West and Central Africa to Paris, Geneva and Zurich, which could be transit points to other European countries (Chaber, et al., 2010; Wood, et al., 2014). However, more research is required to determine how much great ape meat is involved in this trade (Chaber, et al., 2010).

#### *Illegal live trade*

The illegal trade in live great apes has been recognized as a threat to their survival since the 1980s (e.g. SC66 Doc. 48.2); however, the current scale of this trafficking is unknown. Indications of ongoing live trade include records from the United Nations Office on Drugs and Crime (UNODC) of 208 live apes seized since 2000 (UNODC, 2016). In 2017 alone, 20 great ape traffickers were arrested, and 12 live chimpanzees were seized in three African great ape range States, Cameroon, Gabon and Guinea (EAGLE Network, 2017). In most cases of illegal live ape capture and trafficking, adults must be killed to obtain infants (see, for example, Plumptre, et al., 2015; Humle, et al., 2016; Singleton, et al., 2017). Evidence indicates that illegal trade in live great apes is a secondary effect of habitat loss and poaching or conflict-related killing. Great apes are often captured and traded opportunistically, as opposed to poachers entering the forest to deliberately capture infants (e.g., Nijman, 2009; Davis, et al., 2013; Plumptre, et al., 2016b, 2016c; Humle, et al., 2016; Singleton, et al., 2017).

Known instances of great apes being targeted for capture and subsequent trade include Guinea, where highly organised syndicates have used fraudulent CITES papers to trade chimpanzees with China (CITES mission to Guinea in 2011).



**Figure 3.** Main international routes for the illegal trade of live great apes (Stiles, et al., 2013).

#### *Combined impacts of economic development*

Many great ape range States are rich in natural resources, for example, Guinea, DRC and Indonesia. The expanding extractives sector is one of the key drivers collectively accelerating the threats faced by great apes. This poses new challenges for great ape conservation, due to the difficulties in reconciling conservation priorities with urgently needed economic development. Alongside the direct impact of habitat loss caused by mines, logging concessions, roads and other infrastructure (e.g., transmission lines or dams), new development attracts huge numbers of people from far away in search of new opportunities (Rainer, et al., 2014). This often leads to uncontrolled additional development, including artisanal mining and farming, and the associated threat of bushmeat poaching. For example, Poulsen and Clark (2010) described poaching for bushmeat inside logging concessions, and Spira, et al. (2017) discuss the increase in bushmeat poaching of gorillas and chimpanzees around mines.

### **5. International agreements specific to great ape range States**

The Agreement on the Conservation of Gorillas and Their Habitats (or Gorilla Agreement) was developed under the auspices of Convention on the Conservation of Migratory Species of Wild Animals (CMS) and has been in force since November 2008. At the 12<sup>th</sup> CMS Conference of the Parties, in 2017, chimpanzees were listed as a migratory species under CMS.

The Gorilla Agreement is an important international legal instrument, as it is a Multilateral Environmental Agreement covering all 10 gorilla range States. The agreement provides governments, IGOs, NGOs, scientists, local people and the international community at large with a legally-binding framework to maintain and rehabilitate gorilla populations and their habitats. The Agreement is administered by the CMS Secretariat, which works closely with GRASP and benefits from partnerships with other organizations, including CITES and the IUCN Species Survival Commission (SSC). Central African Republic, Congo, DRC, Gabon, Nigeria, Rwanda and Uganda are Parties to the agreement. Equatorial Guinea has signed but not ratified, and Angola and Cameroon have not signed yet (Gorilla Agreement, 2008: [www.cms.int/en/legalinstrument/gorilla-agreement](http://www.cms.int/en/legalinstrument/gorilla-agreement)).

As many great ape habitats extend beyond political state borders, land-use planning and protected areas management require cross-border cooperation. However, transboundary conservation efforts can easily become complex. To be successful, they require high-level political commitment, sustained financial and technical support, cross-sectorial collaboration, strong institutional coordination and inclusion of all impacted stakeholders in the decision-making process. Examples of particularly encouraging initiatives include transboundary collaboration between DRC, Rwanda and Uganda

in the Greater Virunga Landscape (Refisch & Jenson, 2016); the Sangha Trinational, which involves Cameroon, Central African Republic and Congo; the Transboundary Peace Park for Sierra Leone and Liberia, which unites the Rainforest National Park in Sierra Leone and Gola Forest National Park in Liberia; the Mayombe Transboundary Initiative encompassing Luki Reserve in DRC, Dimonika Reserve in Congo and Maiombe National Park in Cabinda, Angola (Ron & Refisch, 2013); and the Heart of Borneo Initiative uniting the interior regions of Indonesian and Malaysian Borneo and Brunei Darussalam.

## 6. Challenges and solutions

Almost all great ape populations are in decline in both Africa and Asia, some so drastically that the population viability is in doubt. Following similar developments in Southeast Asia, the African landscape will soon experience a dramatic change through large-scale investments in extractive industries, transport infrastructure and commercial agriculture (Edwards, et al., 2014; Wich, et al., 2014; Laurance, et al., 2015; Rainer, et al., 2015; Sloan, et al., 2016). While these industries can help lift people out of poverty, operations must be sensitive to the biodiversity values and ecosystem services on which local communities and wildlife heavily rely. Hence, inclusive and integrated land-use planning processes and implementation of biodiversity-friendly business practices are required if such economic activities are to be accommodated sustainably in great ape habitats. The most sensitive ecosystems and biologically-rich great ape habitats should be protected, ideally through designation as off-limits to habitat modification. In this respect, a number of initiatives seek to minimize the negative impacts of industrial activities (agriculture and extractives) on biodiversity. These efforts include the Forestry Stewardship Council (FSC) for sustainable logging, the Roundtable for Sustainable Palm Oil (RSPO), the Business and Biodiversity Offsets Programme (BBOP) and associated standards (BBOP Standard on Biodiversity Offsets), the International Finance Cooperation (IFC) Performance Standard 6, the Equator Principles, the Global Oil and Gas Industry Association for Environmental and Social issues (IPIECA), and the International Council on Mining and Metals (ICMM). The description of each initiative and its potential benefit for the conservation of great apes and their habitats goes beyond the purpose of this report.

Law enforcement alone is insufficient to halt illegal trafficking of live great apes or their body parts. Arrests and convictions for holding or selling great apes (or bushmeat of any kind, including ape meat) rarely occur (Nijman, 2005). Stiles, et al., (2013) reported just 27 arrests for great ape trade between 2005 and 2011 – one quarter of which were never prosecuted. Peer-reviewed studies found that weak enforcement and lenient sentencing were inadequate deterrents of crime, recording only one prosecution for orangutan trading between 2006 and 2016 in two provinces of West Kalimantan, Indonesia (Freund, et al., 2017; Nijman, 2017). Freund, et al. (2017) found that not one of 145 orangutan cases reported and 133 rescues/confiscations resulted in prosecution or charges. Nijman (2017) found that at least 440 formal confiscations and surrenders of orangutans to law enforcement agencies across Indonesia between 1993 and 2016 resulted in only seven successful convictions. Reported convictions have been criticized for light sentences, which are unlikely to deter further crime. For example, four oil palm plantation workers were sentenced to 8–10 months in prison for killing two orangutans, and four people were sentenced to only eight months in prison for killing 20 orangutans (Jakarta Post, 2012). In addition, corruption can render wildlife law enforcement ineffectual, as deterrence becomes difficult to achieve (Bennett, 2015). Many of the world's great ape range States score poorly on the Corruption Perceptions Index (Transparency International, 2018). In many cases, convicted and incarcerated criminals have been able to bribe their way to liberty (Martini, 2013; WWF & TRAFFIC, 2015; Wyatt & Ngoc Cao, 2015). Without convictions and effective deterrence, it will not be possible to reduce the rates of illegal killing that threaten great apes. In this context, understanding the whole chain of actors involved in the bushmeat trade is necessary in order to address the threat. Caspary, et al. (2001) described the entire bushmeat network around the Tai National Park in Côte d'Ivoire, including urban bushmeat markets and restaurants.

Improvements in law enforcement efforts are underway in several countries. The EAGLE network, which started in Cameroon, has now been replicated in Benin, Congo, Côte d'Ivoire, Gabon, Guinea, Madagascar, Senegal, Togo and Uganda, with increasingly visible success (EAGLE Network, 2018). Partnerships between EAGLE members and national governments result in environmental crimes being detected, followed up and brought to justice. Their focus is usually ivory, as well as on great ape meat and live trade. For example, efforts by the EAGLE network member LAGA, in collaboration with the government in Cameroon, resulted in a record number of eight court judgments against 13 wildlife traffickers in 2017, with two traffickers receiving maximum penalties of three years imprisonment (EAGLE Network, 2017). In Indonesia in 2016, three orangutan smugglers were sentenced to up to 2.5 years in prison (Tempo, 2016), and in 2017, an orangutan killer was given a sentence of two years and nine months in prison (Robin des Bois, 2016; the organisation produces a quarterly newsletter of wildlife illegal trade records across the world). More work is needed to build on these efforts and establish deterrence against further illegal capture, killing, holding or sale of great apes across all range States.

In the context of the CITES mandate, more research is required to understand the scale of illegal cross-border trade in live great apes and their body parts. As detailed above, bushmeat crosses borders with little or no control. It is often difficult to identify body parts to species level (meat can be smoked and/or cut into sections), or by country of origin.

Both illegal meat and live trade are facilitated by corruption, lack of law enforcement, and in some range States, impunity. High profit margins and low risks to wildlife criminals have meant that people engaged in other types of crime (e.g. trafficking of drugs) have added wildlife crime to their activities. Furthermore, e-commerce has given suppliers and smugglers unprecedented access to new markets (Estrada, et al., 2017). The development of social media outlets as venues for wildlife trade is responsible for an explosion of illegal trade in numerous species that are now sold illicitly via the Internet (CITES CoP17 Doc. 29; Harrison, et al., 2016). CITES Decisions 17.92-17.96 direct Parties, the CITES Secretariat and the CITES Standing Committee to review and develop measures to combat this wildlife cybercrime (CITES CoP17 Decisions, 2016). The online trade is stimulated by the countless photos on social media of people holding wild animals taken in various situations (e.g. rescue facilities, zoos). Such images can influence the perceptions people have of those species' status and promote the idea of possessing them as pets (e.g. Ross, et al., 2008; Leighty, et al., 2015). Great apes are highly symbolic in the global illegal wildlife trade. The media is increasingly covering this topic and regular reports give cause for continued concern (e.g. Shukman, 2017). However, until recently, there has been a significant lack of verified quantitative and qualitative data on the trade in great apes, including the circumstances surrounding confiscations, making it hard to define longer-term strategies to combat this high-profile issue. To address this gap, GRASP, together with its technical partner, the UNEP-World Conservation Monitoring Centre (UNEP-WCMC), has developed the first phase of the Apes Seizure database (<https://database.un-grasp.org/>). This database includes data on live ape trade and the illegal trade in great ape parts and carcasses. The application was launched at the 17<sup>th</sup> CITES Conference of the Parties, in October 2016, in Johannesburg, and is currently undergoing a data validation process. It is anticipated that the database will provide analytics on illegal trade for a Status Report to the CITES CoP18 in 2019.

Further development of the Apes Seizure database can work hand-in-hand with the application of new advances in genetic recognition technology. According to CITES regulations, when possible, live victims of illegal wildlife trade should be repatriated to their countries of origin. However, when a great ape is confiscated, it carries little identification other than its DNA. Substantial genetic data collected to date on many wild great ape populations are a good starting point for a genetic recognition resource (e.g. Goldberg, 1997; Hvilson, et al., 2013). Additional reference samples will be required, with particular attention paid to populations less well-documented by existing research projects. In cases of illegally-trafficked great apes or great ape meat, DNA can be analysed to inform decisions on repatriation of live individuals to their country of origin, and to identify the source population, thus better directing future law enforcement efforts. Progress has been made in the development of mobile DNA sequencers (e.g. MiniIon from Oxford Nanopore Technologies). Providing sequencing units to law enforcement and customs officers, along with the necessary training, would enable genetic sampling on the spot and improve the identification of bushmeat or the origin of live animals.

In 2016, CITES introduced a new annual report format for illegal trade, but only a very small number of records of illegal trade in great apes have been included in the reports submitted so far.

Given the wide range of different threats and their complex interlinkages, it is reasonable to assume that addressing any one of these threats alone will not be enough to achieve conservation goals. Great ape habitats continue to be degraded because developmental spatial planning does not take conservation into account, spatial land-use planning is rarely done at a national scale, and law enforcement is weak, or a combination of these factors. Human populations continue to grow rapidly (especially in Sub-Saharan Africa, doubling every 30 years, and in Indonesia where the population will double in about 70 years; World Bank, 2018), putting ever-increasing pressure on land and natural resources. Therefore, great ape meat and live apes will thus continue to appear in the illegal wildlife trade. A holistic approach is required to tackle these challenges effectively, with strong political commitment and coordination between the various actors involved in law enforcement and conservation, including cross-border collaboration. Responses to address the drivers of illegal bushmeat trade and of international illegal live ape by criminal cartels are required.

One approach to addressing the complex interlinkages is coordination among different law enforcement stakeholders at national and regional levels, which remains a challenge. The International Consortium on Combatting Wildlife Crime (ICCWC) is one example where main law enforcement actors strive to coordinate and pool efforts. Within this Consortium, CITES, UNODC, INTERPOL and WCO partner to strengthen criminal justice systems and provide coordinated support at national, regional and international levels to combat wildlife and forest crime (<https://cites.org/eng/prog/iccwc.php>).

Often proposed as an opportunity, tourism with great apes is seen as a means of generating revenue to fund conservation efforts and to protect great apes and their habitats, or as a way for local communities to participate in, and benefit from, conservation activities. The success of mountain gorilla tourism has shown that conservation-based great ape tourism has considerable potential; however, it will not be possible to replicate this success at many other sites. The number of significant risks to great apes that can arise from tourism require a cautious approach. If great ape tourism is not based on sound conservation principles, economic objectives are likely to take precedence, the consequences of which could be detrimental to the great apes and their habitat. See IUCN Best Practice Guidelines for Great Ape Tourism (Macfie & Williamson 2010).

## 7. Recommendations

This report recommends the following actions directed to CITES Parties, recognizing that effective steps to protect great apes involve both great ape range States and the countries that import, or act as trade conduits for, live great apes, great ape meat, other body parts and derivatives, as well as other natural resources extracted from great ape habitats:

1. Recognizing that some CITES Parties may have already undertaken this analysis, all Parties are recommended to review relevant national and regional level legislation, policies and sanctions to ensure adequate protection of great apes through improved legal frameworks, with support from independent legal experts and with reference to the International Consortium in Combating Wildlife Crime (ICWC)'s Wildlife and Forest Crime Analytic Toolkit and USAID's *Measuring Efforts to Combat Wildlife Crime: A Toolkit for Improving Action and Accountability*;
2. Recognizing that many CITES Parties are already undertaking relevant actions in this regard, all Parties are recommended to increase law enforcement efforts by ensuring that existing laws are effectively applied and appropriate judicial processes are adhered to, with reference to the ICWC and USAID toolkits mentioned above. This can be attained by corruption mitigation strategies, and better training of local practitioners and rangers in law enforcement, prosecution evidence gathering, and modern forensic methods, as well as training of customs agents to profile suspect shipments and identify animal species to combat cross-border illegal trade;
3. All Parties are recommended to oblige, by national law, all private actors in the energy, extractives and agricultural sectors to comply with national and international laws and with IUCN and industry best practices in minimising impact on great ape populations and habitat, and should enforce clear penalties for cases of non-compliance<sup>4</sup>. Of particular benefit would be the inclusion of great apes as Species at Risk (SAR) in the FSC standards;
4. Great ape conservation partners are encouraged to contribute to, further develop and use the GRASP Apes Seizure database. These partners include the national law enforcement agencies and wildlife departments likely to hold data on seizures of trafficked bushmeat and live animals, relevant conservation NGOs involved in supporting law enforcement and compliance, researchers and wildlife sanctuaries;
5. All Parties are encouraged to accumulate all data at their disposal (e.g. protected area and conservation NGO reports, police reports, judiciary outcomes, etc.) in order to more consistently report illegal trade data to CITES in a timely manner in accordance with the annual illegal trade report format<sup>5</sup>, and should share data on illegal trade in great apes and great ape parts with GRASP for inclusion in the GRASP Apes Seizure database;
6. All Parties are recommended to use existing DNA sequence databases to establish the origin of confiscated great apes, and/or to support the potential repatriation of live animals;
7. All range State Parties are encouraged to consolidate data from protected area authorities, research and conservation organizations, commercial partners or other relevant stakeholders, in order to contribute great ape survey data and other relevant information to the IUCN SSC A.P.E.S. database by contacting Dr. Tenekwetch Sop (tenekwetch\_sop@eva.mpg.de);
8. All Parties that have not already done so are encouraged to ratify and fully implement the UN Convention against Transnational Organized Crime;
9. Recognizing the negative impact that corruption plays at all levels in support of the illegal capture and trade in great apes, all Parties that have not already done so are encouraged to ratify and fully implement the UN Convention against Corruption, and to solicit the support of national and international NGOs which specialize in combatting corruption, to enhance efforts to eradicate corruption and ensure appropriate judiciary process as it pertains to wildlife law enforcement;

---

<sup>4</sup> Industry best practices bodies are, for forestry: Forestry Stewardship Council (FSC), and IUCN guidelines on reduced impact logging and great apes (Morgan & Sanz, 2007), and on great apes and FSC standards (Morgan, et al., 2013); for industry in general: the Business and Biodiversity Offsets Program (BBOP) and BBOP Standard on Biodiversity Offsets, International Finance Corporation (IFC) Performance Standard 6, Equator Principles, IPIECA (global oil and gas industry association for environmental and social issues), and the International Council on Mining and Metals (ICMM).

<sup>5</sup> <https://cites.org/sites/default/files/reports/E-Guidelines-IllegalTR.pdf>

10. All Parties are encouraged to study dietary alternatives to bushmeat, and to monitor the implementation and impacts of projects piloting bushmeat alternatives, in order to determine their efficacy;
11. All range State Parties are encouraged to adopt legally binding transboundary agreements and foster collaborative judiciary proceedings with respect to evidence exchange, sentencing and extradition to enhance enforcement of illegal cross border trade in live apes, ape parts and bushmeat;
12. All Parties are encouraged to reject any applications for trade in potentially wild caught great apes (permits with source codes “W”, “U” or “I”). Given the rarity of these taxa and the large numbers of captive-bred great apes currently held in zoos and other ex-situ collections, wild-caught great apes are not acceptable for trade among zoos, safari parks or other educational or scientific institutions except under extraordinary circumstances.

The following recommendation is directed to the CITES Secretariat:

13. The Secretariat, in collaboration with other funders and partners such as UNODC where appropriate, is encouraged to provide technical and financial assistance to range States to further strengthen their criminal justice responses to wildlife crime, including tracing and recovering the proceeds of crime and clamping down on corruption.

The following recommendation is directed to private and public donors:

14. All private and public donors are encouraged to provide financial support to enable these recommendations to be implemented.

## 8. Main sources of information

Information on great ape population estimates and change over time has been derived from the IUCN SSC A.P.E.S. database. However, data from the more recent publications has not been entered yet. In these cases, we used information from the IUCN Red List to ensure that the information presented in this report is as up-to-date as possible.

## 9. Acknowledgements

We thank the members of the GRASP secretariat, namely, in alphabetical order, Milena Beekmann, Vinita Ganesh, Amelia Holmes, Dylan Jones, Emily Massingham, Tess Nicholls and Pauliina Upla for their contribution to this document and support throughout the process. Special thanks go to Milena Beekmann for her coordination efforts and her role in the peer-review process. For their contributions to the IUCN Red List assessment texts used in this report, we also thank Claudine André, Anna Behm Masozera, Christophe Boesch, Thomas Breuer, Damien Caillaud, Genevieve Campbell, Osiris Doumbé, Andrew Dunn, Andrew Fowler, Gabriella Fredriksson, Barbara Fruth, Takeshi Furuichi, Mary Gonder, Melvin Gumal, Jefferson Hall, John Hart, Terese Hart, Jena Hickey, Cleve Hicks, Simon Husson, Rachel Ikemeh, Inaoyom Imong, Kathryn Jeffery, Jessica Junker, Kathelijne Kooops, Andrew Marshall, Erik Meijaard, Bethan Morgan, David Morgan, Janet Nackoney, Denis Ndeloh, Aaron Nicholas, Radar Nishuli, Stuart Nixon, Graham Usher, Matt Nowak, Babafemi Ogunjemite, Alex Piel, Lilian Pintea, Andrew Plumpton, Gay Reinartz, Puji Rianti, Crickette Sanz, Ian Singleton, Volker Sommer, Emma Stokes, Samantha Strindberg, Jo Thompson and Suci Utami-Atmoko.

## 10. References

- Abernethy, K., Coad, L., Ilambu, O., Makiloutila, F., Easton J. and Akiak, J. (2010). Wildlife hunting, consumption and trade in the Oshwe sector of the Salonga-Lukenie-Sankuru Landscape. *WWF DRC*, 122.
- Aebischer, T., Siguindo, G., Rochat, E., Arandjelovic, M., Heilman, A., Hickisch, R., Vigilant, L., Joost, S. and Wegmann, D. (2017). ‘First quantitative survey delineates the distribution of chimpanzees in the Eastern Central African Republic’. *Biological Conservation* 213: 84–94.
- Allendorf, F.W., Luikart, G. and Aitken, S.N. (2013). *Conservation and the genetics of populations*. 2nd ed. Hoboken: Wiley-Blackwell.
- Ancrenaz, M., Gimenez, O., Ambu, L., Ancrenaz, K., Andau, P., Goossens, B., Payne, J., Tuuga, A. and Lackman-Ancrenaz, I. (2005). ‘Aerial surveys give new estimates for orangutans in Sabah, Malaysia’. *PLoS Biology* 3: 30–37.

- Ancrenaz, M., Ambu, L., Sunjoto, I., Ahmad, E., Manokaran, K., Meijaard, E. and Lackman, I. (2010). 'Recent surveys in the forests of Ulu Segama Malua, Sabah, Malaysia, show that orangutans (*P.p. morio*) can be maintained in slightly logged forests'. *PLoS One* 5: e11510.
- Ancrenaz, M., Cheyne, S.M., Humle, T. and Robbins, M.M. (2015). 'Impacts of industrial agriculture on ape ecology'. In: Arcus Foundation (ed.) *State of the Apes: Industrial Agriculture and Ape Conservation*. Cambridge, UK: Cambridge University Press.
- Ancrenaz, M., Gumal, M., Marshall, A.J., Meijaard, E., Wich, S.A. and Husson, S. (2016). *Pongo pygmaeus*. *The IUCN Red List of Threatened Species 2016*: e.T17975A17966347. <http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T17975A17966347.en>. Downloaded on 05 September 2017.
- Arandjelovic, M., Head, J., Rabanal, L.I., Schubert, G., Mettke, E., Boesch, C., Robbins, M.M. and Vigilant, L. (2011). 'Non-invasive genetic monitoring of wild central chimpanzees'. *PLoS One* 6: e14761.
- Bennett, E.L. (2015). 'Legal ivory trade in a corrupt world and its impact on African elephant populations'. *Conservation Biology* 29: 54–60.
- Bergl, R.A., Dunn, A., Fowler, A., Imong, I., Ndeloh, D., Nicholas, A. and Oates, J.F. (2016). *Gorilla gorilla* ssp. *diehli*. *The IUCN Red List of Threatened Species 2016*: e.T39998A102326240. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T39998A17989492.en>. Downloaded on 12 September 2017.
- Bermejo, M., Rodriguez-Teijeiro, J.D., Illera, G., Barroso, A., Vila, C. and Walsh, D.P. (2006). 'Ebola outbreak killed 5000 gorillas'. *Science* 314(5805): 1564–1564.
- Boyer, K. (2011). 'Chimpanzee conservation in light of impending iron ore mining project in S.E Senegal'. Graduate Masters thesis. Iowa State University.
- Brcic, T.M., Amarasekaran, B. and McKenna, A. (2010). Final report of the Sierra Leone national chimpanzee census project. Freetown, Sierra Leone: Tacugama Chimpanzee Sanctuary.
- Bryson-Morrison, N., Matsuzawa, T. and Humle, T. (2016). 'Chimpanzees in an anthropogenic landscape: examining food resources across habitat types at Bossou, Guinea, West Africa'. *International Journal of Primatology* 78: 1237–1249.
- Campbell, G., Kuehl, H., N'Goran Kouamé, P. and Boesch, C. (2008). 'Alarming decline of West African chimpanzees in Côte d'Ivoire'. *Current Biology* 18: R903–R904.
- Campbell, G. and Hougbedji, M. (2015). Conservation status of the West African chimpanzee (*Pan troglodytes verus*) in Togo and Benin. *Unpublished report to Primate Action Fund*.
- Campbell-Smith, G., Sembiring, R. and Linkie, M. (2012). 'Evaluating the effectiveness of human–orangutan conflict mitigation strategies in Sumatra'. *Journal of Applied Ecology* 49: 367–375.
- Carvalho, J.S., Marques, T.A. and Vicente, L. (2013). 'Population status of *Pan troglodytes verus* in Lagoas de Cufada Natural Park, Guinea-Bissau'. *PLoS One* 8: e71527.
- Caspary, H.U., Koné, I., Prouot, C. and De Pauw, M. (2001). La chasse et la filière viande de brousse dans l'espace Tai, Cote d'Ivoire. Tropenbos-Côte d'Ivoire Série 2, ISBN: 90-5113-048-1.
- Chaber, A.L., Allebone-Webb, S., Lignereux, Y., Cunningham, A.A. and Rowcliffe, J.M. (2010). 'The scale of illegal meat importation from Africa to Europe via Paris'. *Conservation Letters* 3: 317–321.
- CITES CoP17 Decisions (2016). [Online] <https://cites.org/eng/dec/valid17/81840>. Consulted on 22 March 2018.
- CITES (2017). Annexes. [Online] <https://cites.org/eng/app/appendices.php>
- Coad, L., Schleicher, J., Milner-Gulland, E.J., Marthews, T.R., Starkey, M., Manica, A., Balmford, A., Mbombe, W., Diop Binen, T.R. and Abernethy, K.A. (2013). 'Social and ecological change over a decade in a village hunting system, Central Gabon'. *Conservation Biology* 27: 270–280.

- Colom, A. (2006). The socioeconomic aspects of natural resource use and management by local communities in the Salonga-Lukenie-Sankuru landscape: guidelines for conservation and livelihood improvement. WWF, Democratic Republic of Congo, 250 pp.
- Danquah, E., Oppong, S.K., Akom, E. and Sam, M. (2012). 'Preliminary survey of chimpanzees and threatened monkeys in the Bia-Goaso forest block in southwestern Ghana'. *African Primates* 7: 163–174.
- Davis, J.T., Mengersen, K., Abram, N.K., Ancrenaz, M., Wells, J.A. and Meijaard, E. (2013). 'It's not just conflict that motivates killing of orangutans'. *PLoS One* 8: e75373.
- Dowsett, R.J. and Dowsett-Lemaire, F. ed. (1991). *Flore et Faune du bassin du Kouliou (Congo) et leur exploitation*. Liège, Belgium: Tauraco Press.
- Dunn, A., Bergl, R., Byler, D., Eben-Ebai, S., Etiendem, D.N., Fotso, R., Ikfuingei, R., Imong, I., Jameson, C., Macfie, L., Morgan, B., Nchanji, A., Nicholas, A., Nkembi, L., Omeni, F., Oates, J., Pokempner, A., Sawyer, S. and Williamson, E.A. (2014). *Revised regional action plan for the conservation of the Cross River gorilla (Gorilla gorilla diehli): 2014–2019*. IUCN SSC Primate Specialist Group and Wildlife Conservation Society, New York.
- EAGLE Network (2017). 'The EAGLE Network Annual Report 2017'. [Online] <http://www.eagle-enforcement.org/data/files/eagle-network-annual-report-2017.pdf>
- EAGLE Network (2018). 'Our Members'. [Online] [www.eagle-enforcement.org/members/](http://www.eagle-enforcement.org/members/)
- Edwards, D.P., Sloan, S., Weng, L., Dirks, P., Sayer, J. and Laurance, W.F. (2014). 'Mining and the African environment'. *Conservation Letters* 7: 302–311.
- Estrada, A., Garber, P.A., Rylands, A.B., Roos, C., Fernandez-Duque, E., Di Fiore, A., Nekaris, K.A.I., Nijman, V., Heymann, E.W., Lambert, J.E., Rovero, F., Barelli, C., Setchell, J.M., Gillespie, T.R., Mittermeier, R.A., Arregoitia, L.V., De Guinea, M., Gouveia, S., Dobrovolski, R., Shane, S., Shane, N., Boyle, S.A., Fuentes, A., MacKinnon, K.C., Amato, K.R., Meyer, A.L., Wich, S., Sussman, R.W., Pan, R., Koné, I. and Li, B. (2017). 'Impending extinction crisis of the world's primates: Why primates matter'. *Science Advances* 3: e1600946.
- Fa, J.E., Seymour, S., Dupain, J., Amin, R., Albrechtsen, L. and Macdonald, D. (2006). 'Getting to grips with the magnitude of exploitation: bushmeat in the Cross-Sanaga rivers region, Nigeria and Cameroon'. *Biological Conservation* 129: 497–510.
- Fa, J. and Brown, D. (2009). 'Impacts of hunting on mammals in African tropical moist forests: A review and synthesis'. *Mammal Review* 39: 231–264.
- FAO and WFP (2015). The State of Food Insecurity in the World 2015. Meeting the 2015 international hunger targets: taking stock of uneven progress. Food and Agriculture Organization Publications, Rome.
- Foerster, S., Wilkie, D., Morelli, G.A., Demmer, J., Starkey, M., Telfer, P., Steil, M. and Lewbel, A. (2012). 'Correlates of bushmeat hunting among remote rural households in Gabon, Central Africa'. *Conservation Biology* 26: 335–344.
- Freund, C., Rahman, E. and Knott, C. (2017). 'Ten years of orangutan related wildlife crime investigation in West Kalimantan, Indonesia'. *American Journal of Primatology* 79: e22620.
- Fruth, B., Hickey, J.R., André, C., Furuichi, T., Hart, J., Hart, T., Kuehl, H., Maisels, F., Nackoney, J., Reinartz, G., Sop, T., Thompson, J. and Williamson, E.A. (2016). *Pan paniscus*. *The IUCN Red List of Threatened Species 2016*: e.T15932A102331567. <http://dx.doi.org/10.2305>
- Galat, G., Galat-Luong, A. and Nizinski, G. (2008). Our cousins chimpanzees and baboons face global warming by digging wells to filtrate drinking water. In: *13th IWRA World Water Congress, IUCN*.
- Garriga, R.M., Marco, I., Casas-Díaz, E., Amarasekaran, B. and Humle, T. (2017). Perceptions of challenges to subsistence agriculture, and crop foraging by wildlife and chimpanzees *Pan troglodytes verus* in unprotected areas in Sierra Leone. *Oryx* 1–14.
- Gaveau, D.L., Sloan, S., Molidena, E., Yaen, H., Sheil, D., Abram, N.K., Ancrenaz, M., Nasi, R., Quinones, M., Wielaard, N. and Meijaard, E. (2014). 'Four decades of forest persistence, clearance and logging on Borneo'. *PLoS One* 9: e101654.

- Gilardi, K.V., Gillespie, T.R., Leendertz, F.H., Macfie, E.J., Travis, D.A., Whittier, C.A. and Williamson, E.A. (2015). Best practice guidelines for health monitoring and disease control in great ape populations. Gland, Switzerland: IUCN SSC Primate Specialist Group. 56 pp. <https://portals.iucn.org/library/node/45793>
- Ginn, L.P., Robison, J., Redmond, I. and Nekaris, K.A.I. (2013). ‘Strong evidence that the West African chimpanzee is extirpated from Burkina Faso’. *Oryx* 47: 325–326.
- Global Forest Watch (2017). [Online] [www.globalforestwatch.org/map/6/0.74/15.68/ALL/grayscale/none/581](http://www.globalforestwatch.org/map/6/0.74/15.68/ALL/grayscale/none/581). Consulted on 22 March 2018.
- Goldberg, T.L. (1997). ‘Inferring the geographic origins of ‘refugee’ chimpanzees in Uganda from mitochondrial DNA sequences’. *Conservation Biology* 11: 1441–1446.
- Goodall, J. (1986). *The Chimpanzees of Gombe: Patterns of Behaviour*. Cambridge, Massachusetts: Belknap Press.
- Gorilla Agreement (2008). [Online] [www.cms.int/en/legalinstrument/gorilla-agreement](http://www.cms.int/en/legalinstrument/gorilla-agreement). Consulted on 22 March 2018.
- Gray, M., McNeilage, A., Fawcett, K., Robbins, M.M., Ssebide, B., Mbula, D. and Uwingeli, P. (2010). ‘Censusing the mountain gorillas in the Virunga Volcanoes: complete sweep method versus monitoring’. *African Journal of Ecology* 48: 588–599.
- Gray, M., Roy, J., Vigilant, L., Fawcett, K., Basabose, A., Cranfield, M., Uwingeli, P., Mburanumwe, I., Kagoda, E. and Robbins, M.M. (2013). ‘Genetic census reveals increased but uneven growth of a critically endangered mountain gorilla population’. *Biological Conservation* 158: 230–238.
- Greengrass, E. (2016). ‘Commercial hunting to supply urban markets threatens mammalian biodiversity in Sapo National Park, Liberia’. *Oryx* 50: 397–404.
- Hakizimana, D. and Huynen, M.C. (2013). ‘Chimpanzee (*Pan troglodytes schweinfurthii*) population density and abundance in Kibira National Park, Burundi’. *Pan Africa News* 20: 16–19.
- Hall, J.S., White, L.J., Inogwabini, B.I., Omari, I., Morland, H.S., Williamson, E.A., Saltonstall, K., Walsh, P., Sikubwabo, C., Bonny, D., Kaleme, P.K., Vedder, A. and Freeman, K. (1998). ‘Survey of Grauer’s gorillas (*Gorilla gorilla graueri*) and eastern chimpanzees (*Pan troglodytes schweinfurthii*) in the Kahuzi-Biega National Park lowland sector and adjacent forest in eastern Democratic Republic of Congo’. *International Journal of Primatology*, 19: 207–235.
- Ham, R. (1998). Nationwide chimpanzee census and large mammal survey: Republic of Guinea. Unpublished report to the European Union, Conakry, Guinea.
- Harrison, J., Roberts, D. and Hernandez-Castro, J. (2016). ‘Assessing the extent and nature of wildlife trade on the dark web’. *Conservation Biology* 30: 900–904.
- Hedrick, P.W. and Kalinowski, S.T. (2000). ‘Inbreeding depression in conservation biology’. *Annual Review of Ecology and Systematics* 31: 139–162.
- Hickey, J.R., Granjon, A.C., Vigilant, L., Eckardt, W., Gilardi, K.V., Babaasa, D., Ruzigandekwe, F., Leendertz, F.H. and Robbins, M.M. (in prep.). Virunga 2015–2016 surveys: monitoring mountain gorillas, other select mammals, and illegal activities.
- Hicks, C., Darby, L., Hart, J., Swinkels, J., January, N. and Menken, S. (2010). ‘Trade in orphans and bushmeat threatens one of the Democratic Republic of the Congo’s most important populations of eastern chimpanzees (*Pan troglodytes schweinfurthii*)’. *African Primates* 7: 1–18.
- Hockings, K. and Humle, T. (2009). *Best Practice Guidelines for the Prevention and Mitigation of Conflict Between Humans and Great Apes*. Gland, Switzerland: IUCN SSC Primate Specialist Group. 40 pp. [Online] <https://portals.iucn.org/library/node/9324>
- Humle, T., Boesch, C., Campbell, G., Junker, J., Koops, K., Kuehl, H and Sop, T. (2016), *Pan troglodytes* ssp. *verus*. *The IUCN Red List of Threatened Species 2016*: e.T15935A102327574. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T15935A17989872.en>

- Hvilsom, C., Frandsen, P., Børsting, C., Carlsen, F. Sallé, B., Simonsen, B.T. and Siegismund, H.R. (2013). 'Understanding geographic origins and history of admixture among chimpanzees in European zoos, with implications for future breeding programmes'. *Heredity* 110: 586–593.
- ICCWC (International Consortium on Combating Wildlife Crime) (2012). *Wildlife and Forest Crime Analytic Toolkit (Revised edition)*. New York: United Nations. [Online] [http://www.unodc.org/documents/Wildlife/Toolkit\\_e.pdf](http://www.unodc.org/documents/Wildlife/Toolkit_e.pdf). Downloaded on May 6, 2018.
- Imong, I., Robbins, M.M., Mundry, R., Bergl, R.A. and Kühl, H.S. (2014). 'Informing conservation management about structural versus functional connectivity: a case study of Cross River gorillas'. *American Journal of Primatology* 76: 978–988.
- Inogwabini, B.I., Matungila, B., Mbende, L., Abokome, M. and Wa Tshimanga, T. (2007). 'Great apes in the Lake Tumba landscape, Democratic Republic of Congo: newly described populations'. *Oryx* 41: 532–538.
- IUCN (2014). *Regional Action Plan for the Conservation of Western Lowland Gorillas and Central Chimpanzees 2015–2025*. Gland, Switzerland: IUCN SSC Primate Specialist Group.
- IUCN (2018). *The IUCN Red List of Threatened Species. Version 2018-1*. [Online] [www.iucnredlist.org](http://www.iucnredlist.org)
- IUCN and ICCN. (2012). *Bonobo (Pan paniscus): Conservation Strategy 2012–2022*. Gland, Switzerland: IUCN SSC Primate Specialist Group & Institut Congolais pour la Conservation de la Nature.
- IUCN SSC A.P.E.S. database (2017). [Online] <http://apesportal.eva.mpg.de/database/archive>
- Jakarta Post (2012). 'Orangutan Killers Sent to Prison in Kalimantan Case'. [Online] (23 May 2012). <http://jakartaglobe.id/archive/orangutan-killers-sent-to-prison-in-kalimantan-case/>. Consulted on 22 March 2018.
- JGI, TNC, CBSG, FZS, WCS, WD-MNRT, TAWIRI and USFWS. (2011). Tanzania chimpanzee conservation action planning workshop report. Jane Goodall Institute, Arlington, USA.
- Junker, J., Boesch, C., Freeman, T., Mundry, R., Stephens, C. and Kühl, H. S. (2015). 'Integrating wildlife conservation with conflicting economic land-use goals in a West African biodiversity hotspot'. *Basic and Applied Ecology* 16: 690–702.
- Kirkby, A., Spira, C., Bahati, B., Twendilongé, A., Kujirakwinja, D., Plumptre, A.J., Wieland, M. and Nishuli, R. (2015). Investigating artisanal mining and bushmeat around protected areas: Kahuzi-Biega National Park and Itombwe Reserve. Report to USAID and Arcus Foundation. Wildlife Conservation Society, New York.
- Köndgen, S., Kühl, H., N'Goran, P.K., Walsh, P.D., Schenk, S., Ernst, N., Biek, R., Formenty, P., Mätz-Rensing, K., Schweiger, B., Junglen, S., Ellerbrok, H., Nitsche, A., Briese, T., Lipkin, W.I., Pauli, G., Boesch, C. and Leendertz, F. (2008). 'Pandemic human viruses cause decline of endangered great apes'. *Current Biology* 18: 260–264.
- Kormos, R., Boesch, C., Bakarr, M.I. and Butynski, T.M. (2003). *West African chimpanzees: status survey and conservation action plan*. Gland, Switzerland and Cambridge, UK: IUCN SSC Primate Specialist Group.
- Kormos, R., Kormos, C.F., Humle, T., Lanjouw, A., Rainer, H., Victorine, R., Mittermeier, R.A., Diallo, M.S., Rylands, A.B. and Williamson, E.A. (2014). 'Great apes and biodiversity offset projects in Africa: The case for national offset strategies'. *PLoS One* 9: e111671.
- Kühl, H., Maisels, F., Ancrenaz, M. and Williamson, E.A. (2008). *Best Practices Guidelines for Surveys and Monitoring of Great Ape Populations*. Gland, Switzerland: IUCN SSC Primate Specialist Group.
- Kühl, H., Sop, T., Williamson, E.A., Mundry, R., Brugière, D., Campbell, G., Cohen, H., Danquah, E., Ginn, L., Herbinger, I., Jones, S., Junker, J., Kormos, R., Kouakou, C., N'Goran, P., Normand, E., Shutt-Phillips, K., Tickle, A., Vendras, E., Welsh, A., Wessling, E. and Boesch, C. (2017). 'The Critically Endangered western chimpanzee declines by 80%'. *American Journal of Primatology* 79: e22681.
- Laurance, W., Sloan, S., Weng, L. and Sayer, J.A. (2015). 'Estimating the environmental costs of Africa's massive "development corridors"'. *Current Biology* 25: 3202–3208.

- Leendertz, F., Pauli, G., Maetz-Rensing, K., Boardman, W., Nunn, C., Ellerbrok, H., Jensen, S., Junglen, S. and Christophe, B. (2006). Pathogens as drivers of population declines: 'The importance of systematic monitoring in great apes and other threatened mammals'. *Biological Conservation* 131: 325–337.
- Leendertz, S.A.J., Wich, S.A., Ancrenaz, M., Bergl, R.A., Gonder, M.K., Humle, T. and Leendertz, F.H. (2017). 'Ebola in great apes – current knowledge, possibilities for vaccination, and implications for conservation and human health'. *Mammal Review* 47: 98–111.
- Leighty, K., Valuska, A., Grand, A., Bettinger, T., Mellen, J., Ross, S., Boyle, P. and Ogden, J. (2015). 'Impact of visual context on public perceptions of non-human primate performers'. *PLoS One* 10: e0118487.
- Macfie, E.J. & Williamson, E.A. (2010). 'Best practices guidelines for great ape tourism'. Gland, Switzerland: IUCN SSC Primate Specialist Group. <https://portals.iucn.org/library/node/9636>
- Maisels, F., Strindberg, S., Breuer, T., Greer, D., Jeffery, K. and Stokes, E. (2016a). *Gorilla gorilla* ssp. *gorilla*. *The IUCN Red List of Threatened Species 2016*: e.T9406A17989591. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T9406A17989591.en>
- Maisels, F., Strindberg, S., Greer, D., Jeffery, K., Morgan, D.L. and Sanz, C. (2016b). *Pan troglodytes* ssp. *troglodytes*. *The IUCN Red List of Threatened Species 2016*: e.T15936A17990042. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T15936A17990042.en>
- Maisels, F., Bergl, R.A. and Williamson, E.A. (2016c). *Gorilla gorilla*. *The IUCN Red List of Threatened Species 2016*: e.T9404A102330408. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T9404A17963949.en>
- Martini, M. (2013). Wildlife crime and corruption: In which way does corruption exacerbate the problem of poaching and illegal wildlife trade in Southern Africa and how can anti-corruption contribute to the fight against it? U4 Expert Answer (15 February 2013), Number 367. U4 Anti-Corruption Resource Centre. [www.transparency.org/files/content/corruptionqas/367\\_Wildlife\\_Crimes\\_and\\_Corruption.pdf](http://www.transparency.org/files/content/corruptionqas/367_Wildlife_Crimes_and_Corruption.pdf)
- Matsuzawa, T., Humle, T. and Sugiyama, Y. (eds.). (2011). *The Chimpanzees of Bossou and Nimba*. Japan: Springer.
- Meijaard, E., Buchori, D., Hadiprakarsa, Y., Utami-Atmoko, S.S., Nurcahyo, A., Tjiu, A., Prasetyo, D., Nardiyono, Christie, L., Ancrenaz, M., Abadi, F., Antoni, N.G., Armayadi, D., Dinato, A., Ella, Gumelar, P., Indrawan, T.P., Kussaritano, Munajat, C., Priyono, W.P., Purwanto, Y., Puspitasari, D., Putra, M.S.W., Rahmat, A., Ramadani, H., Sammy, J., Siswanto, D., Syamsuri, M., Andayani, N., Wu, H., Wells, J.A. and Mengersen, K. (2011). 'Quantifying killing of orangutans and human-orangutan conflict in Kalimantan, Indonesia'. *PLoS One* 6: e27491.
- Mitchell M.W., Locatelli S., Ghobrial L., Pokempner, A.A., Sesink Clee, P.R., Abwe, E.E., Nicholas, A., Nkambi, L., Anthony, N.M., Morgan, B.J., Fotso, R., Peeters, M., Hahn, B.H. and Gonder, M.K. (2015). 'The population genetics of wild chimpanzees in Cameroon and Nigeria suggests a positive role for selection in the evolution of chimpanzee subspecies'. *BMC Evolutionary Biology* 15: 3.
- Morgan, D. and Sanz, C. (2007). *Best Practice Guidelines for Reducing the Impact of Commercial Logging on Great Apes in Western Equatorial Africa*. Gland, Switzerland: IUCN SSC Primate Specialist Group. <https://portals.iucn.org/library/node/9059>
- Morgan, B., Adeleke, A., Basse, T., Bergl, R., Dunn, A., Fotso, R., Gadsby, E., Gonder, K., Greengrass, E., Koulagna, D.K., Mbah, G., Nicholas, A., Oates, J., Omeni, F., Saidu, Y., Sommer, V., Sunderland-Groves, J., Tiebou, J. and Williamson, E.A. (2011). Regional action plan for the conservation of the Nigeria-Cameroon chimpanzee (*Pan troglodytes ellioti*). San Diego, USA: IUCN SSC Primate Specialist Group and Zoological Society of San Diego.
- Morgan, D., Mundry, R., Sanz, C., Ayina, C.E., Strindberg, S., Lonsdorf, E. and Kühl, H.S. (2017). 'African apes coexisting with logging: Comparing chimpanzee (*Pan troglodytes troglodytes*) and gorilla (*Gorilla gorilla gorilla*) resource needs and responses to forestry activity'. *Biological Conservation* 218: 277–286.
- Morgan, D., Sanz, C., Greer, D., Rayden, T., Maisels, F. and Williamson, E.A. (2013). *Great Apes and FSC: Implementing 'Ape Friendly' Practices in Central Africa's Logging Concessions*. Gland, Switzerland: IUCN SSC Primate Specialist Group. <https://portals.iucn.org/library/node/10378>
- Murai, M., Ruffler, H., Berlemont, A., Campbell, G., Esono, F., Agbor, A., Mbomio, D., Ebana, A., Nze, A. and Kühl, H. (2013). 'Priority areas for large mammal conservation in Equatorial Guinea'. *PLoS One* 8: e75024.

- Nakamura, M. and Nishida, T. (2012). 'Long-term field studies of chimpanzees at Mahale Mountains National Park, Tanzania'. In: Kappeler, P.M. and Watts, D. (eds). *Long-Term Field Studies of Primates* (pp. 339–356). Berlin Heidelberg: Springer-Verlag.
- Nasi, R., Brown, D., Wilkie, D., Bennett, E.L., Tutin, C., van Tol, G. and Christophersen, T. (2008). Conservation and use of wildlife-based resources: the bushmeat crisis. Technical Series no. 33: 50 pp. Secretariat of the Convention on Biological Diversity, Montreal, Canada, and Centre for International Forestry Research, Bogor, Indonesia.
- Nater, A., Mattle-Greminger, M.P., Nurcahyo, A., Nowak, M.G., de Manuel, M., Desai, T., Grovers, C., Pybus, M., Sonay, T.B., Roos, C., Lameira, A.R., Wich, S.A., Askew, J., Davila-Ross, M., Fredriksson, G., de Valles, G., Casais, F., Prado-Martinez, J., Goossens, B., Verschoor, E.J., Warren, K.S., Singleton, I., Marques, D.A., Pamungkas, J., Perwitasari-Farajallah, D., Rianti, P., Tuuga, A., Gut, I.G., Gut, M., Orozco-terWengel, P., van Schaik, C.P., Bertranpetit, J., Anistimova, M., Scally, A., Marques-Bonet, T., Meijaard, E. and Krützen, M. (2017). 'Morphometric, behavioural and genomic evidence for a new orangutan species'. *Current Biology* 27: 3487–3498.
- Ndiaye, S. (2011). Conservation du Chimpanzé au Sénégal. Etat des Connaissances et Réactualisation du Plan d'Actions. Report for USAID/Wula Nafaa.
- Nijman, V. (2005). *In Full Swing: An Assessment of trade in Orang-utans and Gibbons on Java and Bali, Indonesia*. [1]TRAFFIC Southeast Asia, Selangor, Malaysia.
- Nijman, V. (2009). *An assessment of trade in gibbons and orang-utans in Sumatra, Indonesia*. [1]TRAFFIC Southeast Asia, Selangor, Malaysia.
- Nijman, V. (2017). 'Orangutan trade, confiscations, and lack of prosecutions in Indonesia'. *American Journal of Primatology* 79: e22652.
- Nishida, T., Corp, N., Hamai, M., Hasegawa, T., Hiraiwa-Hasegawa, M., Hosaka, K., Hunt, K., Itoh, N., Kawanaka, K., Matsumoto-Oda, A., Mitani, J., Nakamura, M., Norikoshi, K., Sakamaki, T., Turner, L., Uehara, S. and Zamma, K. (2003). 'Demography, female life history, and reproductive profiles among the chimpanzees of Mahale'. *American Journal of Primatology* 59: 99–121.
- N'Goran, K.P., Ndomba, D.L. and Beukou, G.B. (2016). Rapport de l'inventaire des grands et moyens mammifères dans le segment RCA du paysage trinational de la Sangha. Bureau Régional du WWF pour l'Afrique. Unpublished report.
- Nowak, M.G., Rianti, P., Wich, S.A., Meijaard, E. and Fredriksson, G. (2017). *Pongo tapanuliensis*. *The IUCN Red List of Threatened Species 2017*: e.T120588639A120588662. <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T120588639A120588662.en>. Downloaded on 24 March 2018.
- Oates, J.F., Doumbe, O., Dunn, A., Gonder, M.K., Ikemeh, R., Imong, I., Morgan, B.J., Ogunjemite, B. and Sommer, V. (2016). *Pan troglodytes ssp. ellioti*. *The IUCN Red List of Threatened Species 2016*: e.T40014A17990330. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T40014A17990330.en>. Downloaded on 06 September 2017.
- Ondoua Ondoua, G., Beodo Moundjim, E., Mambo Marindo, J.C., Jiagho, R., Usongo, L. and Williamson, E.A. (2017). *An Assessment of Poaching and Wildlife Trafficking in the Garamba-Bili-Chinko Transboundary Landscape*. Cambridge, UK: TRAFFIC.
- Plumptre, A.J., Rose, R., Nangendo, G., Williamson, E.A., Didier, K., Hart, J., Mulindahabi, F., Hicks, C., Griffin, B., Ogawa, H., Nixon, S., Pintea, L., Vosper, A., McClennan, M., Amsini, F., McNeilage, A., Makana, J.R., Kanamori, M., Hernandez, A., Piel, A., Stewart, F., Moore, J., Zamma, K., Nakamura, M., Kamenya, S., Idani, G., Sakamaki, T., Yoshikawa, M., Greer, D., Tranquilli, S., Beyers, R., Furuichi, T., Hashimoto, C. and Bennett, E. (2010). *Eastern Chimpanzee (Pan troglodytes schweinfurthii): Status Survey and Conservation Action Plan 2010–2020*. Gland, Switzerland: IUCN SSC Primate Specialist Group.
- Plumptre, A.J., Nixon, S., Critchlow, R., Vieilledent, G., Nishuli, R., Kirkby, A., Williamson, E.A., Hall, J.S. and Kujirakwinja, D. (2015). *Status of Grauer's gorilla and chimpanzee in eastern Democratic Republic of Congo: Historical and current distribution and abundance*. New York: Wildlife Conservation Society, Fauna & Flora International and Institut Congolais pour la Conservation de la Nature.
- Plumptre, A.J., Nixon, S., Kujirakwinja, D.K., Vieilledent, G., Critchlow, R., Williamson, E.A., Nishuli, R., Kirkby, A.E. and Hall, J.S. (2016a). 'Catastrophic decline of world's largest primate: 80% loss of Grauer's gorilla (*Gorilla beringei graueri*) population justifies Critically Endangered status'. *PLoS One* 11: e0162697.

- Plumptre, A., Hart, J.A., Hicks, T.C., Nixon, S., Piel, A.K. and Pintea, L. (2016b). *Pan troglodytes* ssp. *schweinfurthii*. *The IUCN Red List of Threatened Species 2016*: e.T15937A17990187. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T15937A17990187.en>
- Plumptre, A., Nixon, S., Caillaud, D., Hall, J.S., Hart, J.A., Nishuli, R. and Williamson, E.A. (2016c). *Gorilla beringei* ssp. *graueri*. *The IUCN Red List of Threatened Species 2016*: e.T39995A102328430. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T39995A17989838.en>
- Plumptre, A., Robbins, M. and Williamson, E.A. (2016). *Gorilla beringei*. *The IUCN Red List of Threatened Species 2016*: e.T39994A102325702. <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T39994A17964126.en>
- Poulsen, J.R. and Clark, C.J. (2010). Congo Basin timber certification and biodiversity conservation. *Biodiversity Conservation in Certified Forests* 51: 55–60.
- Rainer, H., White, A. and Lanjouw, A. (2014). *State of the Apes: Extractive Industries and Ape Conservation*. Cambridge, UK: Cambridge University Press.
- Rainer, H., White, A. and Lanjouw, A. (2015). *State of the Apes: Industrial Agriculture and Ape Conservation*. Cambridge, UK: Cambridge University Press.
- Redford, K. (1992). ‘The empty forest’. *BioScience* 42: 412–422.
- Refisch, J. and Koné, I. (2005). ‘Impact of commercial hunting on monkey populations in the Tai region, Cote d'Ivoire’. *Biotropica* 37: 136–144.
- Refisch, J. and Jenson, J. (2016). ‘Transboundary collaboration in the Greater Virunga landscape: From gorilla conservation to conflict-sensitive transboundary landscape management’. In: Bruch, C., Muffett, C. and Nichols, S. (eds.). *Governance, Natural Resources and Post-Conflict Peacebuilding*. Oxford, UK: Earthscan from Routledge.
- Regnaut, S. and Boesch, C. (2012). Offset strategy for the Guinean Alumina project. Conakry, Republic of Guinea: Wild Chimpanzee Foundation.
- Robbins, M., Gray, M., Kümpel, N., Lanjouw, A., Maisels, F., Mugisha, A., Spelman, L. and Williamson, L. (2008). *Gorilla beringei* ssp. *beringei*. *The IUCN Red List of Threatened Species 2008*: e.T39999A10292321. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T39999A10292321.en>
- Robbins, M.M., Gray, M., Fawcett, K.A., Nutter, F.B., Uwingeli, P., Mburanumwe, I., Kagoda, E., Basabose, A., Stoinski, T.S., Cranfield, M.R., Byamukama, J., Spelman, L.H. and Robbins, A.M. (2011). ‘Extreme conservation leads to recovery of the Virunga Mountain gorillas’. *PLoS One* 6: e19788.
- Robin des Bois (2016). ‘Three Sentenced to Prison over Orangutan Illegal Trade.’ [Online] (23rd March, 2016), <https://en.tempo.co/read/news/2016/03/23/206756195/Three-Sentenced-to-Prison-over-Orangutan-Illegal-Trade>
- Ron, T. and Refisch, J. (2013). Towards a transboundary protected area complex in the Mayombe forest ecosystems: Strategic Plan. UNEP, GRASP and IUCN.
- Ross, S.R., Lukas, K.E., Lonsdorf, E.V., Stoinski, T.S., Hare, B., Shumaker, R. and Goodall, J. (2008). ‘Inappropriate use and portrayal of chimpanzees’. *Science* 319(5869): 1487.
- Roy, J., Vigilant, L., Gray, M., Wright, E., Kato, R., Kabano, P., Basabose, A., Tibenda, E., Kühl, H. and Robbins, M. (2014). ‘Challenges in the use of genetic mark-recapture to estimate the population size of Bwindi mountain gorillas (*Gorilla beringei beringei*)’. *Biological Conservation* 180: 249–261.
- Sá, R., Ferreira da Silva, M.J., Sousa, F.M. and Minhós, T. (2012). ‘The trade and ethnobiological use of chimpanzee body parts in Guinea-Bissau: Implications for Conservation’. *TRAFFIC Bulletin* 24: 31–34.
- Sakamaki, T., Mulavwa, M. and Furuichi, T. (2009). ‘Flu-like epidemics in wild bonobos (*Pan paniscus*) at Wamba, the Luo Scientific Reserve, Democratic Republic of Congo’. *Pan Africa News* 16: 1–4.
- Santika, T., Ancrenaz, M., Wilson, K., Spehar, S., Abram, N., Banes, G., Campbell-Smith, G., Curran, L., d’Arcy, L., Delgado, R., Erman, A., Goossens, B., Hartanto, H., Houghton, M., Husson, S., Kühl, H., Lackman, I., Leiman, A., Llano Sanchez, K., Makinuddin, N., Marshall, A., Meididit, A., Mengersen, K., Musnanda, Nardiyono, Nurcahyo, A., Odom, K., Panda, A., Prasetyo, D., Purnomo, Rafiastanto, A., Raharjo, S., Ratnasari, D., Russon, A., Santana, A.,

- Santoso, E., Sapari, I., Sihite, J., Suyoko, A., Tjiu, A., Utami-Atmoko, S., van Schaik, C., Voigt, M., Wells, J., Wich, S., Willems, E. and Meijaard, E. (2017). 'First integrative trend analysis for a great ape species in Borneo'. *Scientific Reports* 7: 4839.
- Shukman, D. (2017). 'The secret trade in baby chimps: Full report'. *BBC News*. [Online] (31st Jan 2017). [www.bbc.com/news/science-environment-38804688](http://www.bbc.com/news/science-environment-38804688). Consulted on 22 March 2018.
- Singleton, I., Wich, S.A., Nowak, M., Usher, G. and Utami-Atmoko, S.S. (2017). *Pongo abelii*. *The IUCN Red List of Threatened Species 2017*: e.T121097935A115575085. <http://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T121097935A115575085.en>. Downloaded on 24 March 2018.
- Sloan, S., Bertzky, B. and Laurance, W.F. (2016). 'African development corridors intersect key protected areas'. *African Journal of Ecology* 55: 731–737.
- Spira, C., Kirkby, A., Kujirakwinja, D. and Plumptre, A. (2017). 'The socio-economics of artisanal mining and bushmeat hunting around protected areas: Kahuzi-Biega National Park and Itombwe Nature reserve, eastern Democratic Republic of Congo'. *Oryx* 1–9.
- Steel, L., Colom, A., Maisels, F. and Shapiro, A. (2008). The scale and dynamics of wildlife trade originating in the south of Salonga-Lukenie-Sankuru landscape. WWF-WCS.
- Stiles, D., Redmond, I., Cress, D., Nellemann, C. and Formo, R.K. (2013). *Stolen Apes: The Illicit Trade in Chimpanzees, Gorillas, Bonobos, and Orangutans. A Rapid Response Assessment*. Nairobi, Kenya: United Nations Environment Programme, GRID-Arendal.
- Strindberg, S., Maisels, F., Williamson, E.A., Blake, S., Stokes, E.J., Aba'a, R., Abitsi, G., Agbor, A., Ambahe, R.D., Bakabana, P.C., Bechem, M., Berlemont, A., de Semboli, B.B., Boundja, P.R., Bout, N., Breuer, T., Campbell, G., de Wachter, P., Akou, M.E., Mba, F.E., Feistner, A.T.C., Fosso, B., Fotso, R., Greer, D., Inkamba-Nkulu, C., Iyenguet, C.F., Jeffery, K.J., Kokangoye, M., Köhl, H.S., Latour, S., Madzoke, B., Makoumbou, C., Malanda, G.A.F., Malonga, R., Mbolu, V., Morgan, D.B., Motsaba, P., Moukala, G., Mowawa, B.S., Murai, M., Ndzai, C., Nishihara, T., Nzoo, Z., Pintea, L., Pokempner, A., Rainey, H.J., Rayden, T., Ruffler, H., Sanz, C.M., Todd, A., Vanleeuwe, H., Vosper, A., Warren, Y. and Wilkie, D.S. (2018). 'Guns, germs, and trees determine density and distribution of gorillas and chimpanzees in Western Equatorial Africa'. *Science Advances* 4: eaar2964.
- Struebig, M.J., Fischer, M., Gaveau, D.L.A., Meijaard, E., Wich, S.A., Gonner, C., Sykes, R., Wilting, A. and Kramer-Schadt, S. (2015). 'Anticipated climate and land-cover changes reveal refuge areas for Borneo's orangutans'. *Global Change Biology* 21: 2891–2904.
- Tempo (2016). 'Three Sentenced to Prison over Orangutan Illegal Trade.' [Online]. (23rd March 2016). <https://en.tempo.co/read/news/2016/03/23/206756195/Three-Sentenced-to-Prison-over-Orangutan-Illegal-Trade>
- Thomas, L., Buckland, S.T., Rexstad, E.A., Laake, J.L., Strindberg, S., Hedley, S.L., Bishop, J.R.B., Marques, T.A. and Burnham, K.P. (2010). 'Distance software: design and analysis of distance sampling surveys for estimating population size'. *Journal of Applied Ecology* 47: 5–14.
- Tiédoúé, M. R., Normand, E., Diarrassouba, A., Tondossama, A. and Boesch, C. (2016). *Etat de conservation du Parc National de Taï: Rapport de suivi-écologique - phase 11. Rapport de l'Office Ivoirien des Parcs et Réserves/Direction de Zone Sud-ouest, Soubre, Côte d'Ivoire, 31 pp.*
- Transparency International. (2018). 'Corruption Perceptions Index 2017'. [Online] [www.transparency.org/news/feature/corruption](http://www.transparency.org/news/feature/corruption)
- Tweh, C.G., Lormie, M.M., Kouakou, C.Y., Hillers, A., Köhl, H.S. and Junker, J. (2015). 'Conservation status of chimpanzees *Pan troglodytes verus* and other large mammals in Liberia: A nationwide survey'. *Oryx* 49: 710–718.
- United Nations Office on Drugs and Crime, World Drug Report (2016). United Nations.
- USAID (2015). *Measuring efforts to combat wildlife crime: A toolkit for improving action and accountability*. United States Agency for International Development.
- Van der Meer, I.P. (2014). *Inventory of the vegetation structure and food availability for the Western chimpanzee (Pan troglodytes verus) in the Boé region, Guinea-Bissau*. Internship report. Ecology Group of Wageningen University and Chimbo Foundation.

Van der Meer, D. (2016). Can the Random Encounter Model be a useful tool in estimating chimpanzee density? Internship report. VHL University of Applied Sciences and Chimbo Foundation.

Voigt, M., Wich, S., Ancrenaz, M., Meijaard, E., Abram, N., Banes, G., Campbell-Smith, G., d’Arcy, L.J., Delgado, R.A., Erman, A., Gaveau, D., Goossens, B., Heinicke, S., Houghton, M., Husson, S.J., Leiman, A., Sanchez, K.L., Makenuddin, N., Marshall, A., Meididit, A., Miettinen, J., Mundry, R., Musnanda, Nardiyono, Nurcahyo, A., Russon, A., Santika, T., Sihite, J., Spehar, S., Struebig, M., Sulbaran-Romero, E., Tjiu, A., Wells, J., Wilson, K. and Köhl, H. (2018). ‘Global demand for natural resources eliminated more than 100,000 Bornean orangutans’. *Current Biology* 28: 761–769.

Walsh, P., Abernethy, K., Bermejo, M., Beyers, R., De Wachter, P., Akou, M., Huijbregts, B., Mambounga, D., Toham, A., Kilbourn, A., Lahm, S., Latour, S., Maisels, F., Mbina, C., Mihindou, Y., Ndong Obiang, S., Effa, E., Starkey, M., Telfer, P., Thibault, M., Tutin, C., White, L. and Wilkie, D. (2003). ‘Catastrophic ape decline in western equatorial Africa’. *Nature* 422(6932): 611–614.

WCF (2012). Etat de la faune et des menaces dans les aires protégées terrestres et principales zones de forte biodiversité de République de Guinée. Unpublished report, Wild Chimpanzee Foundation, Conakry, Guinea.

WCF (2014). Study of a key area for the preservation of chimpanzee in West Africa: Preliminary inventory along the Bafing River, Republic of Guinea. Unpublished report.

Wenceslau, J.F.C. (2014). ‘Bauxite mining and chimpanzees population distribution, a case study in the Boé sector, Guinea-Bissau’. Master Thesis. VHL University of Applied Sciences and Chimbo Foundation.

Wicander, S. and Coad, L. (2015). *Learning our lessons: A review of alternative livelihood projects in Central Africa*. ECI, University of Oxford, Oxford, UK and IUCN, Gland, Switzerland.

Wich, S.A., Utami-Atmoko, S.S., Setia, T.M., Rijksen, H.D., Schürmann, C., van Hooff, J.A.R.A.M. and van Schaik, C.P. (2004). ‘Life history of wild Sumatran orangutans (*Pongo abelii*)’. *Journal of Human Evolution* 47: 385–398.

Wich, S.A., Meijaard, E., Marshall, A.J., Husson, S., Ancrenaz, M., Lacy, R.C., van Schaik, C.P., Sugardjito, J., Simorangkir, T., Traylor-Holzer, K., Doughty, M., Supriatna, J., Dennis, R., Gumal, M., Knott, C.D. and Singleton, I. (2008). ‘Distribution and conservation status of the orangutan (*Pongo* spp.) on Borneo and Sumatra: how many remain?’. *Oryx* 42: 329–339.

Wich, S.A., Swartz, K.B., Hardus, M.E., Lameira, A.R., Stromberg, E. and Shumaker, R.W. (2009). ‘A case of spontaneous acquisition of a human sound by an orangutan’. *Primates* 50: 56–64.

Wich, S., Riswan, J., Refisch, J. and Nellemann, C. (2011). *Orangutans and the economics of sustainable forest management in Sumatra*. UNEP/GRASP/PanEco/YEL/ICRAF/Grid Arendal.

Wich, S., Fredriksson, G.M., Usher, G., Peters, H.H., Priatna, D., Basalamah, F., Susanto, W. and Köhl, H. (2012a). ‘Hunting of Sumatran orangutans and its importance in determining distribution and density’. *Biological Conservation* 146: 163–169.

Wich S.A., Gaveau, D., Abram, N., Ancrenaz, M., Baccini, A., Brend, S., Curran, L., Delgado, R.A., Erman, A., Fredriksson, G.M., Goossens, B., Husson, S.J., Lackman, I., Marshall, A.J., Naomi, A., Molidena, E., Nardiyono, Nurcahyo, A., Odom, K., Panda, A., Purnomo, Rafiastanto, A., Ratnasari, D., Santana, A.H., Sapari, I., van Schaik, C.P., Sihite, J., Spehar, S., Santoso, E., Suyoko, A., Tiju, A., Usher, G., Atmoko, S.S.U., Willems, E.P. and Meijaard, E. (2012b). ‘Understanding the impacts of land-use policies on a threatened species: is there a future for the Bornean orangutan?’. *PLoS One* 7: e49142.

Wich, S.A., Garcia-Ulloa, J., Köhl, H.S., Humle, T., Lee, J.S. and Koh, L.P. (2014). ‘Will oil palm’s homecoming spell doom for Africa’s great apes?’. *Current Biology* 24: 1659–1663.

Wich, S.A., Struebig, M., Refisch, J., Wilting, A., Kramer-Schadt, S. and Meijaard, E. (2015). The future of the Bornean orangutan: Impacts of change in land cover and climate. Nairobi, Kenya: UNEP/GRASP.

Wich, S., Singleton, I., Nowak, M., Utami Atmoko, S., Nisam, G., Arif, S., Putra, R., Ardi, R., Fredriksson, G., Usher, G., Gaveau, D. and Köhl, H. (2016). ‘Land-cover changes predict steep declines for the Sumatran orangutan (*Pongo abelii*)’. *Science Advances* 2: e1500789.

Wilkie, D.S., Bennett, E.L., Peres, C.A. and Cunningham, A.A. (2011). 'The empty forest revisited'. *Annals of the New York Academy of Sciences* 1223: 120–128.

Wilkie, D.S., Wieland, M., Boulet, H., Le Bel, S., van Vliet, N., Cornelis, D., BriacWarnon, V., Nasi, R. and Fa, J.E. (2016). 'Eating and conserving bushmeat in Africa'. *African Journal of Ecology* 54: 402–414.

Wood, K.L., Tenger, B., Morf, N.V. and Kratzer, A. (2014). Report to CITES: Cites-listed species at risk from illegal trafficking in bushmeat; Results of a 2012 study in Switzerland's international airports. 129 pp. Tengwood Organization & Zürich Institute of Forensic Medicine, University of Zürich, Forensic Genetics, Switzerland.

World Bank (2018). World Development Indicators. [Online] <http://data.worldbank.org/data-catalog/world-development-indicators>

WWF and TRAFFIC (2015). Strategies for fighting corruption in wildlife conservation: A primer. WWF and TRAFFIC Wildlife Crime Initiative. 23 pp. [Online] [http://d2ouvy59p0dg6k.cloudfront.net/downloads/wci\\_strategies\\_for\\_fighting\\_corruption\\_wildlife\\_conservation.pdf](http://d2ouvy59p0dg6k.cloudfront.net/downloads/wci_strategies_for_fighting_corruption_wildlife_conservation.pdf)

Wyatt, T. and Ngoc Cao, A. (2015). Corruption and wildlife trafficking. U4 Anti-Corruption Resource Centre. U4 Issue May 2015, No 11. 54 pp. [Online] <http://issuu.com/cmi-norway/docs/150603144854-73fa106fbd71418f9a0afa481e93443f/1?e=0>

Ziegler, S., Fa, J.E., Wohlfart, C., Streit, B., Jacob, S. and Wegmann, M. (2016). 'Mapping bushmeat hunting pressure in Central Africa'. *Biotropica* 48: 405–412.

#### Annex I. Acronyms

A.P.E.S. database	Ape Populations, Environments and Surveys database
CITES CoP	CITES Conference of the Parties
CMS	Convention on the Conservation of Migratory Species of Wild Animals
DNA	Deoxyribonucleic acid
GRASP	Great Apes Survival Partnership
ICCN	Institut Congolais pour la Conservation de la Nature
IGO	Intergovernmental organisation
INTERPOL	International Criminal Police Organisation
IUCN	International Union for Conservation of Nature
JGI	Jane Goodall Institute
NGO	Non-governmental organisation
LAGA	The Last Great Ape Organisation Cameroon
PanAf	The Pan African Programme
PSG	Primate Specialist Group
SGA	Section on Great Apes
SSC	Species Survival Commission
TRAFFIC	The Wildlife Trade Monitoring Network
UNEP	United Nations Environment Programme
USAID	United States Agency for International Development